



## Literature review

**PhD candidate:** François Leroy

**Programme:** Environmental Earth Sciences

**Department:** Spatial Sciences

*"Mapping biodiversity changes across  
spatio-temporal scales"*

**Advisor:** doc. Ing. Petra Šímová, Ph.D.

**Consultant:** Mgr. Petr Keil, PhD

**Beginning of study:** October 2020

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Materials and Methods</b>	<b>3</b>
<b>3</b>	<b>On the link between spatial grain and trends of biodiversity</b>	<b>9</b>
<b>4</b>	<b>A lack of consistency</b>	<b>13</b>
4.1	Spatial replicates is not a standard . . . . .	13
4.2	Metric heterogeneity . . . . .	14
4.3	On the temporal scale . . . . .	14
<b>5</b>	<b>Future directions</b>	<b>15</b>
<b>6</b>	<b>Conclusion</b>	<b>16</b>
	<b>References</b>	<b>17</b>
	<b>Supplementary materials</b>	<b>22</b>

# 1. Introduction

Human life quality is intrinsically linked to ecosystems state that he is living in. Indeed, ecosystems services extend in a large spectrum of mechanisms including nutrient cycle, food production, or climate and water cycle regulation (Pereira et al., 2012). Some of those ecosystem functions are managed by bird biodiversity such as seed dispersal, controls pests or pollinate plant. Unfortunately, anthropogenic stressors like habitat loss, over exploitation, pollution or introduction of invasive species could lead biodiversity to its sixth mass extinction (Barnosky et al., 2011).

We now know that the loss of global biodiversity is unprecedented and political decisions has been stated in order to limit it (*e.g.* of the Convention on Biological Diversity & Centre, 2006, pp. 2010, 2002). However, current scientific literature has also shown that temporal trends in local changes of biodiversity can be opposite to trends at larger scales (*e.g.* Chase et al., 2019). Thus, current changes in biodiversity is far more complex than a simple global decrease: most of the ecosystems undergo alterations of their communities with changes in species composition (Blowes et al., 2019; Dornelas et al., 2013). Wonders persist about how the trend of these different metrics of biodiversity are link to the spatial and temporal scales used when measured.

In order to investigate this link between spatio-temporal scales and biodiversity metrics, birds is a relevant taxon. Thanks to the many ornithological monitoring and surveys, we now have a large number of long, high-quality time series on bird populations (Bejček & Stastný, 2016; Kamp et al., 2021; Sauer et al., 2013). Birds are easy to observe, easy to identify and thus many volunteers are motivated to conduct standardized sampling. Given their ability to change quickly of locations, their presence is also a good indicator for ecosystem health and thus several standardized metrics have been created to assess their populations.

However, studying biodiversity can be confusing, especially because several choices must be done. Firstly, the level at which you are looking at the biodiversity must be chosen (*e.g.* taxonomic, functional, phylogenetic diversity). Secondly, one must decide which metric is the most appropriate for his study. There are many facets of biodiversity that can be measured by different metrics depending on the objective of your study. Measures of static biodiversity are commonly used such as species richness or  $\alpha$  diversity (*i.e.* number of species, Whittaker, 1960), the Shannon index (Shannon, 1948), the Simpson

index (Simpson, 1949) or the Hill number (Hill, 1973). The later three biodiversity indexes take into account the relative abundances of the species and can be considered as the *quality* of the biodiversity. On an other hand, the spatial and temporal  $\beta$  diversity will measure the species turnover and can be measured thanks to Whittaker's (Whittaker, 1972), Sørensen's (Sørensen, 1948) or Jaccard's (Jaccard, 1912) dissimilarity indexes (*e.g.* Keil et al., 2012). All these metrics assess the taxonomic diversity, *i.e.* they use the species as unit. However, it has also been shown that functional and phylogenetic diversity can provide supplementary information on the community structure and its dynamic (*e.g.* McGill et al., 2006; Mouquet et al., 2012; Webb et al., 2002).

An other measure of biodiversity of great interest is the abundance of the populations. As a matter of fact, as individuals react to disturbances of ecosystems by disappearing, the population trends are considered as resulting of the ecosystems health. Overall populations are impossible to assess, but determining the abundance of few indicator species can give a relevant summary on how is going the entire ecosystem (Gregory et al., 2005). These family of metrics are called the multi-species indicators (MSI) and we have seen the emergence of several of them according to the ecosystem considered. We can cite here the farmland bird indicator, woodland bird indicator or one of the most informative, which summarizes these two previous metrics: the Wildland Bird Indicator (Gregory et al., 1999; Gregory & Strien, 2010). The main idea of these metrics is to compute the geometric mean of abundance of few key species over time.

Finally a last class of indicators, not considered here, take into account both species and ecosystems feature into a summarizing index. The most known ones are the Red List Index (S. Butchart et al., 2005; S. H. M. Butchart et al., 2007; S. H. M. Butchart et al., 2004) or the Biodiversity Change Index (Normander et al., 2012).

Here, I propose to review articles assessing the temporal trends of different macroecological avian biodiversity metrics and to look at which spatial and temporal scales these studies have been done. I decided to consider the most common indicators used to assess biodiversity at the community level and higher, such as the diversity indexes (*e.g.* species richness, functional diversity...) or the abundance indexes (McGill et al., 2015). Summarizing the trends of these qualitative and/or quantitative avian biodiversity indexes along with their spatial and temporal scales will help to see more clearly how the trends of biodiversity are linked to spatio-temporal scales. It is also important to demonstrate that the information about the sampling plan (*i.e.* spatial scale, time span, temporal scales etc) is not systematically indicated in the scientific literature and can bring confusion to the analysis and comparisons of their

trends. I believe that this review can help to have a better overview of the current knowledge on the link between trends of biodiversity metrics of bird populations and the spatio-temporal scales.

specify that it is mainly for continental/coastal birds but no look at islandic communities and change  
\*Stable\* to \*No trend\*

## 2. Materials and Methods

For this review, articles of interest were the ones assessing temporal trends of the most common indicators (*i.e.* metrics) of avian biodiversity and specifying spatial and temporal scales. For this, I used the “*advanced search*” tool of the ISI Web of Science Core collection database with these four following queries:

1. AB = ((biodiversity OR species richness OR diversity) AND (temporal trend\* OR dynamic\*) AND (bird\* OR avia\*)) which resulted in 1346 references.
2. AB = ((biodiversity change index) AND (bird\* OR avia\*) AND trend\*) which resulted in 60 references.
3. AB = ((species richness) AND (bird\* OR avia\*) AND trend\*) which resulted in 313 references.
4. ALL=(birds AND species richness AND temporal trend) which resulted in 88 references.

For each query, the title and abstract of the articles were reviewed. When the temporal trend was explicitly specified (either visually or literally), the material and method part was read in order to collect the *spatial grain* of the trend (*i.e.* the area at which the trend is assessed), its *temporal grain* (*i.e.* the time span at which data have been gathered at one census session), the *spatial extent* (*i.e.* the entire area at which the study applies), the *temporal extent* and the *beginning and ending years* of the study as well as the *general trend* of the metric (Table 2.1). Alternatively, the articles which were often referred to in the relevant articles were also explored.

I decided to take into account only the articles for which there was spatial replicates, *i.e.* where the trend of the metric was assessed at several locations with the same spatial grain. With these replications, the trend reported at one spatial grain is more reliable. However, at larger spatial grains (*i.e.* national, continental or global scales), spatial replicates are rarer. Thus, I decided to take note of these unreplicated

trends only when they were found in a article were spatial replicates were used at smaller scales.

Spatial grain sizes are discretized into four levels: *local*  $\leq 25 \text{ Km}^2$ , *regional*  $> 25 \text{ Km}^2$ , *national* when an entire country is considered and *global* at the worldwide scale.

Concerning the trend assessment, different papers contain the *p-value* or directly specify the significant trend of the metric. However, a portion of papers gives only visual representations of the trend. For those, the standard error was used when displayed. For the very few only giving the trend, **the rule of thumb was applied**. Information can be found in the column *Note* of the Table 6.1 of the supplementary material. Moreover, the final trend retained (*i.e.* either *Increase*, *Stable* or *Decrease*) doesn't reflect all the fluctuations of the metric through time but rather the difference between the starting and ending points.

I have found 31 references in which authors were both determining the temporal trend of a metric of interest and explicitly defining the grain size. However, only 17 of them are using spatial replicates and are thus relevant for this study (Table 2.1). The classes of metric are: *Species richness*, *Evenness*, *Abundance*, *Diversity*, *Temporal beta-diversity*, *Spatial beta-diversity*, *Functional diversity*, *Functional evenness*, *Functional richness*. Some of this classes contain several different indexes. This is the case for instance for the class *Diversity*, which can be defined by either the Shannon or the Simpson index, or for the class *Abundance* which contains various multi-species indicators (for more information, see Table 6.1).

Table 2.1: Trends of different metrics of biodiversity at various spatial and temporal scales

Reference	Metric	Spatial grain (Km <sup>2</sup> )	Temporal grain (year)	Spatial extent (Km <sup>2</sup> )	Temporal extent (year)	Years	Country	Trend
Barnagaud et al., <a href="#">2017</a>	Evenness	Local	1.0	9834000	41	1970-2011	USA	Increase
	SR	Local	1.0	9834000	41	1970-2011	USA	Increase
	Abundance	National	1.0	520475	27	1990-2016	Czech Rep., Switzerland, Denmark, Germany	Stable
	SR	Local	5.0	2800000	30	1982–2011	USA, Canada	Stable
	SR	Regional	5.0	2800000	30	1982–2011	USA, Canada	Stable
	SR	Regional	5.0	2800000	30	1982–2011	USA, Canada	Stable
	SR	Local	5.0	2800000	30	1982–2011	USA, Canada	Increase
	SR	Local	5.0	2800000	30	1982–2011	USA, Canada	Increase
	Abundance	Regional	1.0	643801	14	2007-2020	France	Decrease
	Abundance	Regional	1.0	643801	14	2007-2020	France	Decrease
Chiron et al., <a href="#">2013</a>	Abundance	Regional	1.0	643801	14	2007-2020	France	Decrease
	Abundance	Regional	1.0	643801	14	2007-2020	France	Decrease
	Abundance	Regional	1.0	643801	14	2007-2020	France	Decrease
	Diversity	Local	1.0	242495	13	1994-2006	UK	Increase
Davey et al., <a href="#">2012</a>	Evenness	Local	1.0	242495	13	1994-2006	UK	Increase
	SR	Local	1.0	242495	13	1994-2006	UK	Increase

Table 2.1: Trends of different metrics of biodiversity at various spatial and temporal scales (*continued*)

Reference	Metric	Spatial grain (Km <sup>2</sup> )	Temporal grain (year)	Spatial extent (Km <sup>2</sup> )	Temporal extent (year)	Years	Country	Trend
Harrison et al., 2014	Abundance	Local	1.0	200000	18	1994-2011	Great Britain, UK	Increase
	Abundance	Local	1.0	200000	18	1994-2011	Great Britain, UK	Stable
	Abundance	Local	1.0	200000	18	1994-2011	Great Britain, UK	Stable
Harrison et al., 2016	Abundance	Local	0.5	242495	20	1994-2013	UK	Increase
	Abundance	Local	0.5	242495	20	1994-2013	UK	Stable
Jarzyna and Jetz, 2018	Abundance	Local	0.5	242495	20	1994-2013	UK	Stable
	SR	Regional	1.0	9834000	45	1969-2013	USA	Increase
	SR	Regional	1.0	9834000	45	1969-2013	USA	Increase
	SR	Regional	1.0	9834000	45	1969-2013	USA	Increase
	SR	National	1.0	9834000	45	1969-2013	USA	Increase
	SR	Global	1.0	148940000	45	1969-2013	World	Decrease
	Temporal beta-diversity	Regional	1.0	9834000	45	1969-2013	USA	Increase
	Temporal beta-diversity	Regional	1.0	9834000	45	1969-2013	USA	Increase
	Temporal beta-diversity	Regional	1.0	9834000	45	1969-2013	USA	Increase



Table 2.1: Trends of different metrics of biodiversity at various spatial and temporal scales (*continued*)

Reference	Metric	Spatial grain (Km <sup>2</sup> )	Temporal grain (year)	Spatial extent (Km <sup>2</sup> )	Temporal extent (year)	Years	Country	Trend
	Temporal beta-diversity	National	1.0	9834000	45	1969-2013	USA	Increase
	Temporal beta-diversity	Global	1.0	148940000	45	1969-2013	World	Stable
Pilotto et al., <a href="#">2020</a>	Abundance	Local	NA	10180000	NA	NA	Europe	Stable
	Diversity	Local	NA	10180000	NA	NA	Europe	Increase
	SR	Local	NA	10180000	NA	NA	Europe	Increase
	Temporal beta-diversity	Local	NA	10180000	NA	NA	Europe	Stable
Ram et al., <a href="#">2017</a>	Abundance	Local	1.0	350000	18	1998-2015	Sweden	Increase
	SR	Regional	1.0	350000	18	1998-2015	Sweden	Increase
Reif et al., <a href="#">2013</a>	Spatial beta-diversity	Local	1.0	79000	23	1982-2004	Czech Rep.	Stable
	SR	Local	1.0	79000	23	1982-2004	Czech Rep.	Stable
	SR	National	1.0	79000	23	1982-2004	Czech Rep.	Stable
Schipper et al., <a href="#">2016</a>	Abundance	Local	5.0	24710000	40	1971-2010	Canada, USA, Mexico	Increase
	Diversity	Local	5.0	24710000	40	1971-2010	Canada, USA, Mexico	Increase
	Diversity	Local	5.0	24710000	40	1971-2010	Canada, USA, Mexico	Increase

Table 2.1: Trends of different metrics of biodiversity at various spatial and temporal scales (*continued*)

Reference	Metric	Spatial grain (Km <sup>2</sup> )	Temporal grain (year)	Spatial extent (Km <sup>2</sup> )	Temporal extent (year)	Years	Country	Trend
∞	Functional diversity	Local	5.0	24710000	40	1971-2010	Canada, USA, Mexico	Decrease
	Functional evenness	Local	5.0	24710000	40	1971-2010	Canada, USA, Mexico	Increase
	Functional richness	Local	5.0	24710000	40	1971-2010	Canada, USA, Mexico	Increase
	SR	Local	5.0	24710000	40	1971-2010	Canada, USA, Mexico	Increase
	Sorte and Boecklen, 2005	Local	1.0	9834000	36	1968-2003	USA	Decrease
	Evenness	Local	1.0	9834000	36	-	USA	Decrease
	SR	Local	1.0	9834000	36	1968-2003	USA	Increase
	Van Turnhout et al., 2007	Regional	4.0	41543	28	1973-2000	Netherlands	Increase
	SR	Local	4.0	41543	28	1973-2000	Netherlands	Increase
	SR	National	4.0	41543	28	1973-2000	Netherlands	Increase
Wretenberg et al., 2010	SR	Local	1.0	1800	11	1994-2004	Sweden	Decrease
Inger et al., 2015	Abundance	National	1.0	10180000	22	1980-2001	Europe	Decrease
Donald et al., 2001	Abundance	National	NA	10180000	21	1970-1990	Europe	Decrease

### 3. On the link between spatial grain and trends of biodiversity

The median spatial extent of the 17 articles is  $5.20475 \times 10^5 \text{ Km}^2$ , with the smallest area of  $1800 \text{ Km}^2$  and the greatest representing the global emerged surface (*i.e.*  $1.4894 \times 10^8 \text{ Km}^2$ ). Overall, there were 11 *Decrease*, 31 *Increase* and 14 *Stable* reliable trends (*i.e.* spatially replicated) across the literature, without consideration of any metric or spatial scales.

One have to keep in mind that the proportions of the trends presented here is not representative of the overall the scientific literature. Taking only the articles with spatial replicates reduces the number of studies and thus doesn't reflect the overall dynamic of biodiversity, which is not the purpose of this review. As a matter of fact, meta-analysis by Fraixedas et al. (2020) indicates that around half of bird related indicators are declining and that only 20% of them are increasing.

In our case, local scales are more represented than the others and the number of papers decreases with the increasing spatial scale (Figure 3.1). This is expected, as the spatial replications get more demanding in organization and resources as the grain size enlarges. The *Increase* of the metrics seems to be dominating at smaller scales. On an other hand, the proportion of *Decrease* is more important at regional scales than at local scales. At the global scale, no *Increase* were found.

Replicated species richness and abundance metrics are very represented in the scientific literature (Figure 3.2). The less common trend of abundance is *Increase*, whilst *Decrease* and *Stability* are both as common. Diversity indexes (*i.e.* Sørensen and Jaccard) were always found increasing and temporal  $\beta$ -diversity was found most of the time increasing and never decreasing. The small frequency of the other metrics doesn't allow one to discuss the trend proportions.

**At local scales**, species richness experiences dominantly increase (Figure 3.3). Evenness indexes, *i.e.* taxonomic and functional evenness, are mainly found increasing. Concerning the abundance indexes, the stability dominates but the increase is also often witnessed. In contrast, **at regional scale**, abundance metrics always decrease, whilst temporal  $\beta$ -diversity always increase. Species richness experiments mainly increases, then stability but no decrease is reported. Concerning **national and global scales**, one have to keep in mind that spatial replicates are less frequent and that most of the trends reported here




Figure 3.1: Proportion of *Increase*, *Decrease* or *Stable* trends for each spatial scale




Figure 3.2: Proportion of *Increase*, *Decrease* or *Stable* trends for each of the metric

for these two spatial scales are not replicated. Exceptions are made by Bowler et al. (2021) and Donald et al. (2001). The former showed negative trends in abundance indexes for Denmark and Germany and positive trends for Switzerland and Czech Republic, indicating that one can not conclude about general trend at national scale (here referred as *Stable*). For Donald et al. (2001), trends of mean populations were computed for 30 european countries and most of them were negatives. No positive trend seem dominant at these spatial scales. Global trends only come from Jarzyna and Jetz (2018).




Figure 3.3: Proportion of *Increase*, *Decrease* or *Stable* trends for each metric. Each panel represent one spatial scale

Number of studies using spatial replicates is limited. Thus, observing general rules about spatial scaling of trends of macroecological metrics is not possible with the current state of the scientific literature. However, concerning species richness, we can see that the decrease is global, but that this decrease is rare at lower spatial scales. In fact, increases are more often observed, confirming the high perturbations that biodiversity is undergoing (Dornelas et al., 2014; Vaidyanathan, 2021). This analysis goes along with the temporal  $\beta$ -diversity which is always observed either stable or increasing, sign of instability of the ecosystems.

On an other hand, the decrease of abundances seem to appear mainly at regional or national scales. This may be due to the fact that local scale samplings are done usually outside of very anthropized ecosystems, such as cities. Increasing the scale to regional and national increases the proportion of cities and farmland area, which can have negative impact on bird populations (Reif, 2013).

## 4. A lack of consistency

### 4.1 Spatial replicates is not a standard

Articles computing the trend of a metric with spatial replicates were limited, either due to a lack of data or because the trend was assessed for the spatial extent of the data. For instance, the Breeding Bird Surveys (*e.g.* Kamp et al., 2021; Sauer et al., 2013) follow standardized sampling plan with spatial replications (*i.e.* census plots). However, not all the trend reported for the BBS are summarized at their specific grain sizes and were sometimes computed at their respective national scales, therefore without spatial replication. For instance, the common method encountered to assess population trends (*i.e.* abundance indexes) is to learn a predictive model from the data, predict the target feature (*i.e.* abundance) and then compute the metric and its trend from the output of the model at the national spatial extent (*e.g.* Doxa et al., 2010; Eglington & Pearce-Higgins, 2012; Jiguet et al., 2005; Jiguet et al., 2012; Sauer et al., 2017). These analysis are essential from a conservation point of view and are a majority (Fraixedas et al., 2020): they give valuable information about ecosystem health at large spatial extent and are thus used as arguments by decision-makers. However, they represent just one trend amongst all the possible trends that can be found at this spatial scales and thus doesn't give information about the link between spatial grain and temporal trend.

Even less articles were computing the trends of metrics with spatial replicates at various spatial grain. This was the case for only Chase et al. (2019) and Jarzyna and Jetz (2018). It is also important to cite here Jarzyna et al. (2015), who did iteration computing of temporal change community metrics (*i.e.* temporal dissimilarity, temporal turnover, extinction and colonization) at several spatial scales. However, the temporal trend of these metrics weren't considered and are therefore not reported in Table 2.1.

## 4.2 Metric heterogeneity

Macroecology has seen its metrics blooming these last decades, in search of a multiparametric indexes taking into account all the components of an ecosystem, as do the composite indicators (*e.g.* the red list index, S. H. M. Butchart et al., 2007; biodiversity change indexes, Normander et al., 2012). The latter have proven very effective in order to assess conservation policies and are able to synthesize the ecosystem complexities. This has been especially the case for the birds, who are of great interest in assessing the dynamic of ecosystem health (Fraixedas et al., 2020). Thus, studies focusing only on macroecological metrics are less common than expected. Coupled with the need of spatial replicates, studying the link between the trend of macroecological metrics and spatio-temporal scales still needs to be explored.

## 4.3 On the temporal scale

The oldest study started in 1968 and the median duration is 23 years, with a minimum time span of 11 years and a maximum of 45.

Determination of the temporal grain was way more complicated, as there is no consensus on the subject. Usually, the temporal grain of the sampling is specified, but sometimes with inaccuracies (*e.g.* “in the early morning”). However, the temporal grain of the sampling plan doesn’t represent the final temporal grain. For instance, some metrics are summed over a certain spatial extent (*e.g.* summing the species richness over an atlas square, like in Van Turnhout et al., 2007). In this case, the temporal grain should also be summed, but this was not specified. On an other hand, when the metric is averaged over a spatial grain, the sampling temporal grain can be considered.

Nevertheless, an other problem arises when the trend is computed. Usually, one compute the metric out of the data and considers the value to be representative of half a year or a year. In this case, there is no consensus on what should be the temporal grain: should one keep the grain of the sampling? Should one consider it to be the time-span covered by the sampling? In my analysis, the latter has been used (Table 2.1).

For the cases where the metric was determined out of model (*e.g.* Harrison et al., 2014), the final temporal grain was most of the time explicitly given, with an order of length of the year.



Figure 1 is a log-log plot showing the relationship between Temporal grain (year) on the y-axis and Spatial extent (Km²) on the x-axis. The y-axis ranges from 1 to 5, and the x-axis ranges from 1e+03 to 1e+08. A blue line represents a positive linear relationship on the log-log scale. Data points are black dots. Most points follow the trend, but there are outliers at higher temporal grain values (4 and 5 years) for intermediate spatial extents.

Spatial extent (Km²)	Temporal grain (year)
1e+03	1
1e+04	1.3
3e+04	4
8e+04	1
2e+05	1
3e+05	0.5
4e+05	1
6e+05	1
8e+05	1
3e+06	5
1e+07	1
3e+07	5
1e+08	1

## 5. Future directions

15

## 6. Conclusion

As birds represent one of the most data furnished taxa of vertebrates, reviewing the scientific literature on this topic gives a glimpse of what still needs to be done to better understand the scaling of global biodiversity dynamic.

# References

- Barnagaud, J.-Y., Gaüzère, P., Zuckerberg, B., Princé, K., & Svenning, J.-C. (2017). Temporal changes in bird functional diversity across the United States. *Oecologia*, 185(4), 737–748. <https://doi.org/10.1007/s00442-017-3967-4>
- Barnosky, A. D., Matzke, N., Tomiya, S., Wogan, G. O. U., Swartz, B., Quental, T. B., Marshall, C., McGuire, J. L., Lindsey, E. L., Maguire, K. C., Mersey, B., & Ferrer, E. A. (2011). Has the Earth's sixth mass extinction already arrived? [Number: 7336 Publisher: Nature Publishing Group]. *Nature*, 471(7336), 51–57. <https://doi.org/10.1038/nature09678>
- Bejček, V., & Stastný. (2016). Velké ptačí mapování. *Vesmír*. Retrieved December 10, 2020, from <https://vesmir.cz/cz/on-line-clanky/2016/04/velke-ptaci-mapovani.html>
- Blowes, S. A., Supp, S. R., Antão, L. H., Bates, A., Bruelheide, H., Chase, J. M., Moyes, F., Magurran, A., McGill, B., Myers-Smith, I. H., Winter, M., Bjorkman, A. D., Bowler, D. E., Byrnes, J. E. K., Gonzalez, A., Hines, J., Isbell, F., Jones, H. P., Navarro, L. M., ... Dornelas, M. (2019). The geography of biodiversity change in marine and terrestrial assemblages [Publisher: American Association for the Advancement of Science Section: Research Article]. *Science*, 366(6463), 339–345. <https://doi.org/10.1126/science.aaw1620>
- Bowler, D., Richter, R. L., Eskildsen, D., Kamp, J., Moshøj, C. M., Reif, J., Strebel, N., Trautmann, S., & Voříšek, P. (2021). Geographic variation in the population trends of common breeding birds across central Europe. *Basic and Applied Ecology*, 56, 72–84. <https://doi.org/10.1016/j.baae.2021.07.004>
- Butchart, S., Stattersfield, A., Baillie, J., Bennun, L., Stuart, S., Akçakaya, H., Hilton-Taylor, C., & Mace, G. (2005). Using Red List Indices to measure progress towards the 2010 target and beyond [Publisher: Royal Society]. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 360(1454), 255–268. <https://doi.org/10.1098/rstb.2004.1583>
- Butchart, S. H. M., Akçakaya, H. R., Chanson, J., Baillie, J. E. M., Collen, B., Quader, S., Turner, W. R., Amin, R., Stuart, S. N., & Hilton-Taylor, C. (2007). Improvements to the Red List Index [Publisher: Public Library of Science]. *PLOS ONE*, 2(1), e140. <https://doi.org/10.1371/journal.pone.0000140>
- Butchart, S. H. M., Stattersfield, A. J., Bennun, L. A., Shutes, S. M., Akçakaya, H. R., Baillie, J. E. M., Stuart, S. N., Hilton-Taylor, C., & Mace, G. M. (2004). Measuring Global Trends in the Status of Biodiversity: Red List Indices for Birds (Walt V. Reid, Ed.). *PLoS Biology*, 2(12), e383. <https://doi.org/10.1371/journal.pbio.0020383>
- Chase, J. M., McGill, B. J., Thompson, P. L., Antão, L. H., Bates, A. E., Blowes, S. A., Dornelas, M., Gonzalez, A., Magurran, A. E., Supp, S. R., Winter, M., Bjorkman, A. D., Bruelheide, H., Byrnes, J. E. K., Cabral, J. S., Elahi, R., Gomez, C., Guzman, H. M., Isbell, F., ... O'Connor, M. (2019). Species richness change across spatial scales. *Oikos*, 128(8), 1079–1091. <https://doi.org/10.1111/oik.05968>
- Chiron, F., Princé, K., Paracchini, M. L., Bulgheroni, C., & Jiguet, F. (2013). Forecasting the potential impacts of CAP-associated land use changes on farmland birds at the national level. *Agriculture, Ecosystems & Environment*, 176, 17–23. <https://doi.org/10.1016/j.agee.2013.05.018>
- Davey, C. M., Chamberlain, D. E., Newson, S. E., Noble, D. G., & Johnston, A. (2012). Rise of the generalists: Evidence for climate driven homogenization in avian communities [eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1466-8238.2011.00693.x>]. *Global Ecology and Biogeography*, 21(5), 568–578. <https://doi.org/10.1111/j.1466-8238.2011.00693.x>
- Donald, P. F., Green, R. E., & Heath, M. F. (2001). Agricultural intensification and the collapse of Europe's farmland bird populations [Publisher: Royal Society]. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 268(1462), 25–29. <https://doi.org/10.1098/rspb.2000.1325>

- Dornelas, M., Gotelli, N. J., McGill, B., Shimadzu, H., Moyes, F., Sievers, C., & Magurran, A. E. (2014). Assemblage Time Series Reveal Biodiversity Change but Not Systematic Loss [Publisher: American Association for the Advancement of Science Section: Report]. *Science*, 344(6181), 296–299. <https://doi.org/10.1126/science.1248484>
- Dornelas, M., Magurran, A. E., Buckland, S. T., Chao, A., Chazdon, R. L., Colwell, R. K., Curtis, T., Gaston, K. J., Gotelli, N. J., Kosnik, M. A., McGill, B., McCune, J. L., Morlon, H., Mumby, P. J., Øvreås, L., Studeny, A., & Vellend, M. (2013). Quantifying temporal change in biodiversity: Challenges and opportunities. *Proceedings of the Royal Society B: Biological Sciences*, 280(1750), 20121931. <https://doi.org/10.1098/rspb.2012.1931>
- Doxa, A., Bas, Y., Paracchini, M. L., Pointereau, P., Terres, J.-M., & Jiguet, F. (2010). Low-intensity agriculture increases farmland bird abundances in France [\_eprint: <https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-2664.2010.01869.x>]. *Journal of Applied Ecology*, 47(6), 1348–1356. <https://doi.org/10.1111/j.1365-2664.2010.01869.x>
- Eglinton, S. M., & Pearce-Higgins, J. W. (2012). Disentangling the Relative Importance of Changes in Climate and Land-Use Intensity in Driving Recent Bird Population Trends (R. Arlettaz, Ed.). *PLoS ONE*, 7(3), e30407. <https://doi.org/10.1371/journal.pone.0030407>
- Fraixedas, S., Lindén, A., Piha, M., Cabeza, M., Gregory, R., & Lehtikainen, A. (2020). A state of the art review on birds as indicators of biodiversity: Advances, challenges, and future directions. *Ecological Indicators*, 118, 106728. <https://doi.org/10.1016/j.ecolind.2020.106728>
- Gregory, R. D., Gibbons, D. W., Impey, A., & Marchant, J. H. (1999). Generation of the headline indicator of wild bird populations, 48.
- Gregory, R. D., & Strien, A. v. (2010). Wild bird indicators: Using composite population trends of birds as measures of environmental health. *Ornithological Science*, 9(1), 3–22. <https://doi.org/10.2326/osj.9.3>
- Gregory, R. D., van Strien, A., Vorisek, P., Gmelig Meyling, A. W., Noble, D. G., Foppen, R. P., & Gibbons, D. W. (2005). Developing indicators for European birds [Publisher: Royal Society]. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 360(1454), 269–288. <https://doi.org/10.1098/rstb.2004.1602>
- Harrison, P. J., Buckland, S. T., Yuan, Y., Elston, D. A., Brewer, M. J., Johnston, A., & Pearce-Higgins, J. W. (2014). Assessing trends in biodiversity over space and time using the example of British breeding birds [\_eprint: <https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/1365-2664.12316>]. *Journal of Applied Ecology*, 51(6), 1650–1660. <https://doi.org/10.1111/1365-2664.12316>
- Harrison, P. J., Yuan, Y., Buckland, S. T., Oedekoven, C. S., Elston, D. A., Brewer, M. J., Johnston, A., & Pearce-Higgins, J. W. (2016). Quantifying turnover in biodiversity of British breeding birds [\_eprint: <https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/1365-2664.12539>]. *Journal of Applied Ecology*, 53(2), 469–478. <https://doi.org/10.1111/1365-2664.12539>
- Hill, M. O. (1973). Diversity and Evenness: A Unifying Notation and Its Consequences [\_eprint: <https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.2307/1934352>]. *Ecology*, 54(2), 427–432. <https://doi.org/10.2307/1934352>
- Inger, R., Gregory, R., Duffy, J. P., Stott, I., Voříšek, P., & Gaston, K. J. (2015). Common European birds are declining rapidly while less abundant species' numbers are rising [\_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/ele.12387>]. *Ecology Letters*, 18(1), 28–36. <https://doi.org/10.1111/ele.12387>
- Jaccard, P. (1912). The Distribution of the Flora in the Alpine Zone.1 [\_eprint: <https://nph.onlinelibrary.wiley.com/doi/pdf/10.1111/j.1469-8137.1912.tb05611.x>]. *New Phytologist*, 11(2), 37–50. <https://doi.org/10.1111/j.1469-8137.1912.tb05611.x>

- Jarzyna, M. A., & Jetz, W. (2018). Taxonomic and functional diversity change is scale dependent. *Nature Communications*, 9(1), 2565. <https://doi.org/10.1038/s41467-018-04889-z>
- Jarzyna, M. A., Zuckerberg, B., Porter, W. F., Finley, A. O., & Maurer, B. A. (2015). Spatial scaling of temporal changes in avian communities: Scale dependence of community turnover. *Global Ecology and Biogeography*, 24(11), 1236–1248. <https://doi.org/10.1111/geb.12361>
- Jiguet, F., Julliard, R., Couvet, D., & Petiau, A. (2005). Modeling Spatial Trends in Estimated Species Richness using Breeding Bird Survey Data: A Valuable Tool in Biodiversity Assessment. *Biodiversity and Conservation*, 14(13), 3305–3324. <https://doi.org/10.1007/s10531-004-0448-y>
- Jiguet, F., Devictor, V., Julliard, R., & Couvet, D. (2012). French citizens monitoring ordinary birds provide tools for conservation and ecological sciences. *Acta Oecologica*, 44, 58–66. <https://doi.org/10.1016/j.actao.2011.05.003>
- Kamp, J., Frank, C., Trautmann, S., Busch, M., Dröschmeister, R., Flade, M., Gerlach, B., Karthäuser, J., Kunz, F., Mitschke, A., Schwarz, J., & Sudfeldt, C. (2021). Population trends of common breeding birds in Germany 1990–2018. *Journal of Ornithology*, 162(1), 1–15. <https://doi.org/10.1007/s10336-020-01830-4>
- Keil, P., Schweiger, O., Kühn, I., Kunin, W. E., Kuussaari, M., Settele, J., Henle, K., Brotons, L., Pe'er, G., Lengyel, S., Moustakas, A., Steinicke, H., & Storch, D. (2012). Patterns of beta diversity in Europe: The role of climate, land cover and distance across scales [eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-2699.2012.02701.x>]. *Journal of Biogeography*, 39(8), 1473–1486. <https://doi.org/10.1111/j.1365-2699.2012.02701.x>
- McGill, B. J., Dornelas, M., Gotelli, N. J., & Magurran, A. E. (2015). Fifteen forms of biodiversity trend in the Anthropocene. *Trends in Ecology & Evolution*, 30(2), 104–113. <https://doi.org/10.1016/j.tree.2014.11.006>
- McGill, B. J., Enquist, B. J., Weiher, E., & Westoby, M. (2006). Rebuilding community ecology from functional traits. *Trends in Ecology & Evolution*, 21(4), 178–185. <https://doi.org/10.1016/j.tree.2006.02.002>
- Mouquet, N., Devictor, V., Meynard, C. N., Munoz, F., Bersier, L.-F., Chave, J., Coutron, P., Dalecky, A., Fontaine, C., Gravel, D., Hardy, O. J., Jabot, F., Lavergne, S., Leibold, M., Mouillot, D., Münkemüller, T., Pavoine, S., Prinzing, A., Rodrigues, A. S. L., ... Thuiller, W. (2012). Ecophylogenetics: Advances and perspectives [eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1469-185X.2012.00224.x>]. *Biological Reviews*, 87(4), 769–785. <https://doi.org/10.1111/j.1469-185X.2012.00224.x>
- Normander, B., Levin, G., Auvinen, A.-P., Bratli, H., Stabbetorp, O., Hedblom, M., Glimskär, A., & Gudmundsson, G. A. (2012). Indicator framework for measuring quantity and quality of biodiversity—Exemplified in the Nordic countries. *Ecological Indicators*, 13(1), 104–116. <https://doi.org/10.1016/j.ecolind.2011.05.017>
- of the Convention on Biological Diversity, S., & Centre, U. W. C. M. (Eds.). (2006). *Global biodiversity outlook 2*. Secretariat of the Convention on Biological Diversity.
- Pereira, H. M., Navarro, L. M., & Martins, I. S. (2012). Global Biodiversity Change: The Bad, the Good, and the Unknown. *Annual Review of Environment and Resources*, 37(1), 25–50. <https://doi.org/10.1146/annurev-environ-042911-093511>
- Pilotto, F., Kühn, I., Adrian, R., Alber, R., Alignier, A., Andrews, C., Bäck, J., Barbaro, L., Beaumont, D., Beenaerts, N., Benham, S., Boukal, D. S., Bretagnolle, V., Camatti, E., Canullo, R., Cardoso, P. G., Ens, B. J., Everaert, G., Evtimova, V., ... Haase, P. (2020). Meta-analysis of multidecadal biodiversity trends in Europe [Bandiera\_abtest: a Cc\_license\_type: cc\_by Cg\_type: Nature Research Journals Number: 1 Primary\_atype: Research Publisher: Nature Publishing Group Subject\_term: Biodiversity;Climate-change ecology;Macroecology Subject\_term\_id: biodiversity;climate-change-

- ecology;macroecology]. *Nature Communications*, 11(1), 3486. <https://doi.org/10.1038/s41467-020-17171-y>
- Ram, D., Axelsson, A.-L., Green, M., Smith, H. G., & Lindström, Å. (2017). What drives current population trends in forest birds – forest quantity, quality or climate? A large-scale analysis from northern Europe. *Forest Ecology and Management*, 385, 177–188. <https://doi.org/10.1016/j.foreco.2016.11.013>
- Reif, J. (2013). Long-Term Trends in Bird Populations: A Review of Patterns and Potential Drivers in North America and Europe [Publisher: Museum and Institute of Zoology, Polish Academy of Sciences]. *Acta Ornithologica*, 48(1), 1–16. <https://doi.org/10.3161/000164513X669955>
- Reif, J., Prylová, K., Šizling, A. L., Vermouzek, Z., Šťastný, K., & Bejček, V. (2013). Changes in bird community composition in the Czech Republic from 1982 to 2004: Increasing biotic homogenization, impacts of warming climate, but no trend in species richness. *Journal of Ornithology*, 154(2), 359–370. <https://doi.org/10.1007/s10336-012-0900-9>
- Sauer, J. R., Link, W. A., Fallon, J. E., Pardieck, K. L., & Ziolkowski, D. J., Jr. (2013). The North American Breeding Bird Survey 1966–2011: Summary Analysis and Species Accounts. *North American Fauna*, (79 (79)), 1–32. <https://doi.org/10.3996/nafa.79.0001>
- Sauer, J. R., Pardieck, K. L., Ziolkowski, D. J., Jr., Smith, A. C., Hudson, M.-A. R., Rodriguez, V., Berlanga, H., Niven, D. K., & Link, W. A. (2017). The first 50 years of the North American Breeding Bird Survey. *The Condor*, 119(3), 576–593. <https://doi.org/10.1650/CONDOR-17-83.1>
- Schipper, A. M., Belmaker, J., Miranda, M. D. d., Navarro, L. M., Böhning-Gaese, K., Costello, M. J., Dornelas, M., Foppen, R., Hortal, J., Huijbregts, M. A. J., Martín-López, B., Pettorelli, N., Queiroz, C., Rossberg, A. G., Santini, L., Schiffrers, K., Steinmann, Z. J. N., Visconti, P., Rondinini, C., & Pereira, H. M. (2016). Contrasting changes in the abundance and diversity of North American bird assemblages from 1971 to 2010 [eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/gcb.13292>]. *Global Change Biology*, 22(12), 3948–3959. <https://doi.org/10.1111/gcb.13292>
- Shannon, C. E. (1948). A mathematical theory of communication [Conference Name: The Bell System Technical Journal]. *The Bell System Technical Journal*, 27(3), 379–423. <https://doi.org/10.1002/j.1538-7305.1948.tb01338.x>
- Simpson, E. H. (1949). Measurement of Diversity [Bandiera\_abtest: a Cg\_type: Nature Research Journals Number: 4148 Primary\_atype: Research Publisher: Nature Publishing Group]. *Nature*, 163(4148), 688–688. <https://doi.org/10.1038/163688a0>
- Sørensen, T. J. (1948). *A method of establishing groups of equal amplitude in plant sociology based on similarity of species content and its application to analyses of the vegetation on Danish commons*. [OCLC: 4713331]. I kommission hos E. Munksgaard.
- Sorte, F. A. L., & Boecklen, W. J. (2005). Changes in the diversity structure of avian assemblages in North America [eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1466-822X.2005.00160.x>]. *Global Ecology and Biogeography*, 14(4), 367–378. <https://doi.org/10.1111/j.1466-822X.2005.00160.x>
- Studený, A. C., Buckland, S. T., Harrison, P. J., Illian, J. B., Magurran, A. E., & Newson, S. E. (2013). Fine-tuning the assessment of large-scale temporal trends in biodiversity using the example of British breeding birds [eprint: <https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/1365-2664.12026>]. *Journal of Applied Ecology*, 50(1), 190–198. <https://doi.org/10.1111/1365-2664.12026>
- Vaidyanathan, G. (2021). The world's species are playing musical chairs: How will it end? [Bandiera\_abtest: a Cg\_type: News Feature Number: 7870 Publisher: Nature Publishing Group Subject\_term: Biodiversity, Ecology, Environmental sciences]. *Nature*, 596(7870), 22–25. <https://doi.org/10.1038/d41586-021-02088-3>

- Van Turnhout, C. A. M., Foppen, R. P. B., Leuven, R. S. E. W., Siepel, H., & Esselink, H. (2007). Scale-dependent homogenization: Changes in breeding bird diversity in the Netherlands over a 25-year period. *Biological Conservation*, 134(4), 505–516. <https://doi.org/10.1016/j.biocon.2006.09.011>
- Webb, C. O., Ackerly, D. D., McPeck, M. A., & Donoghue, M. J. (2002). Phylogenies and Community Ecology [eprint: <https://doi.org/10.1146/annurev.ecolsys.33.010802.150448>]. *Annual Review of Ecology and Systematics*, 33(1), 475–505. <https://doi.org/10.1146/annurev.ecolsys.33.010802.150448>
- Whittaker, R. H. (1960). Vegetation of the Siskiyou Mountains, Oregon and California [eprint: <https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.2307/1943563>]. *Ecological Monographs*, 30(3), 279–338. <https://doi.org/10.2307/1943563>
- Whittaker, R. H. (1972). Evolution and Measurement of Species Diversity [eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.2307/1218190>]. *TAXON*, 21(2-3), 213–251. <https://doi.org/10.2307/1218190>
- Wretenberg, J., Pärt, T., & Berg, Å. (2010). Changes in local species richness of farmland birds in relation to land-use changes and landscape structure. *Biological Conservation*, 143(2), 375–381. <https://doi.org/10.1016/j.biocon.2009.11.001>

## Supplementary materials



Table 6.1: Supplementary informations about each article

Reference	Spatial grain (Km <sup>2</sup> )	Trend	Note
Barnagaud et al., <a href="#">2017</a>	Local	Increase	Not sure that it is at the road scale: "Taxonomic evenness showed a marginal, yet significant, non-linear increase from close to 0.54 in the first decade to 0.56 in the last decade (Table 1), suggesting a light trend towards a more even distribution of species' abundances among species within local assemblages "
	Local	Increase	Mean change of SR at the road scales Area of the road = $(40/0.8) * (\pi * 4002)$ with a road of 40 Km with point counts spaced by 0.8 Km and a census radius of 400m
Bowler et al., <a href="#">2021</a>	National	Stable	Metric = MSI, as many and as intense increase (i.e. Czech Rep. and Switzerland) than decrease (i.e. Germany and Denmark)
Chase et al., <a href="#">2019</a>	Local	Stable	NA
	Regional	Stable	NA
	Regional	Stable	NA
	Local	Increase	NA
	Local	Increase	NA
Chiron et al., <a href="#">2013</a>	Regional	Decrease	Concerning the spatial scale, predictions are made using the spatial unit of 4 Km <sup>2</sup> and the FBI is computed for each region of France, then meanned. Prediction with baseline scenario
	Regional	Decrease	FBI prediction with CAP greening cenario
	Regional	Decrease	FBI prediction with No Pillar I scenario
	Regional	Decrease	FBI prediction with biofuel scenario
Davey et al., <a href="#">2012</a>	Local	Increase	Metric = Simpson.They predict the metric using a GAM with spatial resolution of 1 Km <sup>2</sup> . Then they show the trend for the mean value of the metric per year
	Local	Increase	NA
	Local	Increase	NA

Table 6.1: Supplementary informations about each article (*continued*)

Reference	Spatial grain (Km <sup>2</sup> )	Trend	Note
Harrison et al., <a href="#">2014</a>	Local	Increase	To assess the metric, they use a GAM to predict the abundance over the entire area of interest (spatial resolution = 1 Km <sup>2</sup> ) and then compute the geometric mean of species abundance = Multi Species Index (as in Studeny et al., <a href="#">2013</a> ) from the prediction. Data used to learn the GAM are sampled from plots of 1 Km <sup>2</sup> . Farmland communities
	Local	Stable	Farmland communities, GoF ( $\lambda = -1$ ) = weighted towards the rare species
	Local	Stable	Farmland communities, GoF ( $\lambda = -2$ ) weighted towards the common species
Harrison et al., <a href="#">2016</a>	Local	Increase	Geometric mean of species abundance, they predict the abundance with resolution of 1 Km <sup>2</sup> and then computed the metric for each 10000 Km <sup>2</sup> cell across Great Britain, Visited twice a year
	Local	Stable	GoF ( $\lambda = -1$ ) = toward rare species" The goodness-of-fit-based measure of biodiversity suggests that both rare and common species made gains through much of Britain in the first half of the time period, and losses in the second half.", Visited twice a year / Increase first half and second second halfGoF ( $\lambda = -1$ )
	Local	Stable	GoF ( $\lambda = -2$ ) = toward common species " The goodness-of-fit-based measure of biodiversity suggests that both rare and common species made gains through much of Britain in the first half of the time period, and losses in the second half.", Visited twice a year / Increase first half and second second half
Jarzyna and Jetz, <a href="#">2018</a>	Regional	Increase	NA
	Regional	Increase	NA
	Regional	Increase	NA
	National	Increase	NA
	Global	Decrease	NA
	Regional	Increase	NA
	Regional	Increase	NA

Table 6.1: Supplementary informations about each article (*continued*)

Reference	Spatial grain (Km <sup>2</sup> )	Trend	Note
	Regional	Increase	NA
	National	Increase	NA
Pilotto et al., <a href="#">2020</a>	Global	Stable	NA
	Local	Stable	"Analyses of the trends in local biodiversity over large spatial scales"
	Local	Increase	Metric = Simpson, "Analyses of the trends in local biodiversity over large spatial scales"
	Local	Increase	"Analyses of the trends in local biodiversity over large spatial scales"
	Local	Stable	"Analyses of the trends in local biodiversity over large spatial scales"
Ram et al., <a href="#">2017</a>	Local	Increase	MSI for forest species, road of 8 Km with no limitations so assumed 200m
	Regional	Increase	SR for forest species meaned over roads, spatial grain = 8* .4 with road of 8 Km and census radius no limitations so assumed 200m
Reif et al., <a href="#">2013</a>	Local	Stable	Jaccard index, pairwise comparisons between transects
	Local	Stable	JPSP data, transect scale
	National	Stable	JPSP data, national scale
Schipper et al., <a href="#">2016</a>	Local	Increase	The metric (i.e. geometric mean) is meaned over each road. Area of the road = 50*(pi*4002) with 50 census point per road and a census radius of 400m
	Local	Increase	Metric = Shannon
	Local	Increase	Metric = Simpson
	Local	Decrease	NA
	Local	Increase	NA
	Local	Increase	NA
	Local	Increase	NA

Table 6.1: Supplementary informations about each article (*continued*)

Reference	Spatial grain (Km <sup>2</sup> )	Trend	Note
Sorte and Boecklen, <a href="#">2005</a>	Local	Decrease	NA
	Local	Decrease	Metric = evenness
	Local	Increase	The metric is meaned over each road. Area of the road = $50 \cdot (\pi \cdot 400^2)$ with 50 census point per road and a census radius of 400m
Van Turnhout et al., <a href="#">2007</a>	Regional	Increase	For each region, the trend is computed using the mean number of species per atlas square
	Local	Increase	Mainly increase of SR but the proportion of negative trend were higher than for the regional scale
	National	Increase	National scale
Wretenberg et al., <a href="#">2010</a>	Local	Decrease	looking at the trend through different environmental policies, " local species richness (i.e. at the scale of sites = 3 hectares) decreased significantly probably as a result of an overall reduced abundance of several species. "
Inger et al., <a href="#">2015</a>	National	Decrease	NA
Donald et al., <a href="#">2001</a>	National	Decrease	The metric is referred as "Mean population" and the trend is estimated for each european country