Bilangan Tersembunyi Dilil

Author: Notdils

INDONESIAN VERSION

Batas Waktu	Batas Memori
1.0 Detik	256 MB



Dilil adalah seorang pencari harta karun yang legendaris. Ia menemukan sebuah scroll kuno yang berisi deretan angka-angka misterius. Scroll tersebut menampilkan N buah bilangan bulat dalam urutan acak. Diketahui, terdapat satu sesi rahasia berisi M buah bilangan ($M < N$) yang merupakan kunci untuk membuka peti harta karun.

Namun, ada yang aneh! semua bilangan dalam sesi rahasia tersebut PASTI muncul di dalam deretan angka-angka utama. Dilil perlu menemukan bilangan-bilangan yang ada di deretan utama tetapi TIDAK termasuk dalam sesi rahasia. Jumlahan dari bilangan-bilangan tersebut adalah kombinasi kunci untuk peti harta!

Diberikan dua buah array A dan B. Array A berisi N buah bilangan bulat. Array B berisi M buah bilangan bulat ($M < N$). Diketahui bahwa seluruh elemen pada array B dipastikan ada di dalam array A dengan posisi yang acak.

Tugasmu adalah menghitung total jumlah (sum) dari semua elemen pada array A yang TIDAK muncul pada array B.

Format Masukan:
Baris pertama berisi dua buah bilangan bulat N dan M ($1 \leq M < N \leq 100$).
Baris kedua berisi N buah bilangan bulat $A_1, A_2, \ldots, A_N$ ($1 \leq A_i \leq 1000$).
Baris ketiga berisi M buah bilangan bulat $B_1, B_2, \ldots, B_M$ ($1 \leq B_i \leq 1000$).
Dipastikan Semua Input $A_1, A_2, \ldots, A_N$ dan $B_1, B_2, \ldots, B_M$ Berbeda.

Format Keluaran:
Sebuah bilangan bulat yang menyatakan hasil penjumlahan elemen array A yang tidak ada pada array B.

Contoh Masukan 1

```
5 2
4 8 2 1 9
8 1
```

Contoh Keluaran 1

```
15
```

Penjelasan:
Elemen array A adalah $\{4, 8, 2, 1, 9\}$. Elemen array B adalah $\{8, 1\}$. Elemen pada A yang tidak ada di B adalah $4, 2, 9$. Total penjumlahan: $4 + 2 + 9 = 15$.

Dilil's Hidden Numbers

Author: Notdils

ENGLISH VERSION

Time limit	Memory Limit
1.0 Second	256 MB



Dilil is a legendary treasure hunter. He discovers an ancient scroll containing a mysterious sequence of numbers. The scroll displays N integers in random order. It is known that there is a secret session containing M numbers ($M < N$) that is the key to opening a treasure chest.

However, something is strange! all numbers in this secret session are guaranteed to appear in the main sequence of numbers. Dilil needs to find the numbers that exist in the main sequence but are NOT included in the secret session. The sum of these numbers is the combination key to the treasure chest!

You are given two arrays: A and B. Array A contains N integers. Array B contains M integers ($M < N$). It is guaranteed that all elements in array B are present in array A at random positions. Your task is to calculate the total sum of all elements in array A that do NOT appear in array B.

Input Format:
The first line contains two integers N and M ($1 \leq M < N \leq 100$).
The second line contains N integers $A_1, A_2, \ldots, A_N$ ($1 \leq A_i \leq 1000$).
The third line contains M integers $B_1, B_2, \ldots, B_M$ ($1 \leq B_i \leq 1000$).
It is guaranteed that all inputs $A_1, A_2, \ldots, A_N$ and $B_1, B_2, \ldots, B_M$ are all unique.

Output Format:
A single integer representing the sum of all elements in array A that are not present in array B.

Sample Input 1

```
5 2
4 8 2 1 9
8 1
```

Sample Output 1

```
15
```

Explanation:
The elements of array A are $\{4, 8, 2, 1, 9\}$. The elements of array B are $\{8, 1\}$. The elements in A that do not appear in B are $4, 2$, and 9. The total sum is: $4 + 2 + 9 = 15$.