

Generalized Integral Representations for Functions with Values in $C(V_{3,3})^*$

Klaus GÜRLEBECK¹ Zhongxiang ZHANG²

Abstract By using the solution to the Helmholtz equation $\Delta u - \lambda u = 0$ ($\lambda \geq 0$), the explicit forms of the so-called kernel functions and the higher order kernel functions are given. Then by the generalized Stokes formula, the integral representation formulas related with the Helmholtz operator for functions with values in $C(V_{3,3})$ are obtained. As application of the integral representations, the maximum modulus theorem for function u which satisfies $Hu = 0$ is given.

Keywords Universal Clifford algebra, Helmholtz equation, Generalized Cauchy-Pompeiu formula

2000 MR Subject Classification 30G35

1 Introduction and Preliminaries

Integral representation formulas in complex analysis and Clifford analysis have been well-developed in [2–9, 11–21, 24–32], etc. These integral representation formulas are powerful tools. In [24], E. Obolashvili obtained some generalized Cauchy-Pompeiu representation formulas. In [25–26], higher order Cauchy-Pompeiu type representation formulas related to the factors and powers of the Helmholtz operator were given in quaternionic analysis in case of taking $\lambda = |h|^2$ with a complex or real number h . In [8], in case of taking $\lambda = |h|^2$ with a hyper-complex number h , we constructed the kernel functions and then got the higher order integral representation formulas for functions with values in a universal Clifford algebra $C(V_{n,n})$. In this paper, by using the idea in [8, 24, 28], we get the explicit expressions of the kernel functions and then get the explicit integral representation formulas for functions with values in $C(V_{3,3})$. These explicit integral representation formulas will play an important role in studying the further properties of the functions with values in $C(V_{3,3})$. In the last section, as application of the integral representations, we give the maximum modulus theorem for function u which satisfies $Hu = 0$.

Let $V_{n,s}$ ($0 \leq s \leq n$) be an n -dimensional ($n \geq 1$) real linear space with basis $\{e_1, e_2, \dots, e_n\}$, $C(V_{n,s})$ be the 2^n -dimensional real linear space with basis

$$\{e_A, A = \{h_1, \dots, h_r\} \in \mathcal{PN}, 1 \leq h_1 < \dots < h_r \leq n\},$$

Manuscript received October 25, 2008. Revised January 21, 2010. Published online December 28, 2010.

¹Institut f. Mathematik/Physik, Bauhaus-Universität Weimar, Coudraystr. 13, D-99423 Weimar, Germany.
E-mail: klaus.guerlebeck@uni-weimar.de

²Corresponding author. School of Mathematics and Statistics, Wuhan University, Wuhan 430072, China.
E-mail: zhangzx9@126.com

*Project supported by Deutscher Akademischer Austausch Dienst (German Academic Exchange Service) and the National Natural Science Foundation of China (No. 10471107).

where N stands for the set $\{1, \dots, n\}$ and $\mathcal{P}N$ denotes the family of all order-preserving subsets of N in the above way. we denote e_\emptyset as e_0 and e_A as $e_{h_1 \dots h_r}$ for $A = \{h_1, \dots, h_r\} \in \mathcal{P}N$. The product on $C(V_{n,s})$ is defined by

$$\begin{cases} e_A e_B = (-1)^{\#((A \cap B) \setminus S)} (-1)^{P(A,B)} e_{A \Delta B}, & \text{if } A, B \in \mathcal{P}N, \\ \lambda \mu = \sum_{A \in \mathcal{P}N} \sum_{B \in \mathcal{P}N} \lambda_A \mu_B e_A e_B, & \text{if } \lambda = \sum_{A \in \mathcal{P}N} \lambda_A e_A, \mu = \sum_{B \in \mathcal{P}N} \mu_B e_B, \end{cases} \quad (1.1)$$

where S stands for the set $\{1, \dots, s\}$, $\#(A)$ is the cardinal number of the set A , the number $P(A, B) = \sum_{j \in B} P(A, j)$, $P(A, j) = \#\{i, i \in A, i > j\}$, the symmetric difference set $A \Delta B$ is also order-preserving in the above way, and $\lambda_A \in \mathcal{R}$ is the coefficient of the e_A -component of the Clifford number λ . We also denote λ_A as $[\lambda]_A$, $\lambda_{\{i\}}$ as $[\lambda]_i$ and λ_0 as $\text{Re } \lambda$. It follows immediately from the multiplication rule (1.1) that e_0 is the identity element written now as 1 and in particular,

$$\begin{cases} e_i^2 = 1, & \text{if } i = 1, \dots, s, \\ e_j^2 = -1, & \text{if } j = s+1, \dots, n, \\ e_i e_j = -e_j e_i, & \text{if } 1 \leq i < j \leq n, \\ e_{h_1} e_{h_2} \dots e_{h_r} = e_{h_1 h_2 \dots h_r}, & \text{if } 1 \leq h_1 < h_2 < \dots < h_r \leq n. \end{cases} \quad (1.2)$$

Thus $C(V_{n,s})$ is a real linear, associative, but non-commutative algebra and it is called the universal Clifford algebra over $V_{n,s}$. In particular, suppose $n = s = 3$. Then $C(V_{n,s})$ is just the universal Clifford algebra $C(V_{3,3})$.

An involution is defined by

$$\begin{cases} \bar{e}_A = (-1)^{\sigma(A) + \#(A \cap S)} e_A, & \text{if } A \in \mathcal{P}N, \\ \bar{\lambda} = \sum_{A \in \mathcal{P}N} \lambda_A \bar{e}_A, & \text{if } \lambda = \sum_{A \in \mathcal{P}N} \lambda_A e_A, \end{cases} \quad (1.3)$$

where $\sigma(A) = \frac{\#(A)(\#(A)+1)}{2}$. From (1.1) and (1.3), we have

$$\begin{cases} \bar{e}_i = e_i, & \text{if } i = 0, 1, \dots, s, \\ \bar{e}_j = -e_j, & \text{if } j = s+1, \dots, n, \\ \overline{\lambda \mu} = \bar{\mu} \bar{\lambda}, & \text{for any } \lambda, \mu \in C(V_{n,s}). \end{cases} \quad (1.4)$$

The $C(V_{n,n})$ -valued $(n-1)$ -differential form

$$d\sigma = \sum_{k=1}^n (-1)^{k-1} e_k d\hat{x}_k^N$$

is exact, where

$$d\hat{x}_k^N = dx^1 \wedge \dots \wedge dx^{k-1} \wedge dx^{k+1} \wedge \dots \wedge dx^n.$$

In this paper, we shall only consider the special case $s = n = 3$. Let Ω be an open non-empty subset of \mathcal{R}^3 . Functions f defined in Ω and with values in $C(V_{3,3})$ will be considered, i.e., $f : \Omega \rightarrow C(V_{3,3})$. $f \in C^{(r)}(\Omega, C(V_{3,3}))$ is clear that all the component functions of $f(x)$ possess the cited property. The operator D which is written as

$$D = \sum_{k=1}^3 e_k \frac{\partial}{\partial x_k} : C^{(r)}(\Omega, C(V_{3,3})) \rightarrow C^{(r-1)}(\Omega, C(V_{3,3})).$$

The operator D acts on the function f from the left and from the right being governed by the rule

$$D[f] = \sum_{k=1}^3 \sum_A e_k e_A \frac{\partial f_A}{\partial x_k}, \quad [f]D = \sum_{k=1}^3 \sum_A e_A e_k \frac{\partial f_A}{\partial x_k}.$$

The classical Riemann boundary value problems and singular integral equations are widely used for the solution to a great field of problems in applied mechanics theory (see [22, 23] etc.). Naturally, it is interesting to generalize the theory of holomorphic functions in the plane to higher dimensions. Especially, it has practical meaning to generalize the classical theory in the plane to the theory of k -regular functions defined in \mathcal{R}^3 with values in $C(V_{3,3})$. For example, R_m ($m > 0$) Riemann boundary value problem for bi-harmonic function was studied in [20], as for the R_m ($m > 0$) Riemann boundary value problem for bi-harmonic functions, and even more general R_m ($m > 0$) Riemann boundary value problem for k -regular functions, the higher order integral representation formulas play a very important role. The higher order integral representation formulas provide a special solution to the R_m ($m > 0$) Riemann boundary value problem. The importance is similar to the classical Cauchy type integral in complex plane. On the other hand, the generalized Liouville theorem for k -regular functions is very important for solving the R_m ($m > 0$) Riemann boundary value problem. In [1, 10], the generalized Liouville theorem for k -regular functions in Clifford analysis was solved under some growth conditions at infinity. It is a very important result, while it is also valuable to refine the too many growth conditions at infinity especially for Riemann boundary value problems. But how to refine the conditions is difficult. For harmonic functions and bi-harmonic functions defined in \mathcal{R}^3 and with values in $C(V_{3,3})$, we know that the growth conditions at infinity of the generalized Liouville theorem can be refined to one of the conditions in [1, 10], and the method depends on the higher order integral representation formulas (see [20]). In view of the above reasons, we are interested in the different types of higher order integral representation formulas in Clifford analysis. As is well-known that the kernel functions play the most important role in obtaining the higher order integral representation formulas. While in many cases, the explicit expressions of the kernel functions are important to study the further properties of functions, for example, the generalized Liouville theorem of k -regular functions in Clifford analysis.

As for the above purposes, we keep our study in the framework of universal Clifford algebra $C(V_{3,3})$ in this paper. On one hand, it has practical meaning in applied mechanics theory; on the other hand, it provides a foundation to study the more general theory of functions with values in $C(V_{n,n})$.

We consider the following operators:

$$L_\lambda u = Du + \lambda u, \quad L_{*\lambda} u = uD - \lambda u, \quad (1.5)$$

$$Lu = Du + uh, \quad L_* u = uD - hu \quad (1.6)$$

and the Helmholtz operator

$$Hu = (\Delta - |h|^2)u, \quad (1.7)$$

where

$$h = \sum_{i=1}^3 h_i e_i. \quad (1.8)$$

2 Generalized Integral Representations

Denote the Helmholtz equation as

$$\Delta u - |h|^2 u = 0. \quad (2.1)$$

Lemma 2.1 *Let $g_3(|x|, |h|) = \frac{1}{|x|} e^{-|h||x|}$. Then $g_3(|x|, |h|)$ is the fundamental solution to (2.1) in \mathcal{R}^3 .*

Lemma 2.2 *Let $g_3(|x|, |h|)$ be defined as in Lemma 2.1. Then*

(i) *$D[g_3](|x|, |h|)e_A - g_3(|x|, |h|)e_A h$ is the fundamental solution to equation*

$$Lu = Du + uh = 0, \quad h = \sum_{i=1}^3 h_i e_i, \quad (2.2)$$

(ii) *$e_A D[g_3](|x|, |h|) + h e_A g_3(|x|, |h|)$ is the fundamental solution to equation*

$$L_* u = uD - hu = 0, \quad h = \sum_{i=1}^3 h_i e_i. \quad (2.3)$$

In the following, Ω is supposed to be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$.

Lemma 2.3 (see [8]) *Let $u, v \in C^1(\Omega, C(V_{3,3})) \cap C(\overline{\Omega}, C(V_{3,3}))$. Then*

$$\operatorname{Re} \left\{ \int_{\Omega} (vL[u] + L_*[v]u) dy \right\} = \operatorname{Re} \left\{ \int_{\partial\Omega} v d\sigma_y u \right\}. \quad (2.4)$$

Lemma 2.4 *Let $u, v \in C^1(\Omega, C(V_{3,3})) \cap C(\overline{\Omega}, C(V_{3,3}))$. Then*

$$\int_{\Omega} (vL_{\lambda}[u] + L_{*\lambda}[v]u) dy = \int_{\partial\Omega} v d\sigma_y u. \quad (2.5)$$

Denote

$$K_1^A(r, h) = \frac{1}{\omega_3} \left(\frac{(\mathbf{y} - \mathbf{x})e_A}{r^3} + \frac{|h|(\mathbf{y} - \mathbf{x})e_A}{r^2} + \frac{e_A h}{r} \right) e^{-|h|r}, \quad (2.6)$$

$$K_{*1}^A(r, h) = \frac{1}{\omega_3} \left(\frac{e_A(\mathbf{y} - \mathbf{x})}{r^3} + \frac{e_A |h|(\mathbf{y} - \mathbf{x})}{r^2} - \frac{h e_A}{r} \right) e^{-|h|r}, \quad (2.7)$$

$$K_1(r, \lambda) = \frac{1}{\omega_3} \left(\frac{\mathbf{y} - \mathbf{x}}{r^3} + \frac{\lambda(\mathbf{y} - \mathbf{x})}{r^2} + \frac{\lambda}{r} \right) e^{-\lambda r}, \quad (2.8)$$

$$K_{*1}(r, \lambda) = \frac{1}{\omega_3} \left(\frac{\mathbf{y} - \mathbf{x}}{r^3} + \frac{\lambda(\mathbf{y} - \mathbf{x})}{r^2} - \frac{\lambda}{r} \right) e^{-\lambda r}, \quad (2.9)$$

where $\mathbf{y} - \mathbf{x} = \sum_{i=1}^3 (y_i - x_i) e_i$, $r = |\mathbf{y} - \mathbf{x}|$ and ω_3 denotes the area of the unit sphere in \mathcal{R}^3 .

Lemma 2.5 *Let $K_1^A(r, h)$ and $K_{*1}^A(r, h)$ be as in (2.6) and (2.7), where $r = |\mathbf{y} - \mathbf{x}|$. Then*

$$\begin{cases} L[K_1^A(r, h)] = 0, & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}, \\ L_*[K_{*1}^A(r, h)] = 0, & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}. \end{cases} \quad (2.10)$$

Lemma 2.6 Let $K_1(r, \lambda)$ and $K_{*1}(r, \lambda)$ be as in (2.8) and (2.9), where $r = |y - x|$. Then

$$\begin{cases} L_\lambda[K_1(r, \lambda)] = L_{*-\lambda}[K_1(r, \lambda)] = 0, & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}, \\ L_{-\lambda}[K_{*1}(r, \lambda)] = L_{*\lambda}[K_{*1}(r, \lambda)] = 0, & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}. \end{cases} \quad (2.11)$$

Remark 2.1 By Lemmas 2.5 and 2.6, $K_1^A(r, 0)\bar{e}_A$, $\bar{e}_A K_{*1}^A(r, 0)$, $K_1(r, 0)$ and $K_{*1}(r, 0)$ are the classical kernel functions.

Lemma 2.7 Let $K_1^A(r, h)$ be as in (2.6). Then

$$\int_{\partial B(x, \delta)} d\sigma_y K_1^A(r, h) = (1 + \delta|h|)e_A e^{-\delta|h|}, \quad (2.12)$$

where $B(x, \delta) = \left\{y, \sum_{i=1}^3 (y_i - x_i)^2 \leq \delta^2\right\}$ and $\partial B(x, \delta)$ is given the induced orientation.

Proof By Stokes formula, we have

$$\begin{aligned} \int_{\partial B(x, \delta)} d\sigma_y K_1^A(r, h) &= \frac{1}{\omega_3} \int_{\partial B(x, \delta)} d\sigma_y \left(\frac{(\mathbf{y} - \mathbf{x})e_A}{\delta^3} + \frac{|h|(\mathbf{y} - \mathbf{x})e_A}{\delta^2} + \frac{e_A h}{\delta} \right) e^{-\delta|h|} \\ &= \frac{1}{\omega_3} \int_{B(x, \delta)} \left(\frac{3}{\delta^3} + \frac{3|h|}{\delta^2} \right) e_A e^{-\delta|h|} d\mathbf{y} \\ &= (1 + \delta|h|)e_A e^{-\delta|h|}. \end{aligned}$$

Thus, the result follows.

Lemma 2.8 Let $K_{*1}^A(r, h)$ be as in (2.7). Then

$$\int_{\partial B(x, \delta)} K_{*1}^A(r, h) d\sigma_y = e_A (1 + \delta|h|) e^{-\delta|h|}, \quad (2.13)$$

where $B(x, \delta) = \left\{y, \sum_{i=1}^3 (y_i - x_i)^2 \leq \delta^2\right\}$ and $\partial B(x, \delta)$ is given the induced orientation.

Theorem 2.1 (Generalized Cauchy-Pompeiu Formula) Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^1(\Omega, C(V_{3,3})) \cap C(\bar{\Omega}, C(V_{3,3}))$. Then for $x \in \Omega$,

$$\begin{aligned} u(x) &= \frac{1}{\omega_3} \int_{\partial\Omega} \left(\frac{\mathbf{y} - \mathbf{x}}{r^3} + \frac{|h|(\mathbf{y} - \mathbf{x})}{r^2} \right) e^{-|h|r} d\sigma_y u(y) - \frac{1}{\omega_3} \int_{\partial\Omega} \frac{e^{-|h|r}}{r} d\sigma_y u(y) h \\ &\quad - \frac{1}{\omega_3} \int_{\Omega} \left\{ \left(\frac{\mathbf{y} - \mathbf{x}}{r^3} + \frac{|h|(\mathbf{y} - \mathbf{x})}{r^2} \right) e^{-|h|r} L[u](y) - \frac{e^{-|h|r}}{r} L[u](y) h \right\} d\mathbf{y}. \end{aligned} \quad (2.14)$$

Proof For $x \in \Omega$, by Lemmas 2.3 and 2.5, we have

$$\operatorname{Re} \int_{\partial\Omega \setminus \partial B(x, \varepsilon)} K_{*1}^A(r, h) d\sigma_y u(y) = \operatorname{Re} \int_{\Omega \setminus B(x, \varepsilon)} K_{*1}^A(r, h) L[u] d\mathbf{y}. \quad (2.15)$$

Obviously,

$$\operatorname{Re} \int_{\partial B(x, \varepsilon)} K_{*1}^A(r, h) d\sigma_y u(y)$$

$$= \operatorname{Re} \int_{\partial B(x, \varepsilon)} K_{*1}^A(r, h) d\sigma_y (u(y) - u(x)) + \operatorname{Re} \int_{\partial B(x, \varepsilon)} K_{*1}^A(r, h) d\sigma_y u(x). \quad (2.16)$$

By Lemma 2.8, it can be proved that

$$\lim_{\varepsilon \rightarrow 0} \operatorname{Re} \int_{\partial B(x, \varepsilon)} K_{*1}^A(r, h) d\sigma_y u(x) = u_A(x) e_A^2. \quad (2.17)$$

In view of $u \in C^1(\Omega, C(V_{3,3})) \cap C(\overline{\Omega}, C(V_{3,3}))$, by Stokes formula, it can be similarly proved as Lemma 2.8 that

$$\lim_{\varepsilon \rightarrow 0} \operatorname{Re} \int_{\partial B(x, \varepsilon)} K_{*1}^A(r, h) d\sigma_y (u(y) - u(x)) = 0. \quad (2.18)$$

Taking limit $\varepsilon \rightarrow 0$ in (2.15), in view of the weak singularity of $K_{*1}^A(r, h)$ and (2.16)–(2.18), we have

$$u_A(x) e_A^2 = \operatorname{Re} \int_{\partial \Omega} K_{*1}^A(r, h) d\sigma_y u(y) - \operatorname{Re} \int_{\Omega} K_{*1}^A(r, h) L[u] dy. \quad (2.19)$$

Thus

$$\begin{aligned} u_A(x) &= \frac{e_A^2}{\omega_3} \operatorname{Re} \int_{\partial \Omega} e_A \left(\frac{\mathbf{y} - \mathbf{x}}{r^3} + \frac{|h|(\mathbf{y} - \mathbf{x})}{r^2} \right) e^{-|h|r} d\sigma_y u(y) - \frac{e_A^2}{\omega_3} \operatorname{Re} \int_{\partial \Omega} h e_A \frac{e^{-|h|r}}{r} d\sigma_y u(y) \\ &\quad - \frac{e_A^2}{\omega_3} \operatorname{Re} \int_{\Omega} \left\{ e_A \left(\frac{\mathbf{y} - \mathbf{x}}{r^3} + \frac{|h|(\mathbf{y} - \mathbf{x})}{r^2} \right) e^{-|h|r} L[u] - h e_A \frac{e^{-|h|r}}{r} L[u] \right\} dy. \end{aligned} \quad (2.20)$$

From (2.20), it is easy to check that

$$\begin{aligned} u_A(x) &= \frac{e_A^2}{\omega_3} \operatorname{Re} \int_{\partial \Omega} e_A \left(\frac{\mathbf{y} - \mathbf{x}}{r^3} + \frac{|h|(\mathbf{y} - \mathbf{x})}{r^2} \right) e^{-|h|r} d\sigma_y u(y) - \frac{e_A^2}{\omega_3} \operatorname{Re} \int_{\partial \Omega} e_A \frac{e^{-|h|r}}{r} d\sigma_y u(y) h \\ &\quad - \frac{e_A^2}{\omega_3} \operatorname{Re} \int_{\Omega} \left\{ e_A \left(\frac{\mathbf{y} - \mathbf{x}}{r^3} + \frac{|h|(\mathbf{y} - \mathbf{x})}{r^2} \right) e^{-|h|r} L[u] - e_A \frac{e^{-|h|r}}{r} L[u] h \right\} dy. \end{aligned} \quad (2.21)$$

Then the result follows.

Theorem 2.2 (Generalized Cauchy-Pompeiu Formula) *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial \Omega$, $u \in C^1(\Omega, C(V_{3,3})) \cap C(\overline{\Omega}, C(V_{3,3}))$. Then for $x \in \Omega$,*

$$\begin{aligned} u(x) &= \frac{1}{\omega_3} \int_{\partial \Omega} u(y) d\sigma_y \left(\frac{\mathbf{y} - \mathbf{x}}{r^3} + \frac{|h|(\mathbf{y} - \mathbf{x})}{r^2} \right) e^{-|h|r} + \frac{1}{\omega_3} \int_{\partial \Omega} h u(y) d\sigma_y \frac{e^{-|h|r}}{r} \\ &\quad - \frac{1}{\omega_3} \int_{\Omega} \left\{ L_*[u](y) \left(\frac{\mathbf{y} - \mathbf{x}}{r^3} + \frac{|h|(\mathbf{y} - \mathbf{x})}{r^2} \right) e^{-|h|r} + h L_*[u](y) \frac{e^{-|h|r}}{r} \right\} dy. \end{aligned} \quad (2.22)$$

By Theorems 2.1 and 2.2, we can deduce the following result.

Theorem 2.3 (Generalized Cauchy Integral Formula) *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial \Omega$, $u \in C^1(\Omega, C(V_{3,3})) \cap C(\overline{\Omega}, C(V_{3,3}))$, and $Lu = 0$ in Ω . Then for $x \in \Omega$,*

$$u(x) = \frac{1}{\omega_3} \int_{\partial \Omega} \left(\frac{\mathbf{y} - \mathbf{x}}{r^3} + \frac{|h|(\mathbf{y} - \mathbf{x})}{r^2} \right) e^{-|h|r} d\sigma_y u(y) - \frac{1}{\omega_3} \int_{\partial \Omega} \frac{e^{-|h|r}}{r} d\sigma_y u(y) h. \quad (2.23)$$

Theorem 2.4 (Generalized Cauchy Integral Formula) *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^1(\Omega, C(V_{3,3})) \cap C(\overline{\Omega}, C(V_{3,3}))$, and $L_*u = 0$ in Ω . Then for $x \in \Omega$,*

$$u(x) = \frac{1}{\omega_3} \int_{\partial\Omega} u(y) d\sigma_y \left(\frac{\mathbf{y} - \mathbf{x}}{r^3} + \frac{|h|(\mathbf{y} - \mathbf{x})}{r^2} \right) e^{-|h|r} + \frac{1}{\omega_3} \int_{\partial\Omega} hu(y) d\sigma_y \frac{e^{-|h|r}}{r}. \quad (2.24)$$

Remark 2.2 In case of $h = 0$, Theorems 2.1–2.4 are respectively the classical Cauchy-Pompeiu formula and Cauchy integral formula in Clifford analysis.

By Lemmas 2.4 and 2.6, the following theorems can be similarly proved as Theorem 2.1.

Theorem 2.5 *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^1(\Omega, C(V_{3,3})) \cap C(\overline{\Omega}, C(V_{3,3}))$, $K_{*1}(r, \lambda)$ be as in (2.9). Then*

$$\int_{\partial\Omega} K_{*1}(r, \lambda) d\sigma_y u(y) - \int_{\Omega} K_{*1}(r, \lambda) L_{\lambda}[u](y) dy = \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \quad (2.25)$$

Theorem 2.6 *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^1(\Omega, C(V_{3,3})) \cap C(\overline{\Omega}, C(V_{3,3}))$, $K_1(r, \lambda)$ be as in (2.8). Then*

$$\int_{\partial\Omega} K_1(r, \lambda) d\sigma_y u(y) - \int_{\Omega} K_1(r, \lambda) L_{-\lambda}[u](y) dy = \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \quad (2.26)$$

Theorem 2.7 *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^1(\Omega, C(V_{3,3})) \cap C(\overline{\Omega}, C(V_{3,3}))$, $K_{*1}(r, \lambda)$ be as in (2.9). Then*

$$\int_{\partial\Omega} u(y) d\sigma_y K_{*1}(r, \lambda) - \int_{\Omega} L_{*-\lambda}[u](y) K_{*1}(r, \lambda) dy = \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \quad (2.27)$$

Theorem 2.8 *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^1(\Omega, C(V_{3,3})) \cap C(\overline{\Omega}, C(V_{3,3}))$, $K_1(r, \lambda)$ be as in (2.8). Then*

$$\int_{\partial\Omega} u(y) d\sigma_y K_1(r, \lambda) - \int_{\Omega} L_{*\lambda}[u](y) K_1(r, \lambda) dy = \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \quad (2.28)$$

Remark 2.3 In case of $\lambda = 0$, Theorems 2.5–2.8 are respectively the classical Cauchy-Pompeiu formula in Clifford analysis.

From Theorems 2.5–2.8, the following results can be directly proved.

Theorem 2.9 *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^1(\Omega, C(V_{3,3})) \cap C(\overline{\Omega}, C(V_{3,3}))$, $K_{*1}(r, \lambda)$ be as in (2.9), and $L_{\lambda}[u] = 0$ in Ω . Then*

$$\int_{\partial\Omega} K_{*1}(r, \lambda) d\sigma_y u(y) = \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \quad (2.29)$$

Theorem 2.10 *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^1(\Omega, C(V_{3,3})) \cap C(\overline{\Omega}, C(V_{3,3}))$, $K_1(r, \lambda)$ be as in (2.8), and $L_{-\lambda}[u] = 0$ in Ω . Then*

$$\int_{\partial\Omega} K_1(r, \lambda) d\sigma_y u(y) = \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \quad (2.30)$$

Theorem 2.11 *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^1(\Omega, C(V_{3,3})) \cap C(\overline{\Omega}, C(V_{3,3}))$, $K_{*1}(r, \lambda)$ be as in (2.9), and $L_{*-\lambda}[u] = 0$ in Ω . Then*

$$\int_{\partial\Omega} u(y) d\sigma_y K_{*1}(r, \lambda) = \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \quad (2.31)$$

Theorem 2.12 *Let Ω be an open bounded non empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^1(\Omega, C(V_{3,3})) \cap C(\overline{\Omega}, C(V_{3,3}))$, $K_1(r, \lambda)$ be as in (2.8), and $L_{*\lambda}[u] = 0$ in Ω , then*

$$\int_{\partial\Omega} u(y) d\sigma_y K_1(r, \lambda) = \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \quad (2.32)$$

Remark 2.4 In case of $\lambda = 0$, Theorems 2.9–2.12 are respectively the classical Cauchy integral formula in Clifford analysis.

Corollary 2.1 *Suppose $\|u\| = O(e^{\theta\|y\|})$ ($\|y\| \rightarrow +\infty$, $y \in \mathcal{R}^3$), $0 < \theta < |h|$ and denote the conditions as follows: (I) $Lu = 0$ in \mathcal{R}^3 ; (II) $L_*u = 0$ in \mathcal{R}^3 ; (III) $L_\lambda[u] = 0$ in \mathcal{R}^3 ; (IV) $L_{-\lambda}[u] = 0$ in \mathcal{R}^3 ; (V) $L_{*-\lambda}[u] = 0$ in \mathcal{R}^3 ; (VI) $L_{*\lambda}[u] = 0$ in \mathcal{R}^3 . If any of the above conditions is satisfied, then $u \equiv 0$ in \mathcal{R}^3 .*

Proof By Theorems 2.3, 2.4, 2.9–2.12, for any $x \in \mathcal{R}^3$, replacing $\partial\Omega$ by $\partial B(x, R)$ in the integral representations, taking limit $R \rightarrow \infty$, the result follows.

3 Generalized Higher Order Integral Representations

Denote

$$K_{m+1}^A(r, h) = \frac{1}{m!\omega_3} \left(\frac{(1-m)(\mathbf{y}-\mathbf{x})e_A}{r^{3-m}} + \frac{|h|(\mathbf{y}-\mathbf{x})e_A}{r^{2-m}} - \frac{me_A\hbar}{r^{2-m}} + \frac{e_A\hbar}{r^{1-m}} \right) e^{-|h|r} \hbar^m, \quad (3.1)$$

$$K_{*m+1}^A(r, h) = \frac{(-1)^m \hbar^m}{m!\omega_3} \left(\frac{(1-m)e_A(\mathbf{y}-\mathbf{x})}{r^{3-m}} + \frac{e_A|h|(\mathbf{y}-\mathbf{x})}{r^{2-m}} + \frac{m\hbar e_A}{r^{2-m}} - \frac{\hbar e_A}{r^{1-m}} \right) e^{-|h|r}, \quad (3.2)$$

$$K_{m+1}(r, \lambda) = \frac{1}{m!\omega_3} \left(\frac{(1-m)(\mathbf{y}-\mathbf{x})}{r^{3-m}} + \frac{\lambda(\mathbf{y}-\mathbf{x})}{r^{2-m}} - \frac{m}{r^{2-m}} + \frac{\lambda}{r^{1-m}} \right) e^{-\lambda r}, \quad (3.3)$$

$$K_{*m+1}(r, \lambda) = \frac{(-1)^m}{m!\omega_3} \left(\frac{(1-m)(\mathbf{y}-\mathbf{x})}{r^{3-m}} + \frac{\lambda(\mathbf{y}-\mathbf{x})}{r^{2-m}} + \frac{m}{r^{2-m}} - \frac{\lambda}{r^{1-m}} \right) e^{-\lambda r}, \quad (3.4)$$

where $\mathbf{y} - \mathbf{x} = \sum_{i=1}^3 (y_i - x_i)e_i$, $|h| = \lambda$, $h = |h|\hbar$, $r = |y - x|$, $m \geq 0$ and ω_3 denotes the area of the unit sphere in \mathcal{R}^3 .

Lemma 3.1 Let $K_{m+1}^A(r, h)$ and $K_{*m+1}^A(r, h)$ be as in (3.1) and (3.2), $m \geq 0$. Then

$$\begin{cases} L[K_{m+1}^A(r, h)] = K_m^A(r, h), & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}, \\ L_*[K_{*m+1}^A(r, h)] = K_{*m}^A(r, h), & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}. \end{cases} \quad (3.5)$$

Lemma 3.2 Let $K_{m+1}^A(r, h)$ and $K_{*m+1}^A(r, h)$ be as in (3.1) and (3.2), $m \geq 0$. Then

$$\begin{cases} L^{m+1}[K_{m+1}^A(r, h)] = 0, & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}, \\ L_*^{m+1}[K_{*m+1}^A(r, h)] = 0, & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}. \end{cases} \quad (3.6)$$

Lemma 3.3 Let $K_{m+1}(r, \lambda)$ and $K_{*m+1}(r, \lambda)$ be as in (3.3) and (3.4), $m \geq 0$. Then

$$\begin{cases} L_\lambda[K_{m+1}(r, \lambda)] = L_{*-\lambda}[K_{m+1}(r, \lambda)] = K_m(r, \lambda), & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}, \\ L_{-\lambda}[K_{*m+1}(r, \lambda)] = L_{*\lambda}[K_{*m+1}(r, \lambda)] = K_{*m}(r, \lambda), & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}. \end{cases} \quad (3.7)$$

Lemma 3.4 Let $K_{m+1}(r, \lambda)$ and $K_{*m+1}(r, \lambda)$ be as in (3.3) and (3.4), $m \geq 0$. Then

$$\begin{cases} L_\lambda^{m+1}[K_{m+1}(r, \lambda)] = L_{*-\lambda}^{m+1}[K_{m+1}(r, \lambda)] = 0, & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}, \\ L_{-\lambda}^{m+1}[K_{*m+1}(r, \lambda)] = L_{*\lambda}^{m+1}[K_{*m+1}(r, \lambda)] = 0, & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}. \end{cases} \quad (3.8)$$

Suppose $\lambda = 0$, then (3.3) and (3.4) can be rewritten as

$$K_{m+1}(r, 0) = \frac{1}{m!\omega_3} \left(\frac{(1-m)(\mathbf{y}-\mathbf{x})}{r^{3-m}} - \frac{m}{r^{2-m}} \right), \quad (3.9)$$

$$K_{*m+1}(r, 0) = \frac{(-1)^m}{m!\omega_3} \left(\frac{(1-m)(\mathbf{y}-\mathbf{x})}{r^{3-m}} + \frac{m}{r^{2-m}} \right). \quad (3.10)$$

Lemma 3.5 Let $K_{m+1}(r, 0)$ and $K_{*m+1}(r, 0)$ be as in (3.9) and (3.10), $m \geq 0$. Then

$$\begin{cases} D[K_{m+1}(r, 0)] = [K_{m+1}(r, 0)]D = K_m(r, 0), & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}, \\ D[K_{*m+1}(r, 0)] = [K_{*m+1}(r, 0)]D = K_{*m}(r, 0), & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}. \end{cases} \quad (3.11)$$

Lemma 3.6 Let $K_{m+1}(r, 0)$ and $K_{*m+1}(r, 0)$ be as in (3.9) and (3.10), $m \geq 0$. Then

$$\begin{cases} D^{m+1}[K_{m+1}(r, 0)] = [K_{m+1}(r, 0)]D^{m+1} = 0, & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}, \\ D^{m+1}[K_{*m+1}(r, 0)] = [K_{*m+1}(r, 0)]D^{m+1} = 0, & y \in \mathcal{R}_x^3 = \mathcal{R}^3 \setminus \{x\}. \end{cases} \quad (3.12)$$

Remark 3.1 By Lemma 3.6, $K_{m+1}(r, 0)$ and $K_{*m+1}(r, 0)$ are the higher order kernel functions. Comparing the forms with the results in [7, 29–31], these kernel functions are different. In case of $m = 0$, $K_{m+1}(r, 0)$ and $K_{*m+1}(r, 0)$ are the classical kernel functions in Clifford analysis. Thus, (3.3) and (3.4) also provide a way to find the kernel functions for $(m+1)$ -regular functions.

Remark 3.2 Denote $H_j^*(x)$, $j \geq 1$ as in [7, 29–31]. Then we have

$$H_j^*(y-x) = \frac{K_j(r, 0) + K_{*j}(r, 0)}{2}, \quad j \geq 1. \quad (3.13)$$

Theorem 3.1 *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^k(\Omega, C(V_{3,3})) \cap C^{k-1}(\overline{\Omega}, C(V_{3,3}))$. Then for $x \in \Omega$,*

$$\begin{aligned} u(x) = & \frac{1}{\omega_3} \left[\sum_{j=1}^k \frac{1}{(j-1)!} \int_{\partial\Omega} \left[\frac{(2-j)(\mathbf{y}-\mathbf{x})}{r^{4-j}} + \frac{|h|(\mathbf{y}-\mathbf{x})}{r^{3-j}} \right] e^{-|h|r} d\sigma_y L^{j-1}[u](y) \hbar^{j-1} \right. \\ & + \sum_{j=1}^k \frac{1}{(j-1)!} \int_{\partial\Omega} \left[\frac{j-1}{r^{3-j}} - \frac{|h|}{r^{2-j}} \right] e^{-|h|r} d\sigma_y L^{j-1}[u](y) \hbar^j \Big] \\ & - \frac{1}{(k-1)!\omega_3} \int_{\Omega} \left[\frac{(2-k)(\mathbf{y}-\mathbf{x})}{r^{4-k}} + \frac{|h|(\mathbf{y}-\mathbf{x})}{r^{3-k}} \right] e^{-|h|r} L^k[u](y) \hbar^{k-1} dy \\ & - \frac{1}{(k-1)!\omega_3} \int_{\Omega} \left[\frac{k-1}{r^{3-k}} - \frac{|h|}{r^{2-k}} \right] e^{-|h|r} L^k[u](y) \hbar^k dy. \end{aligned} \quad (3.14)$$

Proof For $x \in \Omega$, by Lemmas 2.3, 3.1 and 3.2, we have

$$\begin{aligned} & \operatorname{Re} \left\{ \sum_{j=1}^k (-1)^{j-1} \int_{\partial\Omega \setminus \partial B(x, \varepsilon)} K_{*j}^A(r, h) d\sigma_y L^{j-1}[u](y) \right\} \\ & = \operatorname{Re} \left\{ (-1)^{k-1} \int_{\Omega \setminus B(x, \varepsilon)} K_{*k}^A(r, h) L^k[u] dy \right\}. \end{aligned} \quad (3.15)$$

In view of $u \in C^k(\Omega, C(V_{3,3})) \cap C^{k-1}(\overline{\Omega}, C(V_{3,3}))$, by Stokes formula, it can be proved that

$$\lim_{\varepsilon \rightarrow 0} \operatorname{Re} \left\{ \int_{\partial B(x, \varepsilon)} K_{*j}^A(r, h) d\sigma_y L^{j-1}[u](y) \right\} = 0, \quad 2 \leq j \leq k. \quad (3.16)$$

Taking limit $\varepsilon \rightarrow 0$ in (3.15), in view of the weak singularity of $K_{*j}^A(r, h)$ ($j = 1, \dots, k$), by (2.16)–(2.18) and (3.16), we have

$$\begin{aligned} & \operatorname{Re} \left\{ \sum_{j=1}^k (-1)^{j-1} \int_{\partial\Omega} K_{*j}^A(r, h) d\sigma_y L^{j-1}[u](y) \right\} \\ & = u_A(x) e_A^2 + \operatorname{Re} (-1)^{k-1} \int_{\Omega} K_{*k}^A(r, h) L^k[u] dy. \end{aligned} \quad (3.17)$$

Combining (3.2) with (3.17), by the same technique as in (2.21), (3.14) holds.

By Lemmas 2.3, 3.1 and 3.2, similarly, we have the following theorem.

Theorem 3.2 *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^k(\Omega, C(V_{3,3})) \cap C^{k-1}(\overline{\Omega}, C(V_{3,3}))$. Then for $x \in \Omega$,*

$$\begin{aligned} u(x) = & \frac{1}{\omega_3} \left[\sum_{j=1}^k \frac{(-1)^{j-1}}{(j-1)!} \int_{\partial\Omega} \hbar^{j-1} L_*^{j-1}[u](y) d\sigma_y \left[\frac{(2-j)(\mathbf{y}-\mathbf{x})}{r^{4-j}} + \frac{|h|(\mathbf{y}-\mathbf{x})}{r^{3-j}} \right] e^{-|h|r} \right. \\ & - \sum_{j=1}^k \frac{(-1)^{j-1}}{(j-1)!} \int_{\partial\Omega} \hbar^j L_*^{j-1}[u](y) d\sigma_y \left[\frac{j-1}{r^{3-j}} - \frac{|h|}{r^{2-j}} \right] e^{-|h|r} \Big] \\ & - \frac{(-1)^{k-1}}{(k-1)!\omega_3} \int_{\Omega} \hbar^{k-1} L_*^k[u](y) \left[\frac{(2-k)(\mathbf{y}-\mathbf{x})}{r^{4-k}} + \frac{|h|(\mathbf{y}-\mathbf{x})}{r^{3-k}} \right] e^{-|h|r} dy \end{aligned}$$

$$- \frac{(-1)^{k-1}}{(k-1)! \omega_3} \int_{\Omega} \hbar^k L_*^k[u](y) \left[\frac{1-k}{r^{3-k}} + \frac{|h|}{r^{2-k}} \right] e^{-|h|r} dy. \quad (3.18)$$

Remark 3.3 In case of $k = 1$, Theorems 3.1 and 3.2 are just Theorems 2.1 and 2.2. Thus we also call Theorems 3.1 and 3.2 the generalized higher order Cauchy-Pompeiu formulas.

By Theorems 3.1 and 3.2, we have the following result.

Theorem 3.3 Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^k(\Omega, C(V_{3,3})) \cap C^{k-1}(\overline{\Omega}, C(V_{3,3}))$, $L^k u = 0$ in Ω . Then for $x \in \Omega$,

$$\begin{aligned} u(x) = & \frac{1}{\omega_3} \left[\sum_{j=1}^k \frac{1}{(j-1)!} \int_{\partial\Omega} \left[\frac{(2-j)(\mathbf{y}-\mathbf{x})}{r^{4-j}} + \frac{|h|(\mathbf{y}-\mathbf{x})}{r^{3-j}} \right] e^{-|h|r} d\sigma_y L^{j-1}[u](y) \hbar^{j-1} \right. \\ & \left. + \sum_{j=1}^k \frac{1}{(j-1)!} \int_{\partial\Omega} \left[\frac{j-1}{r^{3-j}} - \frac{|h|}{r^{2-j}} \right] e^{-|h|r} d\sigma_y L^{j-1}[u](y) \hbar^j \right]. \end{aligned} \quad (3.19)$$

Theorem 3.4 Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^k(\Omega, C(V_{3,3})) \cap C^{k-1}(\overline{\Omega}, C(V_{3,3}))$, $L_*^k u = 0$ in Ω . Then for $x \in \Omega$,

$$\begin{aligned} u(x) = & \frac{1}{\omega_3} \left[\sum_{j=1}^k \frac{(-1)^{j-1}}{(j-1)!} \int_{\partial\Omega} \hbar^{j-1} L_*^{j-1}[u](y) d\sigma_y \left[\frac{(2-j)(\mathbf{y}-\mathbf{x})}{r^{4-j}} + \frac{|h|(\mathbf{y}-\mathbf{x})}{r^{3-j}} \right] e^{-|h|r} \right. \\ & \left. - \sum_{j=1}^k \frac{(-1)^{j-1}}{(j-1)!} \int_{\partial\Omega} \hbar^j L_*^{j-1}[u](y) d\sigma_y \left[\frac{j-1}{r^{3-j}} - \frac{|h|}{r^{2-j}} \right] e^{-|h|r} \right]. \end{aligned} \quad (3.20)$$

Remark 3.4 In case of $k = 1$, Theorems 3.3 and 3.4 are just Theorems 2.3 and 2.4. Thus we also call Theorems 3.3 and 3.4 the generalized higher order Cauchy integral formulas.

By Lemmas 2.4, 3.3 and 3.4, we have the following result.

Theorem 3.5 Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^k(\Omega, C(V_{3,3})) \cap C^{k-1}(\overline{\Omega}, C(V_{3,3}))$, $K_{*j}(r, \lambda)$, $j = 1, \dots, k$ be as in (3.4). Then

$$\begin{aligned} & \sum_{j=1}^k (-1)^{j-1} \int_{\partial\Omega} K_{*j}(r, \lambda) d\sigma_y L_{\lambda}^{j-1}[u](y) + (-1)^k \int_{\Omega} K_{*k}(r, \lambda) L_{\lambda}^k[u](y) dy \\ = & \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \end{aligned} \quad (3.21)$$

Theorem 3.6 Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^k(\Omega, C(V_{3,3})) \cap C^{k-1}(\overline{\Omega}, C(V_{3,3}))$, $K_j(r, \lambda)$, $j = 1, \dots, k$ be as in (3.3). Then

$$\begin{aligned} & \sum_{j=1}^k (-1)^{j-1} \int_{\partial\Omega} K_j(r, \lambda) d\sigma_y L_{-\lambda}^{j-1}[u](y) + (-1)^k \int_{\Omega} K_k(r, \lambda) L_{-\lambda}^k[u](y) dy \\ = & \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \end{aligned} \quad (3.22)$$

Theorem 3.7 *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^k(\Omega, C(V_{3,3})) \cap C^{k-1}(\overline{\Omega}, C(V_{3,3}))$, $K_{*j}(r, \lambda)$, $j = 1, \dots, k$ be as in (3.4). Then*

$$\begin{aligned} & \sum_{j=1}^k (-1)^{j-1} \int_{\partial\Omega} L_{*-\lambda}^{j-1}[u](y) d\sigma_y K_{*j}(r, \lambda) + (-1)^k \int_{\Omega} L_{*-\lambda}^k[u](y) K_{*k}(r, \lambda) dy \\ &= \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \end{aligned} \quad (3.23)$$

Theorem 3.8 *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^k(\Omega, C(V_{3,3})) \cap C^{k-1}(\overline{\Omega}, C(V_{3,3}))$, $K_j(r, \lambda)$, $j = 1, \dots, k$ be as in (3.3). Then*

$$\begin{aligned} & \sum_{j=1}^k (-1)^{j-1} \int_{\partial\Omega} L_{*\lambda}^{j-1}[u](y) d\sigma_y K_j(r, \lambda) + (-1)^k \int_{\Omega} L_{*\lambda}^k[u](y) K_k(r, \lambda) dy \\ &= \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \end{aligned} \quad (3.24)$$

Remark 3.5 In case of $k = 1$, Theorems 3.5–3.8 are just Theorems 2.5–2.8.

Remark 3.6 In case of $\lambda = 0$, Theorems 3.5–3.8 are the other expressions of the higher order Cauchy-Pompeiu formulas which are different from the results in [7, 29–31].

By Theorems 3.5–3.8, we have the following results.

Theorem 3.9 *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^k(\Omega, C(V_{3,3})) \cap C^{k-1}(\overline{\Omega}, C(V_{3,3}))$, $K_{*j}(r, \lambda)$, $j = 1, \dots, k$ be as in (3.4), $L_{*\lambda}^k u = 0$ in Ω . Then*

$$\sum_{j=1}^k (-1)^{j-1} \int_{\partial\Omega} K_{*j}(r, \lambda) d\sigma_y L_{*\lambda}^{j-1}[u](y) = \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \quad (3.25)$$

Theorem 3.10 *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^k(\Omega, C(V_{3,3})) \cap C^{k-1}(\overline{\Omega}, C(V_{3,3}))$, $K_j(r, \lambda)$, $j = 1, \dots, k$ be as in (3.3), $L_{-\lambda}^k u = 0$ in Ω . Then*

$$\sum_{j=1}^k (-1)^{j-1} \int_{\partial\Omega} K_j(r, \lambda) d\sigma_y L_{-\lambda}^{j-1}[u](y) = \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \quad (3.26)$$

Theorem 3.11 *Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^k(\Omega, C(V_{3,3})) \cap C^{k-1}(\overline{\Omega}, C(V_{3,3}))$, $K_{*j}(r, \lambda)$, $j = 1, \dots, k$ be as in (3.4), $L_{*-\lambda}^k u = 0$ in Ω . Then*

$$\sum_{j=1}^k (-1)^{j-1} \int_{\partial\Omega} L_{*-\lambda}^{j-1}[u](y) d\sigma_y K_{*j}(r, \lambda) = \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \quad (3.27)$$

Theorem 3.12 Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^k(\Omega, C(V_{3,3})) \cap C^{k-1}(\overline{\Omega}, C(V_{3,3}))$, $K_j(r, \lambda)$, $j = 1, \dots, k$ be as in (3.3), $L_*^k u = 0$. Then

$$\sum_{j=1}^k (-1)^{j-1} \int_{\partial\Omega} L_*^{j-1}[u](y) d\sigma_y K_j(r, \lambda) = \begin{cases} u(x), & x \in \Omega, \\ 0, & x \in \mathcal{R}^3 \setminus \overline{\Omega}. \end{cases} \quad (3.28)$$

Remark 3.7 In case of $k = 1$, Theorems 3.9–3.12 are just Theorems 2.9–2.12.

Remark 3.8 In case of $\lambda = 0$, Theorems 3.9–3.12 are the other expressions of the higher order Cauchy integral formulas which are different from the results in [7, 29–31].

4 Maximum Modulus Theorem

In this section, suppose that Ω is a bounded domain in \mathcal{R}^3 , $Hu = 0$ in Ω . We shall give the Maximum Modulus Theorem for u .

Lemma 4.1 Let $u, v \in C^2(\Omega, C(V_{3,3})) \cap C^1(\overline{\Omega}, C(V_{3,3}))$. Then

$$\int_{\Omega} (vH[u] - H[v]u) dy = \int_{\partial\Omega} v d\sigma_y L_\lambda[u] - \int_{\partial\Omega} L_{*- \lambda}[v] d\sigma_y u. \quad (4.1)$$

Remark 4.1 In case of $\lambda = 0$, Lemma 4.1 is just the following well-known Green formula in Clifford analysis:

$$\int_{\Omega} (v\Delta[u] - \Delta[v]u) dy = \int_{\partial\Omega} v d\sigma_y D[u] - \int_{\partial\Omega} [v] D d\sigma_y u. \quad (4.2)$$

Lemma 4.2 Let Ω be a bounded domain in \mathcal{R}^3 , $u \in C^2(\Omega, C(V_{3,3}))$, $Hu = 0$ in Ω and $|u(x)| = \text{constant}$ in Ω . We have

- (I) If $\lambda > 0$, then $u(x) \equiv 0$ in Ω ,
- (II) If $\lambda = 0$, then $u(x)$ must be constant.

Proof Denote $u(x) = \sum_A u_A(x) e_A$. Then $\sum_A u_A^2(x) \equiv C$, from which we have

$$\sum_{i=1}^3 \sum_A \left(\frac{\partial u_A}{\partial x_i} \right)^2 + \sum_A u_A(x) \Delta[u_A] = 0. \quad (4.3)$$

In view of $Hu = 0$ in Ω , thus $Hu_A = 0$ in Ω . Then more clearly,

$$\Delta[u_A] = \lambda^2 u_A. \quad (4.4)$$

Combining (4.3) with (4.4), we have

$$\sum_{i=1}^3 \sum_A \left(\frac{\partial u_A}{\partial x_i} \right)^2 + \sum_A \lambda^2 u_A^2(x) = 0. \quad (4.5)$$

Obviously, if $\lambda > 0$, then $u_A(x) \equiv 0$. Thus $u(x) \equiv 0$.

If $\lambda = 0$, then

$$\frac{\partial u_A(x)}{\partial x_i} = 0, \quad i = 1, 2, 3, \quad A \in \mathcal{PN}, \quad N = \{1, \dots, 3\}, \quad x \in \Omega. \quad (4.6)$$

Consequently, $u(x)$ is a constant.

Remark 4.2 Lemma 4.2 is still valid in \mathcal{R}^3 .

Remark 4.3 By Lemma 4.2, suppose that Ω is a bounded domain in \mathcal{R}^3 , $u \in C^2(\Omega, C(V_{3,3}))$, $|u(x)| = \text{constant}$ in Ω . Denote the conditions as follows: (I) $Lu = 0$ in Ω ; (II) $L_*u = 0$ in Ω ; (III) $L_\lambda[u] = 0$ in Ω ; (IV) $L_{-\lambda}[u] = 0$ in Ω ; (V) $L_{*-\lambda}[u] = 0$ in Ω ; (VI) $L_{*\lambda}[u] = 0$ in Ω ; (VII) $D[u] = 0$ in Ω ; (VIII) $[u]D = 0$ in Ω . If any of the above conditions is satisfied, then $u(x) \equiv C$.

Theorem 4.1 Let Ω be an open bounded non-empty subset of \mathcal{R}^3 with a Liapunov boundary $\partial\Omega$, $u \in C^2(\Omega, C(V_{3,3})) \cap C^1(\overline{\Omega}, C(V_{3,3}))$, and $Hu = 0$ in Ω . Then for $x \in \Omega$,

$$u(x) = \frac{1}{\omega_3} \int_{\partial\Omega} \left(\frac{\mathbf{y} - \mathbf{x}}{r^3} + \frac{\lambda(\mathbf{y} - \mathbf{x})}{r^2} - \frac{\lambda}{r} \right) e^{-\lambda r} d\sigma_y u(y) + \frac{1}{\omega_3} \int_{\partial\Omega} \frac{e^{-\lambda r}}{r} d\sigma_y L_\lambda[u](y). \quad (4.7)$$

Proof By Lemma 4.1, taking $v = -\frac{1}{\omega_3} \frac{e^{-\lambda r}}{r}$ in (4.1), obviously, we get $L_{*-\lambda}[v] = K_{*1}(r, \lambda)$. Thus for $x \in \Omega$, we have

$$-\frac{1}{\omega_3} \int_{\partial(\Omega \setminus B(x, \delta))} \frac{e^{-\lambda r}}{r} d\sigma_y L_\lambda[u](y) - \int_{\partial(\Omega \setminus B(x, \delta))} K_{*1}(r, \lambda) d\sigma_y u = 0. \quad (4.8)$$

It is easy to check that

$$\lim_{\delta \rightarrow 0} \left[-\frac{1}{\omega_3} \int_{\partial B(x, \delta)} \frac{e^{-\lambda r}}{r} d\sigma_y L_\lambda[u](y) - \int_{\partial B(x, \delta)} K_{*1}(r, \lambda) d\sigma_y u \right] = -u(x). \quad (4.9)$$

Combining (4.8) with (4.9), (4.7) follows.

Corollary 4.1 Let Ω be a bounded domain in \mathcal{R}^3 , $u \in C^2(\Omega, C(V_{3,3}))$ and $Hu = 0$ in Ω . Then for any $x_0 \in \Omega$,

$$u(x_0) = \frac{1}{\omega_3} \left(\frac{1}{R^2} + \frac{\lambda}{R} \right) e^{-\lambda R} \int_{\partial B(x_0, R)} u(y) dS + \frac{\lambda^2 e^{-\lambda R}}{\omega_3 R} \int_{B(x_0, R)} u(y) dy, \quad (4.10)$$

where R is chosen such that $B(x_0, R) \subset \Omega$.

Remark 4.4 In case of $\lambda = 0$, Corollary 4.1 is just the mean value theorem for harmonic functions.

Remark 4.5 Corollary 4.1 is still valid for the functions satisfying any of the following conditions: (I) $Lu = 0$ in Ω ; (II) $L_*u = 0$ in Ω ; (III) $L_\lambda[u] = 0$ in Ω ; (IV) $L_{-\lambda}[u] = 0$ in Ω ; (V) $L_{*-\lambda}[u] = 0$ in Ω ; (VI) $L_{*\lambda}[u] = 0$ in Ω ; (VII) $D[u] = 0$ in Ω ; (VIII) $[u]D = 0$ in Ω .

Theorem 4.2 (Maximum Modulus Theorem) Suppose that Ω is a bounded domain in \mathcal{R}^3 , $u \in C^2(\Omega, C(V_{3,3}))$ and $Hu = 0$ in Ω . If there exists an $x_0 \in \Omega$, such that

$$|u(x_0)| \geq |u(x)|, \quad x \in \Omega, \quad (4.11)$$

then $u(x)$ is a constant. Moreover, if $\lambda > 0$, then $u(x) \equiv 0$.

Proof (I) $\lambda = 0$, by the mean value theorem of harmonic functions and Lemma 4.2, the result follows.

(II) $\lambda > 0$, suppose $|u(x_0)| > 0$. Taking $R > 0$ such that $B(x_0, R) \subset \Omega$, by Corollary 4.1, we have

$$|u(x_0)| \leq \frac{1}{\omega_3} \left(\frac{1}{R^2} + \frac{\lambda}{R} \right) e^{-\lambda R} |u(x_0)| R^2 \omega_3 + \frac{\lambda^2 e^{-\lambda R}}{\omega_3 R} |u(x_0)| R^3 V_3, \quad (4.12)$$

more clearly,

$$|u(x_0)| \leq \frac{1 + \lambda R + \frac{(\lambda R)^2}{3}}{e^{\lambda R}} |u(x_0)| < |u(x_0)|. \quad (4.13)$$

(4.13) implies $|u(x_0)| = 0$. Thus the result follows.

Corollary 4.2 *Suppose that Ω is a bounded domain in \mathcal{R}^3 , $u \in C^2(\Omega, C(V_{3,3}))$, $Hu = 0$ in Ω and continuous on $\overline{\Omega}$. Then*

$$\max\{|u(x)|, x \in \overline{\Omega}\} = \max\{|u(x)|, x \in \partial\Omega\}. \quad (4.14)$$

Remark 4.6 Theorem 4.2 and Corollary 4.2 are still valid for the functions satisfying any of the following conditions: (I) $Lu = 0$ in Ω ; (II) $L_*u = 0$ in Ω ; (III) $L_\lambda[u] = 0$ in Ω ; (IV) $L_{-\lambda}[u] = 0$ in Ω ; (V) $L_{*-\lambda}[u] = 0$ in Ω ; (VI) $L_{*\lambda}[u] = 0$ in Ω ; (VII) $D[u] = 0$ in Ω , (VIII) $[u]D = 0$ in Ω .

Acknowledgement The authors are very grateful to the reviewers for their helpful suggestions.

References

- [1] de Almeida, R. and Kraußhar, R. S., On the asymptotic growth of entire monogenic functions, *J. Anal. Appl.*, **24**, 2005, 791–813.
- [2] Begehr, H., Iterations of Pompeiu operators, *Mem. Diff. Eq. Math. Phys.*, **12**, 1997, 3–21.
- [3] Begehr, H., Iterated integral operators in Clifford analysis, *J. Anal. Appl.*, **18**, 1999, 361–377.
- [4] Begehr, H., Representation formulas in Clifford analysis. Acoustics, Mechanics, and the Related Topics of Mathematical Analysis, World Scientific, Singapore, 2002, 8–13.
- [5] Begehr, H., Dai, D. Q. and Li, X., Integral representation formulas in polydomains, *Complex Variables, Theory Appl.*, **47**, 2002, 463–484.
- [6] Begehr, H. Zhang, Z. X. and Du, J. Y., On Cauchy-Pompeiu formula for functions with values in a universal Clifford algebra, *Acta Math. Sci.*, **23B**(1), 2003, 95–103.
- [7] Begehr, H., Du, J. Y. and Zhang, Z. X., On higher order Cauchy-Pompeiu formula in Clifford analysis and its applications, *General Math.*, **11**, 2003, 5–26.
- [8] Begehr, H. Zhang, Z. X. and Vu, T. N. H., Generalized integral representations in Clifford analysis, *Complex Var., Elliptic Eqns.*, **51**, 2006, 745–762.
- [9] Brackx, F., Delanghe, R. and Sommen, F., Clifford Analysis, Research Notes in Mathematics, **76**, Pitman Books Ltd, London, 1982.
- [10] Constaes, D., de Almeida, R. and Kraußhar, R. S., On Cauchy estimates and growth orders of entire solutions of iterated Dirac and generalized Cauchy-Riemann equations, *Math. Meth. Appl. Sci.*, **29**, 2006, 1663–1686.
- [11] Delanghe, R., On regular analytic functions with values in a Clifford algebra, *Math. Ann.*, **185**, 1970, 91–111.
- [12] Delanghe, R., On the singularities of functions with values in a Clifford algebra, *Math. Ann.*, **196**, 1972, 293–319.
- [13] Delanghe, R., Morera’s theorem for functions with values in a Clifford algebra, *Simon Stevin*, **43**, 1969–1970, 129–140.
- [14] Delanghe, R. and Brackx, F., Hypercomplex function theory and Hilbert modules with reproducing kernel, *Proc. London Math. Soc.*, **37**, 1978, 545–576.
- [15] Du, J. Y. and Zhang, Z. X., A Cauchy’s integral formula for functions with values in a universal Clifford algebra and its applications, *Complex Var. Theory Appl.*, **47**, 2002, 915–928.
- [16] Franks, E. and Ryan, J., Bounded monogenic functions on unbounded domains, *Contemp. Math.*, **212**, 1998, 71–80.

- [17] Gürlebeck, K., On some operators in Clifford analysis, *Contemp. Math.*, **212**, 1998, 95–107.
- [18] Gürlebeck, K., Kähler, U., Ryan, J., et al., Clifford analysis over unbounded domains, *Adv. Appl. Math.*, **2**(19), 1997, 216–239.
- [19] Gürlebeck, K. and Sprössig, W., Quaternionic Analysis and Elliptic Boundary Value Problems, Akademie-Verlag, Berlin, 1989.
- [20] Gürlebeck, K. and Zhang, Z. X., Some Riemann boundary value problems in Clifford analysis, *Math. Meth. Appl. Sci.*, **33**, 2010, 287–302.
- [21] Iftimie, V., Functions hypercomplex, *Bull. Math. Soc. Sci. Math. R. S. Romania.*, **9**(57), 1965, 279–332.
- [22] Lu, J. K., Boundary value problems of analytic functions, World Scientific, Singapore, 1993.
- [23] Muskhelishvili, N. I., Singular Integral Equations, NauKa, Moscow, 1968.
- [24] Obolashvili, E., Higher order partial differential equations in Clifford analysis, Birkhauser, Boston, Basel, Berlin, 2002.
- [25] Vu, T. N. H., Integral representations in quaternionic analysis related to Helmholtz operator, *Comp. Var. Theory Appl.*, **12**(48), 2003, 1005–1021.
- [26] Vu, T. N. H., Helmholtz Operator in Quaternionic Analysis, Ph. D. Thesis, Free University, Berlin, 2005.
- [27] Vekua, I. N., Generalized Analytic Functions, Pergamon Press, Oxford, 1962.
- [28] Xu, Z. Y., Boundary Value Problems and Function Theory for Spin-invariant Differential Operators, Ph. D. Thesis, Ghent State University, 1989.
- [29] Zhang, Z. X., Integral representations and its applications in Clifford analysis, *Gen. Math. (in Romania)*, **13**(3), 2005, 79–96.
- [30] Zhang, Z. X., On k -regular functions with values in a universal Clifford algebra, *J. Math. Anal. Appl.*, **315**(2), 2006, 491–505.
- [31] Zhang, Z. X., Some properties of operators in Clifford analysis, *Complex Var., Elliptic Eqns.*, **6**(52), 2007, 455–473.
- [32] Zhang, Z. X., On higher order Cauchy-Pompeiu formula for functions with values in a universal Clifford algebra, *Bull. Belg. Math. Soc.*, **14**, 2007, 87–97.