

Essential Norms of Composition Operators Between Hardy Spaces of the Unit Disc*

Luo LUO¹ Kai LI¹

Abstract The authors express the essential norms of composition operators between Hardy spaces of the unit disc in terms of the natural Nevanlinna counting function.

Keywords Essential norm, Composition operators, Hardy spaces

2000 MR Subject Classification 32A35, 47B33

1 Introduction

Let D be the open unit disk in the complex plane and $H(D)$ denote the space of all holomorphic functions in D . For each p ($0 < p < \infty$), the Hardy space $H^p(D)$ is defined by

$$H^p(D) = \left\{ f \in H(D) : \sup_{0 < r < 1} \int_{\partial D} |f(r\xi)|^p d\sigma(\xi) < \infty \right\}, \quad \|f\|_p = \left[\int_{\partial D} |f^*(\xi)|^p d\sigma(\xi) \right]^{\frac{1}{p}},$$

where f^* denotes the radial limit of f and $d\sigma$ is the normalized Lebesgue measure on the boundary ∂D of D . For $1 < p < \infty$, the Hardy space $H^p(D)$ is a Banach space.

Let $\varphi : D \rightarrow D$ be a holomorphic self-map of D . For a holomorphic function f on D , denote the composition $f \circ \varphi$ by $C_\varphi f$ and call C_φ the composition operator induced by φ .

Let X and Y be Banach spaces. For a bounded linear operator $T : X \rightarrow Y$, the essential norm $\|T\|_{e, X \rightarrow Y}$ is defined to be the distance from T to the set of the compact operators $K : X \rightarrow Y$, namely,

$$\|T\|_{e, X \rightarrow Y} = \inf \{ \|T - K\| : K \text{ is compact from } X \text{ into } Y \},$$

where $\|\cdot\|$ denotes the usual operator norm.

J. H. Shapiro [3] expressed the essential norm of the composition operator $C_\varphi : H^2(D) \rightarrow H^2(D)$ in terms of natural Nevanlinna counting function of the inducing map φ .

The natural Nevanlinna counting function for φ , N_φ , provides such a measure. It is defined by

$$N_\varphi(w) = \sum_{z \in \varphi^{-1}\{w\}} \log \left(\frac{1}{|z|} \right), \quad w \in D \setminus \{\varphi(0)\}.$$

As usual, $z \in \varphi^{-1}\{w\}$ is repeated according to the multiplicity of the zero of $\varphi - w$ at z .

Manuscript received February 3, 2010. Revised October 1, 2010. Published online January 25, 2011.

¹Department of Mathematics, University of Science and Technology of China, Hefei 230026, China.

E-mail: lluo@ustc.edu.cn awaypa@mail.ustc.edu.cn

*Project supported by the National Natural Science Foundation of China (No. 10771201) and the Anhui Provincial Natural Science Foundation of China (No. 090416233).

The main goal of this paper is to compute the essential norm of $C_\varphi : H^p(D) \rightarrow H^q(D)$ for $1 < p \leq q < \infty$ in terms of the natural Nevanlinna counting function of the inducing map φ .

In this paper, we get the following theorem.

Theorem 1.1 *Let φ be a holomorphic self-map of D , $1 < p \leq q < \infty$. If $C_\varphi : H^p(D) \rightarrow H^q(D)$ is bounded, then there exist constants C_1 and C_2 , such that*

$$C_1 \limsup_{|a| \rightarrow 1^-} \frac{N_\varphi(a)}{[\log(\frac{1}{|a|})]^{\frac{q}{p}}} \leq \|C_\varphi\|_{e, H^p \rightarrow H^q}^q \leq C_2 \limsup_{|a| \rightarrow 1^-} \frac{N_\varphi(a)}{[\log(\frac{1}{|a|})]^{\frac{q}{p}}}.$$

Particularly, we get the corollary.

Corollary 1.1 For $1 < p \leq q < \infty$, $C_\varphi : H^p(D) \rightarrow H^q(D)$ is compact if and only if

$$\limsup_{|a| \rightarrow 1^-} \frac{N_\varphi(a)}{[\log(\frac{1}{|a|})]^{\frac{q}{p}}} = 0.$$

In the case $p = q = 2$, Theorem 1.1 and Corollary 1.1 were given by J. H. Shapiro [3].

Throughout the paper, C denotes a positive constant, whose value may change from one occurrence to the next one, but it is independent of f and φ .

2 Proof of Theorem 1.1

Recall that a holomorphic function f in D has the Taylor expansion

$$f(z) = \sum_{k=0}^{\infty} a_k z^k.$$

For the Taylor expansion of f and any integer $n \geq 1$, let

$$R_n f(z) = \sum_{k=n}^{\infty} a_k z^k$$

and $K_n = I - R_n$ where $I f = f$ is the identity operator.

The operator K_n has a connection with the following natural question: When does the partial sums of the Taylor expansion of f converge to f in the norm topology of the function space? K. Zhu [5] considered the question for various analytic function spaces on the unit disc. In order to prove our main result, we need some of his results.

Lemma 2.1 *Suppose that X is a Banach space of holomorphic functions in D with the property that the polynomials are dense in X . Then $\|K_n f - f\|_X \rightarrow 0$ as $n \rightarrow \infty$ if and only if $\sup\{\|K_n\| : n \geq 1\} < \infty$.*

Lemma 2.2 *If $1 < p < \infty$, then $\|K_n f - f\|_p \rightarrow 0$ as $n \rightarrow \infty$ for each $f \in H^p(D)$. Moreover, $\sup\{\|R_n\| : n \geq 1\} < \infty$ and $\sup\{\|K_n\| : n \geq 1\} < \infty$.*

Lemmas 2.1 and 2.2 are Proposition 1 in [5].

To prove Theorem 1.1, we also need the following lemmas.

Lemma 2.3 For $0 < p < \infty$, $f \in H(D)$ and φ is a holomorphic self-map of D . Then

$$\|f \circ \varphi\|_p^p = |f(\varphi(0))|^p + \frac{p^2}{2} \int_D |f(w)|^{p-2} |f'(w)|^2 N_\varphi(w) dA(w),$$

where dA is the normalized Lebesgue measure on D .

Lemma 2.3 is the special case of Lemma 2.2 (see the Change of Variable Formula and (2.1) in [4]).

Lemma 2.4 Let ψ be a holomorphic self-map of D . If $\psi(0) \neq 0$ and $0 < r < |\psi(0)|$, then

$$N_\psi(0) \leq \frac{1}{r^2} \int_{rD} N_\psi(w) dA(w).$$

Lemma 2.4 is the special case of Lemma 4.1 in [4].

Lemma 2.5 Let ψ be a holomorphic self-map of D . Let $a \in D$ and let

$$\sigma_a(w) = \frac{a - w}{1 - \bar{a}w}$$

be the Möbius self-map of D that interchanges 0 and a . Then

$$N_\psi \circ \sigma_a = N_{\sigma_a \circ \psi}.$$

Lemma 2.5 is the special case of Lemma 4.2 in [4].

Lemma 2.6 For $0 < p < \infty$, we have $f \in H^p(D)$ and $w \in D$. Then

$$|f(w)| \leq \frac{C \|f\|_p}{(1 - |w|)^{\frac{1}{p}}}.$$

Here C is independent of f .

Lemma 2.6 is the special case of Lemma 2.5 in [4].

Proof of Theorem 1.1 At first, we prove

$$\|C_\varphi\|_{e, H^p \rightarrow H^q}^q \geq C_1 \limsup_{|a| \rightarrow 1^-} \frac{N_\varphi(a)}{[\log(\frac{1}{|a|})]^{\frac{q}{p}}}.$$

For $a \in D$, letting

$$k_a(z) = \left\{ \frac{1 - |a|^2}{(1 - \bar{a}z)^2} \right\}^{\frac{1}{p}},$$

we know $\|k_a\|_p = 1$ and, as $|a| \rightarrow 1^-$, $k_a \rightarrow 0$ uniformly on compact subset of D .

For the moment, fix a compact operator $K : H^p(D) \rightarrow H^q(D)$. Since the family $\{k_a\}$ is bounded in $H^p(D)$, and $k_a \rightarrow 0$ uniformly on compact subsets of D as $|a| \rightarrow 1^-$, we have $\|Kk_a\|_q \rightarrow 0$, as $|a| \rightarrow 1^-$. Thus

$$\|C_\varphi - K\| \geq \limsup_{|a| \rightarrow 1^-} \|(C_\varphi - K)k_a\|_q \geq \limsup_{|a| \rightarrow 1^-} (\|C_\varphi k_a\|_q - \|Kk_a\|_q) = \limsup_{|a| \rightarrow 1^-} \|C_\varphi k_a\|_q.$$

Upon taking the infimum of both sides of this inequality over all compact operators $K : H^p(D) \rightarrow H^q(D)$, we obtain

$$\|C_\varphi\|_{e, H^p \rightarrow H^q} \geq \limsup_{|a| \rightarrow 1^-} \|C_\varphi k_a\|_q. \quad (2.1)$$

By Lemma 2.3,

$$\|C_\varphi k_a\|_q^q = |k_a(\varphi(0))|^q + \frac{q^2}{2} \int_D |k_a(w)|^{q-2} |k'_a(w)|^2 N_\varphi(w) dA(w).$$

So, there is a constant C such that

$$\begin{aligned} \|C_\varphi k_a\|_q^q &\geq C \int_D |k_a(w)|^{q-2} |k'_a(w)|^2 N_\varphi(w) dA(w) \\ &= C \frac{4}{p^2} |a|^2 (1 - |a|^2)^{\frac{q}{p}} \int_D \frac{N_\varphi(w)}{|1 - \bar{a}w|^{2 + \frac{2q}{p}}} dA(w) \\ &= C \frac{4}{p^2} |a|^2 (1 - |a|^2)^{\frac{q}{p} - 2} \int_D \frac{N_\varphi(w)}{|1 - \bar{a}w|^{\frac{2q}{p} - 2}} |\sigma'_a(w)| dA(w) \\ &= C \frac{4}{p^2} |a|^2 (1 - |a|^2)^{\frac{q}{p} - 2} \int_D \frac{N_\varphi(\sigma_a(z))}{|1 - \bar{a}\sigma_a(z)|^{\frac{2q}{p} - 2}} dA(z). \end{aligned}$$

Here $\sigma_a = \sigma_a^{-1}$ is the Möbius self-map of D as in Lemma 2.5, and the change of variable $z = \sigma_a(w)$ was made. Now,

$$\frac{1}{|1 - \bar{a}\sigma_a(z)|} = \frac{|1 - \bar{a}z|}{1 - |a|^2} \geq \frac{1}{2(1 - |a|^2)}, \quad \text{as } |z| \leq \frac{1}{2},$$

so

$$\|C_\varphi k_a\|_q^q \geq \frac{C|a|^2}{(1 - |a|^2)^{\frac{q}{p}}} \int_{\frac{1}{2}D} N_\varphi(\sigma_a(z)) dA(z).$$

Since $\sigma_a \circ \varphi(0) > \frac{1}{2}$, if $|a|$ is sufficiently close to 1, applying Lemmas 2.5 and 2.4, we have

$$\int_{\frac{1}{2}D} N_\varphi(\sigma_a(z)) dA(z) = \int_{\frac{1}{2}D} N_{\sigma_a \circ \varphi}(z) dA(z) \geq 4N_{\sigma_a \circ \varphi}(0) = 4N_\varphi(a).$$

Therefore,

$$\|C_\varphi k_a\|_q^q \geq \frac{C|a|^2 N_\varphi(a)}{(1 - |a|^2)^{\frac{q}{p}}}.$$

Since $\log(\frac{1}{|a|})$ is comparable to $(1 - |a|^2)$, if $|a|$ is sufficiently close to 1, by (2.1), we get

$$\|C_\varphi\|_{e, H^p \rightarrow H^q}^q \geq C_1 \limsup_{|a| \rightarrow 1^-} \frac{N_\varphi(a)}{[\log(\frac{1}{|a|})]^{\frac{q}{p}}}.$$

Now, we turn to prove

$$\|C_\varphi\|_{e, H^p \rightarrow H^q}^q \leq C_2 \limsup_{|a| \rightarrow 1^-} \frac{N_\varphi(a)}{[\log(\frac{1}{|a|})]^{\frac{q}{p}}}.$$

Since, for each n , K_n is compact, we have that $C_\varphi K_n$ is compact and for each n ,

$$\|C_\varphi\|_{e, H^p \rightarrow H^q} = \|C_\varphi R_n + C_\varphi K_n\|_{e, H^p \rightarrow H^q} \leq \|C_\varphi R_n\|. \quad (2.2)$$

Let U denote the closed unit ball in $H^p(D)$, for $f(z) \in U$, by Lemma 2.3,

$$\|C_\varphi R_n f\|_q^q = |R_n f(\varphi(0))|^q + \frac{q^2}{2} \int_D |R_n f(w)|^{q-2} |(R_n f)'(w)|^2 N_\varphi(w) dA(w). \quad (2.3)$$

For a fixed constant r_0 , $\frac{1}{2} < r_0 < 1$, we have

$$\begin{aligned} & \int_D |R_n f(w)|^{q-2} |(R_n f)'(w)|^2 N_\varphi(w) dA(w) \\ &= \int_{r_0 D} |R_n f(w)|^{q-2} |(R_n f)'(w)|^2 N_\varphi(w) dA(w) \\ & \quad + \int_{D \setminus r_0 D} |R_n f(w)|^{q-2} |(R_n f)'(w)|^2 N_\varphi(w) dA(w). \end{aligned} \quad (2.4)$$

Let $M = \sup_{|w| > r_0} \frac{N_\varphi(w)}{[\log(\frac{1}{|w|})]^{\frac{q}{p}}}$. By Lemma 2.6, we have

$$|R_n f(w)|^{q-p} \leq \frac{C \|R_n f\|_p^{q-p}}{(1 - |w|)^{\frac{q-p}{p}}}.$$

Then

$$\begin{aligned} & \int_{D \setminus r_0 D} |R_n f(w)|^{q-2} |(R_n f)'(w)|^2 N_\varphi(w) dA(w) \\ & \leq CM \|R_n f\|_p^{q-p} \int_{D \setminus r_0 D} |R_n f(w)|^{p-2} |(R_n f)'(w)|^2 \frac{[\log(\frac{1}{|w|})]^{\frac{q}{p}-1} [\log(\frac{1}{|w|})]}{(1 - |w|)^{\frac{q-p}{p}}} dA(w). \end{aligned}$$

Since $\log(\frac{1}{|w|}) \leq 2(1 - |w|)$ as $|w| \geq \frac{1}{2}$, we have

$$\begin{aligned} & \int_{D \setminus r_0 D} |R_n f(w)|^{q-2} |(R_n f)'(w)|^2 N_\varphi(w) dA(w) \\ & \leq CM \|R_n f\|_p^{q-p} \int_{D \setminus r_0 D} |R_n f(w)|^{p-2} |(R_n f)'(w)|^2 \log\left(\frac{1}{|w|}\right) dA(w). \end{aligned}$$

For $\varphi(z) = z$, we have $N_\varphi(w) = \log(\frac{1}{|w|})$. By Lemma 2.3, we get

$$\int_{D \setminus r_0 D} |R_n f(w)|^{p-2} |(R_n f)'(w)|^2 \log\left(\frac{1}{|w|}\right) dA(w) \leq C \|R_n f\|_p^p.$$

By Lemma 2.2 and $f(z) \in U$, we get

$$\int_{D \setminus r_0 D} |R_n f(w)|^{q-2} |(R_n f)'(w)|^2 N_\varphi(w) dA(w) \leq CM \|R_n f\|_p^q \leq CM.$$

Using the Cauchy integral formula, for $0 < r < 1$, $w \in rD$, we have

$$(R_n f)'(w) = \frac{1}{2\pi i} \int_{\partial(rD)} \frac{(R_n f)(\xi)}{(\xi - w)^2} d\xi.$$

By the Hölder inequality, letting $r \rightarrow 1^-$, we get

$$|(R_n f)'(w)| \leq \frac{C \|R_n f\|_p}{(1 - |w|)^2}, \quad \text{where } C \text{ is independent of } f.$$

By Lemma 2.6, we get

$$|R_n f(w)|^{q-2} |(R_n f)'(w)|^2 \leq \frac{C \|R_n f\|_p^q}{(1 - |w|)^{\frac{q+4p-2}{p}}}.$$

By Lemma 2.2, we get $\|R_n f\|_p \rightarrow 0$, as $n \rightarrow \infty$. So, as $n \rightarrow \infty$,

$$|R_n f(w)|^{q-2} |(R_n f)'(w)|^2 \rightarrow 0, \quad \text{uniformly on } r_0 D \quad \text{and} \quad |R_n f(\varphi(0))| \rightarrow 0. \quad (2.5)$$

By Lemma 2.3, for $f(z) = z$ and $p = 2$, we get

$$\|\varphi\|_2^2 = |\varphi(0)|^2 + 2 \int_D N_\varphi(w) dA(w).$$

So, by Lemma 2.6, we get

$$\int_{r_0 D} N_\varphi(w) dA(w) \leq C, \quad \text{where } C \text{ is independent of } \varphi. \quad (2.6)$$

Combining (2.2)–(2.6) and letting $n \rightarrow \infty$, we get

$$\|C_\varphi\|_{e, H^p \rightarrow H^q}^q \leq C \sup_{|w| > r_0} \frac{N_\varphi(w)}{[\log(\frac{1}{|w|})]^{\frac{q}{p}}}.$$

Let $r_0 \rightarrow 1^-$. Then

$$\|C_\varphi\|_{e, H^p \rightarrow H^q}^q \leq C_2 \limsup_{|a| \rightarrow 1^-} \frac{N_\varphi(a)}{[\log(\frac{1}{|a|})]^{\frac{q}{p}}}.$$

The proof is completed.

References

- [1] Luo, L., The essential norm of a composition operator on the Bergman space of the unit ball, *Complex Variables*, **49**, 2004, 845–850.
- [2] Luo, L., The essential norm of a composition operator on Hardy space of the unit ball (in Chinese), *Chin. Ann. Math.*, **28A**(6), 2007, 805–810.
- [3] Shapiro, J. H., The Essential Norm of a composition operator, *Ann. of Math.*, **125**, 1987, 375–404.
- [4] Smith, W., Composition operators between Bergman and Hardy spaces, *Trans. Amer. Math. Soc.*, **348**, 1996, 2331–2347.
- [5] Zhu, K., Duality of Bloch spaces and norm convergence of Taylor series, *Michigan Math. J.*, **38**, 1991, 89–101.
- [6] Zhu, K. H., *Operator Theory in Function Space*, Marcel-Dekker, New York, 1990.