

On the Normal Subgroup with Exactly Two G -Conjugacy Class Sizes***

Xianhe ZHAO* Xiuyun GUO**

Abstract Let G be a finite group with a non-central Sylow r -subgroup R , $Z(G)$ the center of G , and N a normal subgroup of G . The purpose of this paper is to determine the structure of N under the hypotheses that N contains R and the G -conjugacy class size of every element of N is either 1 or m . Particularly, it is shown that N is Abelian if $N \cap Z(G) = 1$ and the G -conjugacy class size of every element of N is either 1 or m .

Keywords Normal subgroups, Conjugacy class sizes, Nilpotent groups

2000 MR Subject Classification 20D10, 20D20

1 Introduction

All groups considered in this paper are finite. Let G be a group, π a set of some primes and x an element of G . x^G denotes the conjugacy class containing x , $|x^G|$ denotes the size of x^G , x_π and $x_{\pi'}$ denote π -component and π' -component of x , respectively. Moreover, we write G_π for a Hall π -subgroup of G , $G_{\pi'}$ for a Hall π' -subgroup of G , and n_π for the π -part of n whenever n is a positive integer.

In 1904, Burnside proved that if a group G has a conjugacy class with prime power size, then G is not simple (see [3, Corollary II, p. 322]). Since then, many authors have investigated the relationship between the structure of a group and its conjugacy class sizes (for example, [1, 2, 4, 5, 8–14]). Among these results, a classic result by Itô [8] asserts that a group G is nilpotent if $|x^G| = 1$ or m for every $x \in G$. Recently, Beltrán and Felipe [2] proved that every Hall p' -subgroup of a p -solvable group is nilpotent if $|x^G| = 1$ or m for every p' -element x of G . On the other hand, the structure of a normal subgroup N of a group G was given if N is the union of some G -conjugacy classes (see [9–12]). Now, we are interested in the following question: Let G be a finite group and let N be a normal subgroup of G . If $|x^G| = 1$ or m for every element $x \in N$, is N nilpotent?

Our main result is the following theorem.

Theorem 1.1 *Let G be a finite group with a non-central Sylow r -subgroup R and N a normal subgroup of G containing R . If $|x^G| = 1$ or m for every element x of N , then N is nilpotent.*

Manuscript received March 9, 2008. Published online June 8, 2009.

*College of Mathematics and Information Science, Henan Normal University, Xinxiang 453007, Henan, China; Department of Mathematics, Shanghai University, Shanghai 200444, China.

E-mail: zhaoxianhe989@yahoo.com.cn

**Department of Mathematics, Shanghai University, Shanghai 200444, China. E-mail: xyguo@shu.edu.cn

***Project supported by the National Natural Science Foundation of China (No. 10771132), SGRC (No. GZ 310), the Research Grant of Shanghai University and the Shanghai Leading Academic Discipline Project (No. J50101).

2 Preliminaries

We first list some lemmas which are useful in the proof of our main result.

Lemma 2.1 (see [5, Lemma 1.1]) *Let N be a normal subgroup of a group G and x an element of G . Then*

- (a) $|x^N|$ divides $|x^G|$;
- (b) $|(Nx)^{G/N}|$ divides $|x^G|$.

Corollary 2.1 *Let N be a normal subgroup of a group G and $N \cap Z(G) = 1$. If $|x^G| = 1$ or m for every element x of N , then N is Abelian.*

Proof By the assumption of this corollary, we may assume $|N| = km + 1$, where k is the number of noncentral G -conjugacy classes contained in N . It follows that $(|N|, m) = 1$. By Lemma 2.1, we deduce that N is Abelian.

Lemma 2.2 (see [7, Theorem 33.4]) *Let G be a group. A prime p does not divide any conjugacy class size of G if and only if G has a central Sylow p -subgroup.*

Lemma 2.3 *Let π be a set of some primes and N be a normal subgroup of a group G . If $\bar{x} = xN$ is a π -element, then there exists a π -element x^* of G such that $\bar{x} = \overline{x^*}$.*

Proof Let $o(\bar{x}) = n_0$ and $o(x) = n \cdot m$ such that n is a π -number and $(n, m) = 1$. Then $n_0 \mid n$ and $x^{n_0} \in N$. Since $(n, m) = 1$, there exist integers u and v such that $un + vm = 1$. It follows from $x = x^{un} \cdot x^{vm}$ that $xN = (x^m)^v N$. It is clear that $x^* = (x^m)^v$ is a π -element.

Lemma 2.4 (see [6, Theorem 1]) *Let G be a group acting transitively on a set Ω with $|\Omega| > 1$. Then there exist a prime p and a p -element $x \in G$ such that x acts without fixed point on Ω .*

3 Proof of Theorem 1.1

Now, we are equipped to prove the main result.

Assume that Theorem 1.1 is not true. Let G be a counterexample with minimal order, and $Z(G)$ be the center of the group G . Without loss of generality, we may replace N by $NZ(G)$. Therefore we may assume that $Z(G) \leq N$. We will complete the proof by the following steps.

Step 1 $N_p \not\leq Z(G)$ for any prime divisor $p(\neq r)$ of $|N|$.

If not, there exists a prime divisor $q(\neq r)$ of the order of N such that $N_q \leq Z(G)$ and thus $N_q \trianglelefteq G$. Consider the quotient groups G/N_q and N/N_q . For convenience, we use “ \sim ” to work in the factor group mod N_q . Obviously, \tilde{R} is a non-central Sylow r -subgroup of \tilde{G} , and \tilde{N} is a normal subgroup of \tilde{G} containing \tilde{R} . Let \tilde{x} be an element of \tilde{N} and y an element of G . We may assume that x is a q' -element of N by Lemma 2.3. If $\tilde{x}\tilde{y} = \tilde{y}\tilde{x}$, then $[x, y] \in N_q$. So y normalizes the group $\langle x \rangle \times N_q$, and hence $[x, y] \in \langle x \rangle$. Consequently, $[x, y] = 1$. So $C_{\tilde{G}}(\tilde{x}) = \widetilde{C_G(x)}$. Therefore, $|\tilde{x}^{\tilde{G}}| = |\tilde{G} : C_{\tilde{G}}(\tilde{x})| = |\tilde{G} : \widetilde{C_G(x)}| = |G : C_G(x)| = 1$ or m . This means that the hypotheses of the theorem are inherited by factor group \tilde{G} and \tilde{N} . We conclude that \tilde{N} is nilpotent by the minimal choice of G . Since $N_q \leq Z(G)$ while $Z(G) \leq Z(N)$, N is nilpotent, a contradiction.

In the following, we will consider the quotient groups $G/Z(G)$ and $N/Z(G)$. For convenience, we use “ $-$ ” to work in the factor group mod $Z(G)$.

Step 2 For any $1 \neq \bar{x} \in \bar{N}$, we have

- (i) If $o(\bar{x})$ is a power of a prime p , then $C_N(x)_{p'} \leq Z(C_G(x))$.
- (ii) If $o(\bar{x})$ is not a prime power, then $C_N(x) \leq Z(C_G(x))$.

If $o(\bar{x})$ is a power of p , then by Lemma 2.3 we may assume that x is a p -element. For any p' -element $y \in N \cap C_G(x) = C_N(x)$, since $C_G(xy) = C_G(x) \cap C_G(y) \subseteq C_G(x)$, the assumption of the theorem implies $C_G(xy) = C_G(x)$, and hence $C_G(x) \subseteq C_G(y)$. It follows that $y \in Z(C_G(x))$, and therefore $C_N(x)_{p'} \leq Z(C_G(x))$.

If $o(\bar{x})$ is not a prime power, then $o(x)$ is also not a prime power. So we may assume $x = x_1 x_2 \cdots x_s$, where the order of each x_i is a power of a prime p_i and x_i commutes pairwise with $p_i \neq p_j$ ($i, j = 1, 2, \dots, s$ and $s \geq 2$). Since $o(\bar{x})$ is not a prime power, there at least exist two elements beyond $Z(G)$ among x_i ($1 \leq i \leq s$), say, x_1 and x_2 . Noticing that x, x_1 and x_2 are non-central elements and $C_G(x) = C_G(x_1) \cap C_G(x_2 \cdots x_s) = C_G(x_2) \cap C_G(x_1 x_3 \cdots x_s)$, we have $C_G(x) = C_G(x_1) = C_G(x_2)$ by the assumption of the theorem. On the other hand, (i) implies that $C_N(x)_{p'_1} \leq Z(C_G(x))$ and $C_N(x)_{p'_2} \leq Z(C_G(x))$, while $C_N(x)_{p'_1} C_N(x)_{p'_2} = C_N(x)$. So $C_N(x) \leq Z(C_G(x))$.

Step 3 Let $1 \neq \bar{x}, \bar{y} \in \bar{N}$ and $o(\bar{x})$ be not a prime power. If $C_G(x) \neq C_G(y)$, then $C_N(x) \cap C_N(y) = Z(G)$.

By (ii), we have $C_N(x) \leq Z(C_G(x))$. If there exists an element a such that $1 \neq a \in C_N(x) \cap C_N(y)$ but $a \notin Z(G)$, then $C_G(x) = C_G(a)$, and therefore $C_N(x) = C_N(a)$. Also, $a \in C_N(y)$ implies $y \in C_N(a) = C_N(x) \leq Z(C_G(x))$. It follows that $C_G(x) \subseteq C_G(y)$. Hence $C_G(x) = C_G(y)$, a contradiction.

Step 4 For any $1 \neq \bar{x} \in \bar{N}$, we have

- (iii) If $o(\bar{x})$ is a power of a prime q , then $|\bar{x}^{\bar{G}}|_{q'} = m_{q'}$.
- (iv) If $o(\bar{x})$ is not a prime power, then $|\bar{x}^{\bar{G}}| = m$.

Let $C_{\bar{G}}(\bar{x}) = \bar{H}$. For every element $\bar{y} \in \bar{H}$ of order of a power of a prime t where $t \in q'$, we may assume that y is a t -element and $y \in H$ by Lemma 2.3. Therefore we have that $\langle \bar{x}, \bar{y} \rangle$ is a cyclic group and hence $\langle x, y \rangle$ is Abelian. So $\bar{y} \in \overline{C_G(x)}$. It follows that $|\bar{H}|_t = |\overline{C_G(x)}|_t$. So $|\bar{H}|_{q'} = |C_G(x)|_{q'}$. Therefore

$$|\bar{x}^{\bar{G}}|_{q'} = \frac{|\bar{G}|_{q'}}{|\bar{H}|_{q'}} = \frac{|\bar{G}|_{q'}}{|C_G(x)|_{q'}} = \frac{|G|_{q'}/|Z(G)|_{q'}}{|C_G(x)|_{q'}/|Z(G)|_{q'}} = \frac{|G|_{q'}}{|C_G(x)|_{q'}} = |G : C_G(x)|_{q'} = m_{q'}.$$

Next, if \bar{x} is not a prime power order element, then $o(x)$ is also not a prime power. So we may assume that $x = x_1 x_2 \cdots x_s$, where the order of each x_i is a power of a prime p_i and x_i commute pairwise with $p_i \neq p_j$ ($i, j = 1, 2, \dots, s$ and $s \geq 2$). Since $o(\bar{x})$ is not a prime power, there at least exist two elements beyond $Z(G)$ among x_i ($1 \leq i \leq s$), say, x_1 and x_2 . So $\bar{x} = \bar{x}_1 \bar{x}_2 \bar{x}_3 \cdots \bar{x}_s$. Obviously, $C_{\bar{G}}(\bar{x}) = C_{\bar{G}}(\bar{x}_1) \cap C_{\bar{G}}(\bar{x}_2 \bar{x}_3 \cdots \bar{x}_s) = C_{\bar{G}}(\bar{x}_2) \cap C_{\bar{G}}(\bar{x}_1 \bar{x}_3 \cdots \bar{x}_s)$, and it is clear that $C_{\bar{G}}(\bar{x}) \leq C_{\bar{G}}(\bar{x}_1)$ and $C_{\bar{G}}(\bar{x}) \leq C_{\bar{G}}(\bar{x}_2)$. Hence $|\bar{x}_1^{\bar{G}}| \mid |\bar{x}^{\bar{G}}|$ and $|\bar{x}_2^{\bar{G}}| \mid |\bar{x}^{\bar{G}}|$. By (iii), we have $m_{p'_1} \mid |\bar{x}^{\bar{G}}|$ and $m_{p'_2} \mid |\bar{x}^{\bar{G}}|$. So $|\bar{x}^{\bar{G}}| = m$.

Step 5 If $1 \neq \bar{x} \in \bar{N}$ is not a prime power order element, then $C_{\bar{G}}(\bar{x}) = \overline{C_G(x)}$, particularly, $C_{\bar{N}}(\bar{x}) = \overline{C_N(x)}$.

Since

$$|\bar{x}^{\bar{G}}| = |\bar{G} : C_{\bar{G}}(\bar{x})| \leq |\bar{G} : \overline{C_G(x)}| = |G : C_G(x)| = m$$

while $|\bar{x}^{\bar{G}}| = m$ by (iv), thus we obtain $C_{\bar{G}}(\bar{x}) = \overline{C_G(x)}$. Particularly, $C_{\bar{N}}(\bar{x}) = C_{\bar{G}}(\bar{x}) \cap \bar{N} = \overline{C_G(x)} \cap \bar{N} = \overline{C_G(x) \cap N} = \overline{C_N(x)}$.

Step 6 Let $g_0 \in N - Z(G)$ and \bar{g}_0 be not a prime power order element. Consider the conjugacy class $\bar{g}_0^{\bar{N}}$ of \bar{g}_0 in \bar{N} . Then there exists some non-central element x of N such that $\bar{g}_0^{\bar{N}} \cap \overline{C_N(x)} = \emptyset$.

Consider the conjugacy class g_0^N of g_0 in N . Noting that N operates transitively on the set g_0^N with $|g_0^N| > 1$, we have that there exists an element x of N such that x operates without fixed point on g_0^N by Lemma 2.4. It follows that $g_0^N \cap C_N(x) = \emptyset$. So $\bar{g}_0^{\bar{N}} \cap \overline{C_N(x)} = \emptyset$.

Step 7 There exists a $\{p, r\}$ -element g of N such that \bar{g} is a $\{p, r\}$ -element of \bar{N} for any prime divisor $p(\neq r)$ of $|N|$.

According to Step 1, for any prime divisor $p(\neq r)$ of $|N|$, there exists a non-central p -element in N , say, x . By (i), we have $C_N(x) = C_N(x)_p \times C_N(x)_{p'}$. We claim $C_N(x)_r \not\leq Z(G)$. Otherwise, $C_N(x)_r \leq Z(G)$. As R is non-central, there exists a non-central r -element z such that $z \in N \setminus C_N(x)_r$. So $C_N(x)_r < \langle C_N(x)_r, z \rangle \leq C_G(z)$, in contradiction to $|z^G| = |x^G| = m$. Take $z \in C_N(x)_r \setminus Z(G)$ and let $g = xz$. Thus $\bar{g} = \bar{x}\bar{z}$. It is clear that g and \bar{g} satisfy the requirement of Step 7.

Step 8 If there exist an r -element x of N and a prime divisor $p(\neq r)$ of $|N|$ such that $p \nmid |\overline{C_N(x)}|$, then $|\bar{N}|_p \mid m$.

As

$$|x^N| = |N : C_N(x)| = |\bar{N} : \overline{C_N(x)}|,$$

it leads to $|\bar{N}|_p \mid |x^N|$ since p does not divide $|\overline{C_N(x)}|$. It follows that $|\bar{N}|_p \mid |x^G|$ by Lemma 2.1. The hypotheses of the theorem imply $|\bar{N}|_p \mid m$.

Step 9 $|\overline{C_N(y)}|$ and $|\bar{N}|$ have the same prime divisors for any r -element y of N .

If it is not true, there exist an r -element x_0 of N and a prime divisor p of N such that p does not divide $|\overline{C_N(x_0)}|$. Obviously, $p \neq r$. By Step 8, we have $|\bar{N}|_p \mid m$.

By Step 7, we may take a $\{p, r\}$ -element $g \in N$ such that \bar{g} is a $\{p, r\}$ -element. Applying Step 6, we have that there exists a non-central element x of N such that $\bar{g}^{\bar{N}} \cap \overline{C_N(x)} = \emptyset$. Consider that $\overline{C_N(x)}$ operates on $\bar{g}^{\bar{N}}$ by conjugation. Notice that no element in $\overline{C_N(x)}$ distinct from 1 centralizes any element in $\bar{g}^{\bar{N}}$ by Steps 3 and 5. So all orbits of $\overline{C_N(x)}$ on $\bar{g}^{\bar{N}}$ have the same length $|\overline{C_N(x)}|$, which implies $|\overline{C_N(x)}| \mid |\bar{g}^{\bar{N}}|$. Also $|\bar{g}^{\bar{N}}| \mid |\bar{g}^{\bar{G}}|$ by Lemma 2.1, so

$$|\overline{C_N(x)}| \mid |\bar{g}^{\bar{G}}|. \quad (3.1)$$

On the other hand, it is obvious that $\overline{C_N(g)}$ operates without fixed points on $\bar{g}^{\bar{G}} - \bar{g}^{\bar{G}} \cap \overline{C_N(g)}$. By Steps 3 and 5 once again, we have

$$|\overline{C_N(g)}| \mid (|\bar{g}^{\bar{G}}| - |\bar{g}^{\bar{G}} \cap \overline{C_N(g)}|). \quad (3.2)$$

Since N contains a Sylow r -subgroup R of G , $|\overline{C_N(g)}|_r = |\overline{C_N(x)}|_r$, from which the relationship between (3.1) and (3.2) becomes

$$|\overline{C_N(g)}|_r \mid |\bar{g}^{\bar{G}} \cap \overline{C_N(g)}|. \quad (3.3)$$

Notice $|\bar{N}|_p \mid m$. Also, Step 4 implies that $|\bar{N}|_p \mid |\bar{g}^{\bar{G}}|$. Obviously, $|\overline{C_N(g)}|_p \leq |\bar{N}|_p$. So $|\overline{C_N(g)}|_p \mid |\bar{g}^{\bar{G}}|$. Noticing (3.2), we have

$$|\overline{C_N(g)}|_p \mid |\bar{g}^{\bar{G}} \cap \overline{C_N(g)}|. \quad (3.4)$$

By (3.3) and (3.4), we have

$$|\overline{C_N(g)}|_{\{p,r\}} \mid |\overline{g}^G \cap \overline{C_N(g)}|. \quad (3.5)$$

Noticing that $\overline{C_N(g)}$ is Abelian by (ii), we get

$$|\overline{g}^G \cap \overline{C_N(g)}| = |\overline{C_N(g)}|_{\{p,r\}}, \quad (3.6)$$

a contradiction.

Step 10 The final contradiction.

Let p be a prime divisor of $|\overline{N}|$ with $p \neq r$. Choose a non-central r -element x_0 such that $|\overline{C_N(x_0)}|_p$ is as small as possible. By Step 9 we have $|\overline{C_N(x_0)}|_p > 1$. So, it is clear that we may take a $\{p, r\}$ -element $g \in C_N(x_0)$ such that \overline{g} is also a $\{p, r\}$ -element. Arguing as in Step 9, we may see that (3.1) and (3.2) still hold, and therefore (3.3) also holds.

Noting $C_N(g) = C_N(x_0)$ by Step 3, we have $|\overline{C_N(g)}|_p = |\overline{C_N(x_0)}|_p$. Noticing $|\overline{C_N(x_0)}|_p \leq |\overline{C_N(x)}|_p$ by the choice of x_0 , we have $|\overline{C_N(g)}|_p \leq |\overline{C_N(x)}|_p$. Consequently, $|\overline{C_N(g)}|_p \mid |\overline{g}^G|$ by (3.1). By using a similar argument as in Step 9, (3.4) is obtained. Arguing as in Step 9 once again, we have (3.5), and thus equation (3.6) holds, a contradiction.

Corollary 3.1 *Let G be a finite group with a non-central Sylow r -subgroup R and N a normal subgroup of G containing R . If $|x^G| = 1$ or m for every element x of N , then either N is Abelian or $N_{r'} \leq Z(G)$.*

Proof By Theorem 1.1, we have that N is nilpotent. So, for any $p(\neq r)$ -element x of N , we have $r \nmid |G : C_G(x)|$. If $N_{r'} \not\leq Z(G)$, then $r \nmid m$. So $R \leq Z(N)$. However, since R is non-central, there exists a non-central r -element x_0 such that $N \leq C_N(x_0)$. Thus we have $N_{r'} \leq Z(C_G(x_0))$ by (i) in the proof of Theorem 1.1. So N is Abelian.

Corollary 3.2 (see [8, Theorem 1]) *Let G be a group. If $|x^G| = 1$ or m for every element x in G , then G is nilpotent.*

Proof Let $N = G$. Obviously, N satisfies the hypotheses of Theorem 1.1. So $N = G$ is nilpotent by Theorem 1.1.

Acknowledgement The authors would like to thank the referees and the editors for their carefully reading the manuscript and for their helpful comments.

References

- [1] Baer, R., Group elements of prime power index, *Trans. Amer. Math. Soc.*, **75**(1), 1953, 20–47.
- [2] Beltrán, A. and Felipe, M. J., Finite groups with two p -regular conjugacy class lengths, *Bull. Austral. Math. Soc.*, **67**, 2003, 163–169.
- [3] Burnside, W., *Theory of Groups of Finite Order*, Cambridge University Press, Cambridge, 1911.
- [4] Camina, A. R. and Camina, R. D., Implications of conjugacy class size, *J. Group Theory*, **1**(3), 1998, 257–269.
- [5] Chillag, D. and Herzog, M., On the length of the conjugacy classes of finite groups, *J. Algebra*, **131**(1), 1990, 110–125.
- [6] Fein, B., Kantor, W. M. and Schacher, M., Relative Brauer groups II, *J. Reine Angew. Math.*, **328**, 1981, 39–57.
- [7] Huppert, B., *Character Theory of Finite Groups*, de Gruyter Expositions in Mathematics, **25**, Walter de Gruyter, Berlin, 1998.

- [8] Itô, N., On finite groups with given conjugate type I, *Nagoya Math. J.*, **6**, 1953, 17–28.
- [9] Riese, U. and Shahabi, M. A., Subgroups which are the union of four conjugacy classes, *Comm. Algebra*, **29**(2), 2001, 695–701.
- [10] Shahryari, M. and Shahabi, M. A., Subgroups which are the union of three conjugate classes, *J. Algebra*, **207**(1), 1998, 326–332.
- [11] Shi, W. J., A class of special minimal normal subgroups (in Chinese), *J. Southwest Teachers College*, **9**, 1984, 9–13.
- [12] Wang, J., A special class of normal subgroups (in Chinese), *J. Chengdu Univ. Sci. Tech.*, **4**, 1987, 115–119.
- [13] You, X. Z. and Qian, G. H., A new graph related to conjugacy classes of finite groups (in Chinese), *Chin. Ann. Math.*, **28A**(5), 2007, 631–636.
- [14] Zhang, J. P., Finite groups with many conjugate elements, *J. Algebra*, **170**(2), 1994, 608–624.