

Entanglement Rényi entropy and boson-fermion duality in massless Thirring model

Harunobu Fujimura

September 26, 2023 @ HET Camp

arXiv:2309.11889

collaborators: T.Nishioka and S.Shimamori

1. Introduction
2. Analysis of the ERE and MRI
3. Parameter dependence of the ERE and MRI
4. Summary

Introduction




1. Entanglement in QFT

State of total system : $|0\rangle$

Reduced density matrix : $\rho_{\text{tot}} = |0\rangle\langle 0|$

Total space : $V_{\text{tot}} = V \cup \bar{V}$

Reduced density matrix : $\rho_V \equiv \text{Tr}_{\bar{V}}[\rho_{\text{tot}}]$



- Entanglement Rényi entropy (ERE) :

$$S_n(V) \equiv \frac{1}{1-n} \log \text{Tr}_V [\rho_V^n] , \quad n \in \mathbb{Z}_+ ,$$

- Mutual Rényi information (MRI) :

$$I_n(V_1, V_2) \equiv S_n(V_1) + S_n(V_2) - S_n(V_1 \cup V_2) ,$$

\Rightarrow Correlation between V_1 and V_2 , where $V = V_1 \cup V_2$.

➡ ERE and MRI are important quantities that characterize QFT.

1. Entanglement in QFT

However, Calculating $\text{Tr}_V [\rho_V^n]$ in QFT is very difficult...

$$\text{Tr}_V [\rho_V^n] \sim Z_n$$

Partition function

Known exact results (1+1 dimension) :



- ✓ Free massless fermion. n sheets, N intervals. [Casini, Fosco, Huerta 2005]
- ✓ Arbitrary CFT. But single interval only. [Holzhey, Larsen, Wilczek 1994]
- ✓ Free compact boson. 2 sheets, 2 intervals. [Calabrese, Cardy, Tonni 2011]

➡ There is no exact result for interacting system.

1. Boson-fermion duality

It's difficult to compute the ERE and MRI in interacting system.

⇒ We focused on **boson-fermion duality**. [Karch, Tong, Turner 2019]


fermionic theory \mathcal{T}_F

discrete sym : \mathbb{Z}_2^F

local op : $\hat{\mathcal{O}}^F$

partition function : Z^F

fermionization



bosonization

bosonic theory \mathcal{T}_B

discrete sym : \mathbb{Z}_2^B

local op : $\hat{\mathcal{O}}^B$

partition function : Z^B

fermionization dictionary (1+1 dim, closed manifold) :

$$\mathcal{T}_F = \frac{\mathcal{T}_B \times \text{Kitaev}}{\mathbb{Z}_2^B}$$


couple with the TQFT

gauging \mathbb{Z}_2^B sym

1. Our work

What we did

- Massless Thirring model (1+1 dim, fermion, 4-point interaction)
- Combined conventional method (Replica method) and **boson-fermion duality**
- Got exact results on ERE and MRI in the case of the figure below
- Saw parameter dependence of ERE and MRI



Analysis of the ERE and MRI

The model

Massless Thirring model

$$\mathcal{L}_F = i\bar{\psi}\not{\partial}\psi + \underbrace{\frac{\pi}{2}\lambda(\bar{\psi}\gamma^\mu\psi)(\bar{\psi}\gamma_\mu\psi)}_{\text{interaction}}$$

ψ : Dirac fermion

\mathbb{Z}_2^F : $\psi \rightarrow -\psi$

λ : Thirring coupling

difficult to analyze

Free compact boson

$$\mathcal{L}_B = -\frac{R^2}{8\pi}\partial_\mu\phi\partial^\mu\phi$$

$$\phi \sim \phi + 2\pi$$

ϕ : scalar

$$\mathbb{Z}_2^B : \phi \rightarrow \phi + \pi$$

R : compact boson radius

easy to analyze

fermionize

$$1 + \lambda = \frac{4}{R^2}$$





We fermionize compact boson and get massless Thirring model.

Replica method

Target : $S_2(V) = -\log \text{Tr}_V [\rho_V^2]$

$$\rho_V(\psi_i, \psi_j) = \text{Tr}_{\bar{V}} [\langle \psi_{j,V} | 0 \rangle \langle 0 | \psi_{i,V} \rangle]$$


$$\begin{aligned} \text{Tr}_V [\rho_V^2] &= \sum_{\psi_1, \psi_2} \rho_V(\psi_1, \psi_2) \rho_V(\psi_2, \psi_1) \\ &\sim Z_{\Sigma_{2,2}}^F \end{aligned}$$



$\Sigma_{2,2}$

➡ We have to calculate the partition function $Z_{\Sigma_{2,2}}^F$

Conformal map [Lumin, Mathur 2001]



cross-ratio : $x = \frac{(v_1 - u_1)(v_2 - u_2)}{(u_2 - u_1)(v_2 - v_1)}$



moduli of torus : τ

$$Z_{\Sigma_{2,2}}^F(x) = (\text{UV div}) f(x) Z_T^F(\tau)$$



We can map to torus

Analytical result

$$S_2(V) = -\log f(x)(UV \text{ div}) - \log Z_T^F \xrightarrow{\text{fermionization}}$$

$$S_2(V) = -\log(UV \text{ div}) - \log (\text{sum of } Z_T^B[t]s) \longleftarrow \text{free theory (known), } t : \mathbb{Z}_2^B \text{ gauge field}$$

Exact result

$$S_2(V, \lambda) = S_2(V, 0) - \frac{1}{2} \log \left[\frac{1}{2\vartheta_3^4(\tau)} \sum_{j=2}^4 \vartheta_j^2(\tau(1+\lambda)) \vartheta_j^2\left(\frac{\tau}{1+\lambda}\right) \right]$$

where,

$S_2(V, 0)$: Free ERE \rightarrow consistent with existing result



$\vartheta_j(\tau)$, $j = 2, 3, 4$: Jacobi theta functions

We got the exact ERE and MRI in massless Thirring model

Parameter dependence of the ERE and MRI

Cross-ratio dependence of ERE

$$\text{Deviation : } \Delta S_2(x) = S_2(V, \lambda) - S_2(V, 0)$$



Single interval, CFT [Holzhey et al 1994]


$$S_n = \frac{c}{6} \left(1 + \frac{1}{n} \right) \log \left(\frac{v-u}{\epsilon} \right)$$

c : central charge,
 ϵ : UV cutoff



We got reasonable results

Coupling dependence of ERE



T-duality :


$$\lambda \rightarrow \lambda_{\text{dual}} \equiv -\frac{\lambda}{\lambda + 1}$$

$\lambda > 0$ and $\lambda < 0$ correspond



We investigated the behavior with respect to the coupling constants.

Cross-ratio dependence of MRI



As coupling λ becomes larger,
the entanglement between V_1 and V_2
grows stronger.

Summary

Summary


- ERE and MRI are important quantities that characterize QFT.
- We **exactly** derived the ERE and MRI in an interacting system by **fermionization**.
- We saw the behaviors of ERE and MRI from exact results.

Future work

- ERE and MRI for 3-intervals or more.
- Massive Thirring model.
- Other quantum information measures.

Thank you.

Backup slides : The duality web



Backup slides : Coupling dependence of ERE

