

## SECTION 5.3: MARGINAL AND CONDITIONAL PROBABILITY DISTRIBUTIONS

*Exercise (5.23).* In Example 5.4 and Exercise 5.5, we considered the joint density of  $Y_1$ , the proportion of the capacity of the tank that is stocked at the beginning of the week, and  $Y_2$ , the proportion of the capacity sold during the week, given by

$$f(y_1, y_2) = \begin{cases} 3y_1, & 0 \leq y_2 \leq y_1 \leq 1, \\ 0, & \text{elsewhere.} \end{cases}$$

- (a) Find the marginal density function for  $Y_2$ .

*Solution:*

$$\begin{aligned} f_2(y_2) &= \int_{y_2}^1 3y_1 dy_1 = \left. \frac{3y_1^2}{2} \right|_{y_2}^1 = \frac{3}{2} - \frac{3y_2^2}{2} \\ &= \frac{3}{2}(1 - y_2^2) \end{aligned}$$

□

- (b) For what values of  $y_2$  is the conditional density  $f(y_1|y_2)$  defined?

*Solution:*

$$\begin{aligned} f(y_1|y_2) &= \frac{f(y_1, y_2)}{f_2(y_2)} = \frac{3y_1}{\frac{3}{2}(1 - y_2^2)} = \frac{2y_1}{1 - y_2^2} \\ &= \begin{cases} \frac{2y_1}{1 - y_2^2}, & y_2 \leq y_1 \leq 1 \\ 0, & \text{elsewhere} \end{cases} \end{aligned}$$

So its defined for  $0 \leq y_2 < 1$  to avoid division by 0.

□

- (c) What is the probability that more than half a tank is sold given that three-fourths of a tank is stocked?

*Solution:* We have  $P(Y_2 > 1/2 | Y_1 = 3/4)$

$$\begin{aligned} f_1(y_1) &= \int_0^{y_1} 3y_1 dy_2 = 3y_1 y_2 \Big|_0^{y_1} = 3y_1^2 \\ f(y_2 | y_1) &= \frac{3y_1}{3y_1^2} = \frac{1}{y_1} \\ P(Y_2 > 1/2 | Y_1 = 3/4) &= \int_{1/2}^{3/4} \frac{1}{3/4} dy_2 = \int_{1/2}^{3/4} \frac{4}{3} dy_2 = \frac{4}{3} y_2 \Big|_{1/2}^{3/4} \\ &= \left(\frac{4}{3} * \frac{3}{4}\right) - \left(\frac{4}{3} * \frac{1}{2}\right) = 1 - \frac{4}{6} = \frac{1}{3} \end{aligned}$$

□

*Exercise (5.24).* In Exercise 5.6, we assumed that if a radioactive particle is randomly located in a square with sides of unit length, a reasonable model for the joint density function for  $Y_1$  and  $Y_2$  is

$$f(y_1, y_2) = \begin{cases} 1, & 0 \leq y_1 \leq 1, 0 \leq y_2 \leq 1, \\ 0, & \text{elsewhere.} \end{cases}$$

- (a) Find the marginal density functions for  $Y_1$  and  $Y_2$ .

*Solution:*

$$f_1(y_1) = 1 \text{ for } 0 \leq y_1 \leq 1$$

$$f_2(y_2) = 1 \text{ for } 0 \leq y_2 \leq 1$$

□

- (b) What is  $P(.3 < Y_1 < .5)$ ?  $P(.3 < Y_2 < .5)$ ?

*Solution:* Since its a rectangular region for each

$$P(.3 < Y_1 < .5) = P(.3 < Y_2 < .5) = .2 * 1 = .2$$

□

- (c) For what values of  $y_2$  is the conditional density  $f(y_1 | y_2)$  defined?

*Solution:*

$$f(y_1|y_2) = \frac{f(y_1, y_2)}{f_2(y_2)} = \frac{1}{1} = 1$$

Defined for  $0 \leq y_2 \leq 1$ . □

- (d) For any  $y_2$ ,  $0 \leq y_2 \leq 1$  what is the conditional density function of  $Y_1$  given that  $Y_2 = y_2$ ?

*Solution:* The conditional probability we calculated in the last one so

$$f(y_1|y_2) = \begin{cases} 1, & 0 \leq y_1 \leq 1 \\ 0, & \text{elsewhere} \end{cases}$$

For any  $y_2$  where  $0 \leq y_2 \leq 1$ . □

- (e) Find  $P(.3 < Y_1 < .5|Y_2 = .3)$ .

*Solution:*  $P(.3 < Y_1 < .5|Y_2 = .3) = .2/1 = .2$  □

- (f) Find  $P(.3 < Y_1 < .5|Y_2 = .5)$ .

*Solution:*  $P(.3 < Y_1 < .5|Y_2 = .5) = .2/1 = .2$  □

- (g) Compare the answers that you obtained in parts (a), (d), and (e). For any  $y_2$ ,  $0 \leq y_2 \leq 1$  how does  $P(.3 < Y_1 < .5)$  compare to  $P(.3 < Y_1 < .5|Y_2 = y_2)$ ?

*Solution:* There are all the same answers. □

*Exercise (5.25).* Let  $Y_1$  and  $Y_2$  have joint density function first encountered in Exercise 5.7:

$$f(y_1, y_2) = \begin{cases} e^{-(y_1+y_2)}, & y_1 > 0, y_2 > 0, \\ 0, & \text{elsewhere.} \end{cases}$$

- (a) Find the marginal density functions for  $Y_1$  and  $Y_2$ . Identify these densities as one of those studied in Chapter 4.

*Solution:* Exponential Distribution with

$$\begin{aligned} f_1(y_1) &= \int_0^\infty e^{-(y_1+y_2)} dy_2 = \int_0^\infty e^{-y_1} * e^{-y_2} dy_2 = e^{-y_1} \int_0^\infty e^{-y_2} dy_2 \\ &= e^{-y_1} \left[ -e^{-y_2} \right]_0^\infty = e^{-y_1} \lim_{y_2 \rightarrow \infty} -e^{-y_2} + 1 = e^{-y_1} * (0 + 1) = e^{-y_1} \\ f_2(y_2) &= \int_0^\infty e^{-(y_1+y_2)} dy_1 = e^{-y_2} \int_0^\infty e^{-y_1} dy_1 = e^{-y_2} \end{aligned}$$

□

(b) What is  $P(1 < Y_1 < 2.5)$ ?  $P(1 < Y_2 < 2.5)$ ?

*Solution:*

$$\begin{aligned} P(1 < Y_1 < 2.5) &= \int_1^{2.5} e^{-y_1} dy_1 = \left[ -e^{-y_1} \right]_1^{2.5} = -e^{-2.5} + e^{-1} = 0.28579 \\ P(1 < Y_2 < 2.5) &= \int_1^{2.5} e^{-y_2} dy_2 = 0.28579 \end{aligned}$$

□

(c) For what values of  $y_2$  is the conditional density  $f(y_1|y_2)$  defined?

*Solution:*

$$f(y_1|y_2) = \frac{e^{-(y_1+y_2)}}{e^{-y_2}} = e^{-y_1}$$

Define for  $y_2 > 0$ .

□

(d) For any  $y_2 > 0$ , what is the conditional density function of  $Y_1$  given that  $Y_2 = y_2$ ?

*Solution:*  $f(y_1|y_2) = e^{-y_1}$

□

(e) For any  $y_1 > 0$ , what is the conditional density function of  $Y_2$  given that  $Y_1 = y_1$ ?

*Solution:*  $f(y_2|y_1) = e^{-y_2}$

□

(f) For any  $y_2 > 0$ , how does the conditional density function  $f(y_1|y_2)$  that you obtained in part (d) compare to the marginal density function  $f_1(y_1)$  found in part (a)?

*Solution:* They are the same

□

(g) What does your answer to part (f) imply about marginal and conditional probabilities that  $Y_1$  falls in any interval?

*Solution:* They are the same □

*Exercise (5.27).* In Exercise 5.9, we determined that

$$f(y_1, y_2) = \begin{cases} 6(1 - y_2), & 0 \leq y_1 \leq y_2 \leq 1, \\ 0, & \text{elsewhere} \end{cases}$$

is a valid joint probability density function. Find

(a) the marginal density functions for  $Y_1$  and  $Y_2$ .

*Solution:*

$$\begin{aligned} f_1(y_1) &= \int_{y_1}^1 6(1 - y_2) dy_2 = 6 \left( \int_{y_1}^1 1 dy_2 - \int_{y_1}^1 y_2 dy_2 \right) = 6 \left( y_2 \Big|_{y_1}^1 - \frac{y_2^2}{2} \Big|_{y_1}^1 \right) \\ &= 6 \left( (1 - y_1) - \left( \frac{1}{2} - \frac{y_1^2}{2} \right) \right) = 6 \left( -y_1 + \frac{1}{2} + \frac{y_1^2}{2} \right) = 6 \frac{1 + y_1^2 - 2y_1}{2} \\ &= 3(1 + y_1^2 - 2y_1) = 3(1 - y_1)^2 \\ f_2(y_2) &= \int_0^{y_2} 6(1 - y_2) dy_1 = 6 \int_0^{y_2} (1 - y_2) dy_1 = 6 \left( y_1 - y_2 y_1 \right) \Big|_0^{y_2} \\ &= 6((y_2 - 0) - (y_2^2 - 0)) = 6y_2(1 - y_2) \end{aligned}$$

□

(b)  $P(Y_2 \leq 1/2 | Y_1 \leq 3/4)$ .

*Solution:*

$$\begin{aligned} P &= \int_0^{1/2} \int_{y_1}^{1/2} 6(1 - y_2) dy_2 dy_1 = 6 \int_0^{1/2} \left[ y_2 - \frac{y_2^2}{2} \right]_{y_1}^{1/2} dy_1 \\ &= 6 \int_0^{1/2} \left( (1/2 - y_1) - (1/8 - \frac{y_1^2}{2}) \right) dy_1 = 6 \int_0^{1/2} \left( \frac{3}{8} - y_1 + \frac{y_1^2}{2} \right) dy_1 \\ &= 6 \left[ \frac{3}{8} y_1 - \frac{y_1^2}{2} + \frac{y_1^3}{6} \right]_0^{1/2} = 6 \left[ \frac{3}{16} - \frac{1}{8} + \frac{1}{48} \right] = 6 \left[ \frac{9 - 6 + 1}{48} \right] \\ &= \frac{24}{48} = \frac{1}{2} = 0.5 \\ P(Y_1 < 3/4) &= \int_0^{3/4} 3(1 - y_1)^2 dy_1 = 3 \int_0^{3/4} (1 - 2y_1 + y_1^2) dy_1 = 3 \left( y_1 - y_1^2 + \frac{y_1^3}{3} \right) \Big|_0^{3/4} \\ &= 3 \left( \frac{3}{4} - \frac{9}{16} + \frac{27}{192} \right) = 0.984375 \end{aligned}$$

$$P(Y_2 \leq 1/2 | Y_1 \leq 3/4) = \frac{0.5}{0.984375} = 0.506937$$

□

(c) the conditional density function of  $Y_1$  given  $Y_2 = y_2$ .

$$\text{Solution: } f(y_1|y_2) = \frac{6(1-y_2)}{6y_2(1-y_2^2)} = \frac{1-y_2}{y_2(1-y_2)} = \frac{1}{y_2}$$

□

(d) the conditional density function of  $Y_2$  given  $Y_1 = y_1$ .

$$\text{Solution: } f(y_2|y_1) = \frac{6(1-y_2)}{3(1-y_1)^2} = \frac{2(1-y_2)}{(1-y_1)^2}$$

□

(e)  $P(Y_2 \geq 3/4|Y_1 = 1/2)$ .

*Solution:*

$$f(y_2|y_1 = 1/2) = \frac{2(1-y_2)}{(1-(1/2))^2} = \frac{2(1-y_2)}{1/4} = 8(1-y_2)$$

$$\begin{aligned} P(Y_2 \geq 3/4|y_1 = 1/2) &= \int_{3/4}^1 8(1-y_2)dy_2 = 8 \int_{3/4}^1 (1-y_2)dy_2 \\ &= 8(y_2 - \frac{y_2^2}{2}) \Big|_{3/4}^1 = 8((1 - (3/4)) - ((1/2) - (9/32))) \\ &= 0.25 \end{aligned}$$

□

*Exercise (5.29).* Refer to Exercise 5.11. Find

(a) the marginal density functions for  $Y_1$  and  $Y_2$ .

*Solution:* Triangle with vertices  $(-1, 0)$ ,  $(1, 0)$ ,  $(0, 1)$  has area 1 so  $f(y_1, y_2) = 1$ .

$$f_1(y_1) = \int_0^{1-|y_1|} 1dy_2 = 1 - |y_1|, \quad -1 \leq y_1 \leq 1$$

$$f_2(y_2) = \int_{-(1-y_2)}^{1-y_2} 1dy_1 = 2(1-y_2), \quad 0 \leq y_2 \leq 1$$

□

(b)  $P(Y_2 > 1/2|Y_1 = 1/4)$ .

*Solution:*

$$f(y_2|y_1) = \frac{1}{1 - |y_1|} = \frac{1}{3/4} = \frac{4}{3}, \quad 0 \leq y_2 \leq 3/4$$

$$P(Y_2 > 1/2 | Y_1 = 1/4) = \int_{1/2}^{3/4} \frac{4}{3} dy_2 = \frac{4}{3} \cdot \frac{1}{4} = \frac{1}{3}$$

□

*Exercise (5.32).* Suppose that the random variables  $Y_1$  and  $Y_2$  have joint probability density function,  $f(y_1, y_2)$ , given by (see Exercise 5.14)

$$f(y_1, y_2) = \begin{cases} 6y_1^2 y_2, & 0 \leq y_1 \leq y_2, y_1 + y_2 \leq 2, \\ 0, & \text{elsewhere.} \end{cases}$$

- (a) Show that the marginal density of  $Y_1$  is a beta density with  $\alpha = 3$  and  $\beta = 2$ .

*Solution:*

$$f_1(y_1) = \int_{y_1}^{2-y_1} 6y_1^2 y_2 dy_2 = 3y_1^2 [y_2^2]_{y_1}^{2-y_1} = 3y_1^2 [(2-y_1)^2 - y_1^2]$$

$$= 3y_1^2 [4 - 4y_1] = 12y_1^2 (1 - y_1), \quad 0 \leq y_1 \leq 1$$

Beta( $\alpha = 3, \beta = 2$ ) has density  $\frac{\Gamma(5)}{\Gamma(3)\Gamma(2)} y^2 (1-y) = \frac{24}{2} y^2 (1-y) = 12y^2 (1-y)$ . □

- (b) Derive the marginal density of  $Y_2$ .

*Solution:*

$$f_2(y_2) = \begin{cases} \int_0^{y_2} 6y_1^2 y_2 dy_1 = 2y_2 y_1^3 \Big|_0^{y_2} = 2y_2^4, & 0 \leq y_2 \leq 1 \\ \int_0^{2-y_2} 6y_1^2 y_2 dy_1 = 2y_2 (2-y_2)^3, & 1 < y_2 \leq 2 \end{cases}$$

□

- (c) Derive the conditional density of  $Y_2$  given  $Y_1 = y_1$ .

*Solution:*

$$f(y_2|y_1) = \frac{6y_1^2 y_2}{12y_1^2 (1 - y_1)} = \frac{y_2}{2(1 - y_1)}, \quad y_1 \leq y_2 \leq 2 - y_1$$

□

- (d) Find  $P(Y_2 < 1.1|Y_1 = .60)$ .

*Solution:*

$$f(y_2|y_1 = 0.6) = \frac{y_2}{2(0.4)} = 1.25y_2, \quad 0.6 \leq y_2 \leq 1.4$$

$$P(Y_2 < 1.1|Y_1 = 0.6) = \int_{0.6}^{1.1} 1.25y_2 dy_2 = 0.625[y_2^2]_{0.6}^{1.1} = 0.625(1.21 - 0.36) = 0.53125$$

□

*Exercise (5.33).* Suppose that  $Y_1$  is the total time between a customer's arrival in the store and departure from the service window,  $Y_2$  is the time spent in line before reaching the window, and the joint density of these variables (as was given in Exercise 5.15) is

$$f(y_1, y_2) = \begin{cases} e^{-y_1}, & 0 \leq y_2 \leq y_1 \leq \infty, \\ 0, & \text{elsewhere.} \end{cases}$$

- (a) Find the marginal density functions for  $Y_1$  and  $Y_2$ .

*Solution:*

$$f_1(y_1) = \int_0^{y_1} e^{-y_1} dy_2 = y_1 e^{-y_1}, \quad y_1 > 0$$

$$f_2(y_2) = \int_{y_2}^{\infty} e^{-y_1} dy_1 = -e^{-y_1} \Big|_{y_2}^{\infty} = e^{-y_2}, \quad y_2 > 0$$

□

- (b) What is the conditional density function of  $Y_1$  given that  $Y_2 = y_2$ ? Be sure to specify the values of  $y_2$  for which this conditional density is defined.

*Solution:*

$$f(y_1|y_2) = \frac{e^{-y_1}}{e^{-y_2}} = e^{-(y_1-y_2)}, \quad y_1 \geq y_2$$

Defined for  $y_2 > 0$ .

□

- (c) What is the conditional density function of  $Y_2$  given that  $Y_1 = y_1$ ? Be sure to specify the values of  $y_1$  for which this conditional density is defined.

*Solution:*

$$f(y_2|y_1) = \frac{e^{-y_1}}{y_1 e^{-y_1}} = \frac{1}{y_1}, \quad 0 \leq y_2 \leq y_1$$

Defined for  $y_1 > 0$ .

□



- (d) Is the conditional density function  $f(y_1|y_2)$  that you obtained in part (b) the same as the marginal density function  $f_1(y_1)$  found in part (a)?

*Solution:* No.  $f_1(y_1) = y_1 e^{-y_1}$  while  $f(y_1|y_2) = e^{-(y_1+y_2)}$ . □

- (e) What does your answer to part (d) imply about marginal and conditional probabilities that  $Y_1$  falls in any interval?

*Solution:* They are different, so  $Y_1$  and  $Y_2$  are dependent. □

*Exercise (5.35).* Refer to Exercise 5.33. If two minutes elapse between a customer's arrival at the store and his departure from the service window, find the probability that he waited in line less than one minute to reach the window.

*Solution:* From 5.33(c),  $f(y_2|y_1) = \frac{1}{y_1}$  for  $0 \leq y_2 \leq y_1$ .

$$P(Y_2 < 1 | Y_1 = 2) = \int_0^1 \frac{1}{2} dy_2 = \frac{1}{2}$$

□

*Exercise (5.37).* In Exercise 5.18,  $Y_1$  and  $Y_2$  denoted the lengths of life, in hundreds of hours, for components of types I and II, respectively, in an electronic system. The joint density of  $Y_1$  and  $Y_2$  is given by

$$f(y_1, y_2) = \begin{cases} (1/8)y_1 e^{-(y_1+y_2)/2}, & y_1 > 0, y_2 > 0 \\ 0, & \text{elsewhere.} \end{cases}$$

Find the probability that a component of type II will have a life length in excess of 200 hours.

*Solution:*

$$\begin{aligned} f_2(y_2) &= \int_0^\infty \frac{1}{8} y_1 e^{-(y_1+y_2)/2} dy_1 = \frac{1}{8} e^{-y_2/2} \int_0^\infty y_1 e^{-y_1/2} dy_1 \\ &= \frac{1}{8} e^{-y_2/2} \cdot 4 = \frac{1}{2} e^{-y_2/2} \\ P(Y_2 > 2) &= \int_2^\infty \frac{1}{2} e^{-y_2/2} dy_2 = -e^{-y_2/2} \Big|_2^\infty = e^{-1} \approx 0.368 \end{aligned}$$

□

#### SECTION 5.4: INDEPENDENT RANDOM VARIABLES

*Exercise (5.43).* Let  $Y_1$  and  $Y_2$  have joint density function  $f(y_1, y_2)$  and marginal densities  $f_1(y_1)$  and  $f_2(y_2)$ , respectively. Show that  $Y_1$  and  $Y_2$  are independent if and only if  $f(y_1|y_2) = f_1(y_1)$  for all values of  $y_1$  and for all  $y_2$  such that  $f_2(y_2) > 0$ . A completely analogous argument

establishes that  $Y_1$  and  $Y_2$  are independent if and only if  $f(y_2|y_1) = f_2(y_2)$  for all values of  $y_2$  and for all  $y_1$  such that  $f_1(y_1) > 0$ .

*Solution:* ( $\Rightarrow$ ) If independent,  $f(y_1, y_2) = f_1(y_1)f_2(y_2)$ , so  $f(y_1|y_2) = \frac{f(y_1, y_2)}{f_2(y_2)} = \frac{f_1(y_1)f_2(y_2)}{f_2(y_2)} = f_1(y_1)$ .

( $\Leftarrow$ ) If  $f(y_1|y_2) = f_1(y_1)$ , then  $\frac{f(y_1, y_2)}{f_2(y_2)} = f_1(y_1)$ , so  $f(y_1, y_2) = f_1(y_1)f_2(y_2)$ , which means independent.  $\square$

*Exercise (5.45).* In Exercise 5.1, we determined that the joint distribution of  $Y_1$ , the number of contracts awarded to firm A, and  $Y_2$ , the number of contracts awarded to firm B, is given by the entries in the following table.

$y_1$	$y_2$		
	0	1	2
0	1/9	2/9	1/9
1	2/9	2/9	0
2	1/9	0	0

The marginal probability function of  $Y_1$  was derived in Exercise 5.19 to be binomial with  $n = 2$  and  $p = 1/3$ . Are  $Y_1$  and  $Y_2$  independent? Why?

*Solution:* No. For independence we need  $p(y_1, y_2) = p_1(y_1)p_2(y_2)$  for all  $(y_1, y_2)$ . We have

$$p_1(2) = \binom{2}{2}(1/3)^2 = 1/9 \text{ and } p(2, 0) = 1/9. \text{ If independent, } p_2(0) = 1. \text{ But } p(0, 0) = 1/9 \neq p_1(0)p_2(0) = (4/9)(1) = 4/9. \text{ Not independent. } \square$$

*Exercise (5.47).* In Exercise 5.3, we determined that the joint probability distribution of  $Y_1$ , the number of married executives, and  $Y_2$ , the number of never-married executives, is given by

$$p(y_1, y_2) = \frac{\binom{4}{y_1}\binom{3}{y_2}\binom{2}{3-y_1-y_2}}{\binom{9}{3}},$$

where  $y_1$  and  $y_2$  are integers,  $0 \leq y_1 \leq 3$ ,  $0 \leq y_2 \leq 3$ , and  $1 \leq y_1 + y_2 \leq 3$ . Are  $Y_1$  and  $Y_2$  independent? (Recall your answer to Exercise 5.21.)

*Solution:* No. Sampling without replacement, so dependent.  $\square$

*Exercise (5.49).* In Example 5.4 and Exercise 5.5, we considered the joint density of  $Y_1$ , the proportion of the capacity of the tank that is stocked at the beginning of the week and  $Y_2$ , the proportion of the capacity sold during the week, given by

$$f(y_1, y_2) = \begin{cases} 3y_1, & 0 \leq y_2 \leq y_1 \leq 1, \\ 0, & \text{elsewhere.} \end{cases}$$

Show that  $Y_1$  and  $Y_2$  are dependent.

*Solution:* From 5.23:  $f_1(y_1) = 3y_1^2$ ,  $f_2(y_2) = \frac{3}{2}(1 - y_2^2)$ . Since  $f(y_1, y_2) = 3y_1 \neq f_1(y_1)f_2(y_2)$ , dependent.  $\square$

*Exercise (5.51).* In Exercise 5.7, we considered  $Y_1$  and  $Y_2$  with joint density function

$$f(y_1, y_2) = \begin{cases} e^{-(y_1+y_2)}, & y_1 > 0, y_2 > 0, \\ 0, & \text{elsewhere.} \end{cases}$$

(a) Are  $Y_1$  and  $Y_2$  independent?

*Solution:* Yes.  $f(y_1, y_2) = e^{-(y_1+y_2)} = e^{-y_1} \cdot e^{-y_2} = f_1(y_1)f_2(y_2)$  from 5.25.  $\square$

(b) Does the result from part (a) explain the results you obtained in Exercise 5.25 (d)–(f)? Why?

*Solution:* Yes. Independence implies  $f(y_1|y_2) = f_1(y_1)$ , which is what we found in 5.25.  $\square$

*Exercise (5.53).* In Exercise 5.9, we determined that

$$f(y_1, y_2) = \begin{cases} 6(1 - y_2), & 0 \leq y_1 \leq y_2 \leq 1, \\ 0, & \text{elsewhere} \end{cases}$$

is a valid joint probability density function. Are  $Y_1$  and  $Y_2$  independent?

*Solution:* No. Support region depends on both variables.  $\square$

*Exercise (5.55).* Suppose that, as in Exercise 5.11,  $Y_1$  and  $Y_2$  are uniformly distributed over the triangle shaded in the accompanying diagram. Are  $Y_1$  and  $Y_2$  independent?

*Solution:* No. Support region depends on both variables.  $\square$

*Exercise (5.57).* In Exercises 5.13 and 5.31, the joint density function of  $Y_1$  and  $Y_2$  was given by

$$f(y_1, y_2) = \begin{cases} 30y_1y_2^2, & y_1 - 1 \leq y_2 \leq 1 - y_1, 0 \leq y_1 \leq 1, \\ 0, & \text{elsewhere.} \end{cases}$$

Are the random variables  $Y_1$  and  $Y_2$  independent?

*Solution:* No. Support region depends on both variables.  $\square$

*Exercise (5.59).* If  $Y_1$  is the total time between a customer's arrival in the store and leaving the service window and if  $Y_2$  is the time spent in line before reaching the window, the joint density of these variables, according to Exercise 5.15, is

$$f(y_1, y_2) = \begin{cases} e^{-y_1}, & 0 \leq y_2 \leq y_1 \leq \infty \\ 0, & \text{elsewhere.} \end{cases}$$

Are  $Y_1$  and  $Y_2$  independent?

*Solution:* No. From 5.33,  $f(y_1|y_2) \neq f_1(y_1)$ , so dependent.  $\square$

*Exercise (5.61).* In Exercise 5.18,  $Y_1$  and  $Y_2$  denoted the lengths of life, in hundreds of hours, for components of types I and II, respectively, in an electronic system. The joint density of  $Y_1$  and  $Y_2$  is

$$f(y_1, y_2) = \begin{cases} (1/8)y_1 e^{-(y_1+y_2)/2}, & y_1 > 0, y_2 > 0, \\ 0, & \text{elsewhere.} \end{cases}$$

Are  $Y_1$  and  $Y_2$  independent?

*Solution:* No. Cannot factor as  $g(y_1)h(y_2)$  due to  $y_1$  term.  $\square$

*Exercise (5.63).* Let  $Y_1$  and  $Y_2$  be independent exponentially distributed random variables, each with mean 1. Find  $P(Y_1 > Y_2 | Y_1 < 2Y_2)$ .

*Solution:*

$$\begin{aligned} P(Y_1 > Y_2, Y_1 < 2Y_2) &= \int_0^\infty \int_{y_2}^{2y_2} e^{-y_1} e^{-y_2} dy_1 dy_2 = \int_0^\infty e^{-y_2} (e^{-y_2} - e^{-2y_2}) dy_2 \\ &= \int_0^\infty (e^{-2y_2} - e^{-3y_2}) dy_2 = \frac{1}{2} - \frac{1}{3} = \frac{1}{6} \\ P(Y_1 < 2Y_2) &= \int_0^\infty \int_0^{2y_2} e^{-y_1} e^{-y_2} dy_1 dy_2 = \int_0^\infty e^{-y_2} (1 - e^{-2y_2}) dy_2 = 1 - \frac{1}{3} = \frac{2}{3} \\ P(Y_1 > Y_2 | Y_1 < 2Y_2) &= \frac{1/6}{2/3} = \frac{1}{4} \end{aligned}$$

$\square$

## SECTION 5.6: EXPECTED VALUE OF A FUNCTION OF RANDOM VARIABLES

*Exercise (5.73).* In Exercise 5.3, we determined that the joint probability distribution of  $Y_1$ , the number of married executives, and  $Y_2$ , the number of never-married executives, is given by

$$p(y_1, y_2) = \frac{\binom{4}{y_1} \binom{3}{y_2} \binom{2}{3-y_1-y_2}}{\binom{9}{3}},$$

where  $y_1$  and  $y_2$  are integers,  $0 \leq y_1 \leq 3$ ,  $0 \leq y_2 \leq 3$ , and  $1 \leq y_1 + y_2 \leq 3$ . Find the expected number of married executives among the three selected for promotion. (See Exercise 5.21.)

*Solution:* Hypergeometric:  $\mathbb{E}(Y_1) = n \cdot \frac{r}{N} = 3 \cdot \frac{4}{9} = \frac{4}{3}$ .  $\square$

*Exercise (5.75).* Refer to Exercises 5.7, 5.25, and 5.51. Let  $Y_1$  and  $Y_2$  have joint density function

$$f(y_1, y_2) = \begin{cases} e^{-(y_1+y_2)}, & y_1 > 0, y_2 > 0 \\ 0, & \text{elsewhere.} \end{cases}$$

- (a) What are  $E(Y_1 + Y_2)$  and  $V(Y_1 + Y_2)$ ?

*Solution:*  $Y_1, Y_2$  independent  $\text{Exp}(1)$ , so  $\mathbb{E}(Y_1) = \mathbb{E}(Y_2) = 1$ ,  $V(Y_1) = V(Y_2) = 1$ .

$$\mathbb{E}(Y_1 + Y_2) = 2, \quad V(Y_1 + Y_2) = 2$$

□

- (b) What is  $P(Y_1 - Y_2 > 3)$ ?

*Solution:*

$$P(Y_1 - Y_2 > 3) = \int_3^\infty \int_0^{y_1-3} e^{-y_1} e^{-y_2} dy_2 dy_1 = \int_3^\infty e^{-y_1} (1 - e^{-(y_1-3)}) dy_1 = e^{-3}$$

□

- (c) What is  $P(Y_1 - Y_2 < -3)$ ?

*Solution:* By symmetry,  $P(Y_1 - Y_2 < -3) = P(Y_2 - Y_1 > 3) = e^{-3}$ .

□

- (d) What are  $E(Y_1 - Y_2)$  and  $V(Y_1 - Y_2)$ ?

*Solution:*

$$\mathbb{E}(Y_1 - Y_2) = 0, \quad V(Y_1 - Y_2) = 2$$

□

- (e) What do you notice about  $V(Y_1 + Y_2)$  and  $V(Y_1 - Y_2)$ ?

*Solution:* They are equal: both equal 2.

□

*Exercise (5.77).* In Exercise 5.9, we determined that

$$f(y_1, y_2) = \begin{cases} 6(1 - y_2), & 0 \leq y_1 \leq y_2 \leq 1, \\ 0, & \text{elsewhere} \end{cases}$$

is a valid joint probability density function. Find

- (a)  $E(Y_1)$  and  $E(Y_2)$ .

*Solution:* From 5.27:  $f_1(y_1) = 3(1 - y_1)^2$  and  $f_2(y_2) = 6y_2(1 - y_2)$ .

$$\mathbb{E}(Y_1) = \int_0^1 y_1 \cdot 3(1 - y_1)^2 dy_1 = 3 \int_0^1 (y_1 - 2y_1^2 + y_1^3) dy_1 = 3\left[\frac{1}{2} - \frac{2}{3} + \frac{1}{4}\right] = \frac{1}{4}$$

$$\mathbb{E}(Y_2) = \int_0^1 y_2 \cdot 6y_2(1 - y_2) dy_2 = 6 \int_0^1 (y_2^2 - y_2^3) dy_2 = 6\left[\frac{1}{3} - \frac{1}{4}\right] = \frac{1}{2}$$

□

(b)  $V(Y_1)$  and  $V(Y_2)$ .

*Solution:*

$$\mathbb{E}(Y_1^2) = 3 \int_0^1 y_1^2(1 - y_1)^2 dy_1 = 3 \int_0^1 (y_1^2 - 2y_1^3 + y_1^4) dy_1 = 3\left[\frac{1}{3} - \frac{1}{2} + \frac{1}{5}\right] = \frac{1}{10}$$

$$V(Y_1) = \frac{1}{10} - \frac{1}{16} = \frac{3}{80}$$

$$\mathbb{E}(Y_2^2) = 6 \int_0^1 y_2^3(1 - y_2) dy_2 = 6\left[\frac{1}{4} - \frac{1}{5}\right] = \frac{3}{10}$$

$$V(Y_2) = \frac{3}{10} - \frac{1}{4} = \frac{1}{20}$$

□

(c)  $E(Y_1 - 3Y_2)$ .

*Solution:*

$$\mathbb{E}(Y_1 - 3Y_2) = \frac{1}{4} - 3 \cdot \frac{1}{2} = -\frac{5}{4}$$

□

*Exercise (5.79).* Suppose that, as in Exercise 5.11,  $Y_1$  and  $Y_2$  are uniformly distributed over the triangle shaded in the accompanying diagram with vertices at  $(-1, 0)$ ,  $(1, 0)$ , and  $(0, 1)$ . Find  $E(Y_1 Y_2)$ .

*Solution:*

$$\begin{aligned} \mathbb{E}(Y_1 Y_2) &= \int_{-1}^0 \int_0^{1+y_1} y_1 y_2 dy_2 dy_1 + \int_0^1 \int_0^{1-y_1} y_1 y_2 dy_2 dy_1 \\ &= \int_{-1}^0 \frac{y_1(1+y_1)^2}{2} dy_1 + \int_0^1 \frac{y_1(1-y_1)^2}{2} dy_1 = 0 \end{aligned}$$

By symmetry around  $y_1 = 0$ .

□

*Exercise (5.81).* In Exercise 5.18,  $Y_1$  and  $Y_2$  denoted the lengths of life, in hundreds of hours, for components of types I and II, respectively, in an electronic system. The joint density of  $Y_1$  and  $Y_2$  is

$$f(y_1, y_2) = \begin{cases} (1/8)y_1 e^{-(y_1+y_2)/2}, & y_1 > 0, y_2 > 0, \\ 0, & \text{elsewhere.} \end{cases}$$

One way to measure the relative efficiency of the two components is to compute the ratio  $Y_2/Y_1$ . Find  $E(Y_2/Y_1)$ . [Hint: In Exercise 5.61, we proved that  $Y_1$  and  $Y_2$  are independent.]

*Solution:* Not independent (from 5.61), compute directly:

$$\begin{aligned} \mathbb{E}(Y_2/Y_1) &= \int_0^\infty \int_0^\infty \frac{y_2}{y_1} \cdot \frac{1}{8} y_1 e^{-(y_1+y_2)/2} dy_2 dy_1 \\ &= \frac{1}{8} \int_0^\infty \int_0^\infty y_2 e^{-(y_1+y_2)/2} dy_2 dy_1 = \frac{1}{8} \cdot 4 \cdot 4 = 2 \end{aligned}$$

□

*Exercise (5.87).* Suppose that  $Y_1$  and  $Y_2$  are independent  $\chi^2$  random variables with  $\nu_1$  and  $\nu_2$  degrees of freedom, respectively. Find

(a)  $E(Y_1 + Y_2)$ .

*Solution:*

$$\mathbb{E}(Y_1 + Y_2) = \mathbb{E}(Y_1) + \mathbb{E}(Y_2) = \nu_1 + \nu_2$$

□

(b)  $V(Y_1 + Y_2)$ . [Hint: Use Theorem 5.9 and the result of Exercise 4.112(a).]

*Solution:*

$$V(Y_1 + Y_2) = V(Y_1) + V(Y_2) = 2\nu_1 + 2\nu_2$$

□

## SECTION 5.7: THE COVARIANCE OF TWO RANDOM VARIABLES

*Exercise (5.89).* In Exercise 5.1, we determined that the joint distribution of  $Y_1$ , the number of contracts awarded to firm A, and  $Y_2$ , the number of contracts awarded to firm B, is given by the entries in the following table.

$y_1$	$y_2$		
	0	1	2
0	1/9	2/9	1/9
1	2/9	2/9	0
2	1/9	0	0

Find  $\text{Cov}(Y_1, Y_2)$ . Does it surprise you that  $\text{Cov}(Y_1, Y_2)$  is negative? Why?

*Solution:*

$$\mathbb{E}(Y_1) = 0(4/9) + 1(4/9) + 2(1/9) = 2/3$$

$$\mathbb{E}(Y_2) = 0(4/9) + 1(4/9) + 2(1/9) = 2/3$$

$$\mathbb{E}(Y_1 Y_2) = \sum y_1 y_2 p(y_1, y_2) = 0(1/9) + 0(2/9) + 0(1/9) + 0(2/9) + 1(2/9) + 0 = 2/9$$

$$\text{Cov}(Y_1, Y_2) = \mathbb{E}(Y_1 Y_2) - \mathbb{E}(Y_1)\mathbb{E}(Y_2) = 2/9 - (2/3)(2/3) = 2/9 - 4/9 = -2/9$$

Not surprising: more contracts to one firm means fewer to the other (negative association).  $\square$

*Exercise (5.91).* In Exercise 5.8, we derived the fact that

$$f(y_1, y_2) = \begin{cases} 4y_1 y_2, & 0 \leq y_1 \leq 1, 0 \leq y_2 \leq 1, \\ 0, & \text{elsewhere.} \end{cases}$$

Show that  $\text{Cov}(Y_1, Y_2) = 0$ . Does it surprise you that  $\text{Cov}(Y_1, Y_2)$  is zero? Why?

*Solution:*  $f(y_1, y_2) = 4y_1 y_2 = (2y_1)(2y_2) = f_1(y_1)f_2(y_2)$ , so independent. Thus  $\text{Cov}(Y_1, Y_2) = 0$ . Not surprising: independence implies zero covariance.  $\square$

*Exercise (5.92).* In Exercise 5.9, we determined that

$$f(y_1, y_2) = \begin{cases} 6(1 - y_2), & 0 \leq y_1 \leq y_2 \leq 1, \\ 0, & \text{elsewhere} \end{cases}$$

is a valid joint probability density function. Find  $\text{Cov}(Y_1, Y_2)$ . Are  $Y_1$  and  $Y_2$  independent?

*Solution:* From 5.77:  $\mathbb{E}(Y_1) = 1/4$ ,  $\mathbb{E}(Y_2) = 1/2$ .

$$\begin{aligned} \mathbb{E}(Y_1 Y_2) &= \int_0^1 \int_0^{y_2} y_1 y_2 \cdot 6(1 - y_2) dy_1 dy_2 = 6 \int_0^1 y_2(1 - y_2) \frac{y_2^2}{2} dy_2 \\ &= 3 \int_0^1 (y_2^3 - y_2^4) dy_2 = 3 \left[ \frac{1}{4} - \frac{1}{5} \right] = \frac{3}{20} \\ \text{Cov}(Y_1, Y_2) &= \frac{3}{20} - \frac{1}{4} \cdot \frac{1}{2} = \frac{3}{20} - \frac{1}{8} = \frac{1}{40} \end{aligned}$$

Not independent (from 5.53).  $\square$

*Exercise (5.93).* Let the discrete random variables  $Y_1$  and  $Y_2$  have the joint probability function

$$p(y_1, y_2) = 1/3, \text{ for } (y_1, y_2) = (-1, 0), (0, 1), (1, 0).$$



Find  $\text{Cov}(Y_1, Y_2)$ . Notice that  $Y_1$  and  $Y_2$  are dependent. (Why?) This is another example of uncorrelated random variables that are not independent.

*Solution:*

$$\mathbb{E}(Y_1) = \frac{1}{3}[(-1) + 0 + 1] = 0$$

$$\mathbb{E}(Y_2) = \frac{1}{3}[0 + 1 + 0] = \frac{1}{3}$$

$$\mathbb{E}(Y_1 Y_2) = \frac{1}{3}[(-1)(0) + (0)(1) + (1)(0)] = 0$$

$$\text{Cov}(Y_1, Y_2) = 0 - 0 \cdot \frac{1}{3} = 0$$

Dependent because support is not a rectangle, but uncorrelated.  $\square$

*Exercise (5.95).* Suppose that, as in Exercises 5.11 and 5.79,  $Y_1$  and  $Y_2$  are uniformly distributed over the triangle shaded in the accompanying diagram with vertices at  $(-1, 0)$ ,  $(1, 0)$ , and  $(0, 1)$ .

(a) Find  $\text{Cov}(Y_1, Y_2)$ .

*Solution:* From 5.29:  $f_1(y_1) = 1 - |y_1|$ ,  $f_2(y_2) = 2(1 - y_2)$ . By symmetry,  $\mathbb{E}(Y_1) = 0$ .

$$\mathbb{E}(Y_2) = 2 \int_0^1 y_2(1 - y_2) dy_2 = 2\left[\frac{1}{2} - \frac{1}{3}\right] = \frac{1}{3}$$

$$\text{Cov}(Y_1, Y_2) = \mathbb{E}(Y_1 Y_2) - \mathbb{E}(Y_1)\mathbb{E}(Y_2) = 0 - 0 = 0 \text{ (from 5.79)}$$

$\square$

(b) Are  $Y_1$  and  $Y_2$  independent? (See Exercise 5.55.)

*Solution:* No (from 5.55).  $\square$

(c) Find the coefficient of correlation for  $Y_1$  and  $Y_2$ .

*Solution:*

$$\rho = \frac{\text{Cov}(Y_1, Y_2)}{\sqrt{V(Y_1)V(Y_2)}} = \frac{0}{\sqrt{V(Y_1)V(Y_2)}} = 0$$

$\square$

(d) Does your answer to part (b) lead you to doubt your answer to part (a)? Why or why not?

*Solution:* No. Dependent variables can have zero covariance.  $\square$

*Exercise* (5.99). If  $c$  is any constant and  $Y$  is a random variable such that  $E(Y)$  exists, show that  $\text{Cov}(c, Y) = 0$ .

*Solution:*

$$\text{Cov}(c, Y) = \mathbb{E}(cY) - \mathbb{E}(c)\mathbb{E}(Y) = c\mathbb{E}(Y) - c\mathbb{E}(Y) = 0$$

□