4.4.2

a) We are provided the second order differential equation $y'' - 9y' = 3x^2 - 5\sin(3x)$ and we are tasked with finding the form of the particular solution, by first inspection/observation we find the form to be:

$$y_p = (Ax^2 + Bx + C) + D\cos(3x) + E\sin(3x)$$
 (1)

b) We are also provided with the second order differential equation $y'' + 2y' + y = 2e^{-x} - e^x$, by inspection/observation we conclude that the form of the particular solution is:

$$y_p = Ae^{-x} + Be^x$$

4.4.3

We are tasked with solving the following differential equations via method of undetermined coefficients:

a)
$$y'' + 2y' = 2x + 5 - e^{-2x}$$

$$y'' + 2y' = 2x + 5 - e^{-2x}$$

$$m^{2} + 2m + 0 = 0$$

$$y'_{p} = 2Ax + B - 2Cxe^{-2x} + Ce^{-2x}$$

$$y''_{p} = 2Ax + B - 2Cxe^{-2x} + Ce^{-2x}$$

$$y''_{p} = 2A + 4Cxe^{-2x} - 4Ce^{-2x}$$

$$y''_{p} = 2A + 4Cxe^{-2x} - 4Ce^{-2x}$$

$$2A + 4Cxe^{-2x} - 4Ce^{-2x} + 4Ax + 2B - 4Cxe^{-2x} + 2Ce^{-2x} = 2x + 5 - e^{-2x}$$
$$4Ax + 2(A+B) + 2Ce^{-2x} = 2x + 5 - e^{-2x}$$
$$A = \frac{1}{2}, B = 2, C = \frac{1}{2}$$

Our solution is:

$$y_c + y_p = c_1 + c_2 e^{-2x} + \frac{1}{2}x^2 + 2x + \frac{1}{2}xe^{-2x}$$

b)
$$y'' - 9y' = 2e^{3x}$$
 $y_p = Ae^{3x}$ $y_p = Ae^{3x}$ $y_p = 3Ae^{3x}$ $y'_p = 3Ae^{3x}$ $y'_p = 3Ae^{3x}$ $y''_p = 9Ae^{3x}$ $y''_p = 9Ae^{3x$

Our solution is then:

$$y_c + y_p = c_1 + c_2 e^{9x} - \frac{1}{9} e^{3x}$$

c)
$$y'' + 4y' + 4y = (3+x)e^{-2x}$$

 $y'' + 4y' + 4y = (3+x)e^{-2x}$
 $y_p = Ax^2e^{-2x} + Bx^3e^{-2x}$
 $y_p = Ax^2e^{-2x} + 3Bx^2e^{-2x}$
 $y'_p = 2Axe^{-2x} - 2Ax^2e^{-2x} + 3Bx^2e^{-2x} - 2Bx^3e^{-2x}$
 $(m+2)(m+2) = 0$
 $y''_p = 2Ae^{-2x} + 4Ae^{-2x} - 8Ae^{-2x}$
 $y''_p = 2Ae^{-2x} + 4Ae^{-2x} - 8Ae^{-2x}$

Why did you choose to torture me like this, here is the nasty after plugging in:

$$\begin{split} 2Ae^{-2x} + 4Ae^{-2x} - 8Ae^{-2x} + 4Bx^3e^{-2x} - 12Bx^2e^{-2x} + 6Bxe^{-2x} \\ + 8Axe^{-2x} - 8Ax^2e^{-2x} + 12Bx^2e^{-2x} - 8Bx^3e^{-2x} \\ + Ax^2e^{-2x} + Bx^3e^{-2x} \\ = 3e^{-2x} + xe^{-2x} \end{split}$$

Simplifies beautifully into:

$$2Ae^{-2x} + 6Bxe^{-2x} = 3e^{-2x} + xe^{-2x}$$
$$A = \frac{3}{2}, B = \frac{1}{6}$$

Our solution is then:

$$y_c + y_p = c_1 e^{-2x} + c_2 x e^{-2x} + \frac{3}{2} x^2 e^{-2x} + \frac{1}{6} x^3 e^{-2x}$$