# **Tuples**

(TU) 
$$\frac{e_1 \Rightarrow v_1 \quad \dots \quad e_k \Rightarrow v_k}{(e_1, \dots, e_k) \Rightarrow (v_1, \dots, v_k)}$$

Lists

$$(LI) \quad \frac{e_1 \Rightarrow v_1}{e_1 :: e_2 \Rightarrow v_1 :: v_2}$$

Global definitions

$$(\mathsf{GD}) \quad \frac{f = e \quad e \Rightarrow v}{f \Rightarrow v}$$

## Local definitions

(LD) 
$$\frac{e_1 \Rightarrow v_1 \qquad e_0[v_1/x] \Rightarrow v_0}{\text{let } x = e_1 \text{ in } e_0 \Rightarrow v_0}$$

#### Function calls

(APP) 
$$\frac{e \Rightarrow \text{fun } x \rightarrow e_0 \quad e_1 \Rightarrow v_1 \quad e_0[v_1/x] \Rightarrow v_0}{e \ e_1 \ \Rightarrow \ v_0}$$

# Pattern Matching

$$(PM) \quad \frac{e_0 \Rightarrow v' \equiv p_i[v_1/x_1, \dots, v_k/x_k] \qquad e_i[v_1/x_1, \dots, v_k/x_k] \Rightarrow v}{\operatorname{match} e_0 \operatorname{with} p_1 \rightarrow e_1 \mid \dots \mid p_m \rightarrow e_m \Rightarrow v}$$

## **Built-in operators**

(OP) 
$$\frac{e_1 \Rightarrow v_1}{e_1 \operatorname{op} e_2 \Rightarrow v} \frac{e_2 \Rightarrow v_2 \quad v_1 \operatorname{op} v_2 \Rightarrow v}{e_1 \operatorname{op} e_2 \Rightarrow v}$$

Unary operators are treated analogously.