

# Operations Research Assignment 1

## Linear programming

1. Show if an linear programming in standard form has an optimal feasible solution, then there exists an optimal BFS.
2. Consider the LP in standard form, let  $F$  be the feasible set and  $\{\delta^{(j)}\}_{j \in \mathbb{J}}$  be the set of edge directions of an BFS  $\bar{x}$ . Show

$$F \subseteq \bar{x} + \text{cone} \left( \left\{ \delta^{(j)} \right\}_{j \in \mathbb{J}} \right).$$

3. The unit simplex over  $\mathbb{R}^n$  is denoted by  $\Delta_n$  and is comprised by all nonnegative vectors whose components sum up to one:

$$\Delta_n := \left\{ x \in \mathbb{R}^n \mid x \geq 0, e^T x = 1 \right\},$$

where  $e$  stands for the  $n$ -length column vector of all ones. Show

(i).  $\Delta_n = \text{convex}(\{e_1, e_2, \dots, e_n\})$ , where  $e_i$  is the  $n$ -length column vector whose  $i$ th component is one while all the others are zeros.

(ii). For any  $\{\alpha_i\}_{i=1}^n \subset \mathbb{R}$ , we have

$$\max \{\alpha_1, \alpha_2, \dots, \alpha_n\} = \max_{x \in \Delta_n} (\alpha_1, \alpha_2, \dots, \alpha_n) x.$$

## Optimality Conditions

4. Show that if  $x^*$  is a global minimizer of  $f$  over  $\Omega$ , and  $x^* \in \Omega' \subset \Omega$ . Then  $x^*$  is a global minimizer of  $f$  over  $\Omega'$ .
5. If  $x^*$  is a local minimizer of  $f$  over  $\Omega$  and is a interior point of  $\Omega$ ,  $\Omega \subset \Omega'$ . Then  $x^*$  is a local minimizer of  $f$  over  $\Omega'$ .
6. Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be of class  $C^3$  and  $x^*$  is a local minimizer of  $f$  over a closed set  $\Omega \subset \mathbb{R}^n$ . Review the first-order necessary condition and the second-order necessary condition. Similarly, state and prove the third-order necessary condition.
7. Consider minimizing  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  over a closed set  $\Omega \subset \mathbb{R}_{++}^n$  and

$$f(x) = - \sum_{i=1}^n \log(x_i) \text{ for } x = (x_1, x_2, \dots, x_n)^T \in \Omega.$$

Then all optimal solutions lie on the boundary of  $\Omega$ .

8. Consider the following linear programming:

$$\min -3x_1 - 2x_2 \quad \text{s.t. } x_1 + x_2 \leq 1, x_1 \geq 0, x_2 \geq 0.$$

Use the first-order necessary condition to find the unique optimality feasible solution.

9. Consider minimizing the differentiable function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  over a closed set  $\Omega$ . For any  $x \in \Omega$ , the feasible directions of  $\Omega$  at  $x$  is given by

$$D(x) := \{d \in \mathbb{R}^n \mid d \neq 0, \exists \varepsilon > 0 \text{ such that } x + \alpha d \in \Omega \text{ for all } 0 < \alpha < \varepsilon\},$$

the tangent direction of  $\Omega$  at  $x$  is given by

$$T(x) := \left\{d \in \mathbb{R}^n \mid \exists \{t_k, d^k\}_{k \geq 0} \subset \mathbb{R}_{++} \times \mathbb{R}^n \text{ with } t_k \downarrow 0, d^k \rightarrow d \text{ as } k \rightarrow \infty \text{ and } x + t_k d^k \in \Omega \text{ for all } k \geq 0\right\}.$$

the descent directions of  $f$  at  $x$  is given by

$$F(x) = \left\{d \in \mathbb{R}^n \mid \exists \varepsilon > 0 \text{ such that } f(x + \alpha d) < f(x) \text{ for all } 0 < \alpha < \varepsilon\right\}.$$

Then

- (a)  $\bar{x} \in \Omega$  is a local minimizer if and only if  $D(\bar{x}) \cap F(\bar{x}) = \emptyset$ .
- (b) For any  $x \in \Omega$ ,  $F_0(x) \subset F(x)$ , where

$$F_0(x) := \{d \in \mathbb{R}^n \mid \nabla f(x)^T d < 0\}.$$

- (c)  $D(\bar{x}) \cap F_0(\bar{x}) = \emptyset$  if  $\bar{x} \in \Omega$  is a local minimizer.
  - (d) For any  $x \in \Omega$ ,  $T(x)$  is closed.
  - (e) For any  $x \in \Omega$ ,  $\text{cl}(D(x)) \subset T(x)$ .
  - (f)  $T(\bar{x}) \cap F_0(\bar{x}) = \emptyset$  if  $\bar{x} \in \Omega$  is a local minimizer.
10. Continue to the above question: If  $\Omega := \{x \in \mathbb{R}^n \mid g_i(x) \leq 0 \text{ for } i = 1, 2, \dots, m\}$ , where  $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$  is differentiable for  $i = 1, 2, \dots, m$ . For any  $x \in \Omega$ , the active index set at  $x$  is given by

$$I(x) := \{i \mid i \in \{1, 2, \dots, m\}, g_i(x) = 0\}.$$

Then

- (a) For any  $x \in \Omega$ ,  $G_0(x) \subset D(x)$ , where

$$G_0(x) := \{d \in \mathbb{R}^n \mid \nabla g_i(x)^T d < 0 \text{ for each } i \in I(x)\}.$$

- (b)  $G_0(\bar{x}) \cap F_0(\bar{x}) = \emptyset$  if  $\bar{x} \in \Omega$  is a local minimizer.