

A sharp positivity--rate frontier for quadratic fidelity of reversible spherical generators

Contents

Abstract	2
1. Introduction	2
1.1 Contributions	3
1.2 Priority boundary	3
1.3 Organization	3
1.4 Theorem hierarchy	4
2. Positive reversible spherical generators and sampled harmonic quotients	4
2.1 Notation	4
Proposition 2.1 (sampled residual and aliases)	5
3. Exact covariance and the two-defect decomposition	5
Theorem 3.1 (exact covariance representation)	5
Proposition 3.2 (weighted adjoints and exact traces)	6
Proof architecture	6
4. The sharp positivity--rate--quadratic-fidelity frontier	6
Theorem 4.1 (sharp trace and product bounds)	6
Proof architecture	6
Remark 4.2 (scope of the frontier)	6
5. Equality geometry and exact extremizers	6
Theorem 5.1 (algebraic equality characterization)	6
5.1 Tangent decomposition	7
Theorem 5.2 (local block identity and tight-frame form)	7
Theorem 5.3 (global reversible assembly)	8
Proposition 5.4 (connected antipodal branch)	8
Corollary 5.5 (degree and classification consequences)	8
Corollary 5.6 (global moment consequence)	8
6. Quantitative near-extremizer stability	9
Theorem 6.1 (master stability budget)	9
Corollary 6.2 (exceptional mass and pointwise transfer)	9
Corollary 6.3 (conditional geometric transfers)	9
Proof architecture	10
7. Matching-order positive constructions in dimensions two and three	10
Theorem 7.1 (regular polygons on $\mathbb{S}^1$)	10
7.2 The reflected adaptive-ring construction on $\mathbb{S}^2$	11
Theorem 7.2 (matching order on $\mathbb{S}^2$)	12
Proposition 7.3 (structured support-preserving robustness)	13
8. Exact examples	14
8.1 Three extremal families in every dimension	14
8.2 Platonic examples in $d = 3$	14
8.3 What the examples do and do not show	14
9. Rejected claims and deterministic counterexamples	14
10. Consequences deliberately outside the flagship theorem	15
Appendix A. Proof dependency graph	15

Appendix B. Theorem-to-source-to-test-to-registry map	17
Appendix C. Numerical and falsification protocol	19
Appendix D. Formalization	19
Appendix E. Reproducibility record	20
Appendix F. Limitations and future work	20
Appendix G. Related-theorem and hypothesis-transfer matrix	21
References	23

Scope note. The frontier, equality, and stability theorems are valid in every ambient dimension $d \geq 2$. The matching constructions in Section 7 are proved only for $d=2,3$. No transport or physical-performance conclusion is made in this manuscript.

Abstract

Let $X = (\Omega_i)_{i \in I}$ be a finite indexed sample in $\mathbb{S}^{d-1}$ with positive masses w_i , and let L be a reversible Markov generator with nonnegative jump rates. Assume that constants and coordinate functions are reproduced exactly:

$$L\mathbf{1} = 0, \quad L\Omega = -(d-1)\Omega.$$

We study the residual of L on sampled trace-free quadratic harmonics, quotienting exactly by the quadratic forms that vanish on X . If $r_{\max}$ is the largest outgoing rate and $\mathfrak{D}_2$ is the operator norm of this alias-correct residual in the sampled $\ell^2(w)$ metric, then

$$\mathfrak{D}_2 r_{\max} \geq d(d-1).$$

The constant is sharp in every dimension. We derive an exact covariance factorization into a scalar squared-loss component, with an exact rate-floor-plus-variance identity, and an orthogonal anisotropy component; we characterize equality by common rate and angular loss, with centered tight-frame conditions in every tangent space in the nonantipodal branch, a separate antipodal branch, and global reversible compatibility. We give regular-simplex, cross-polytope, and hypercube extremizers. We also prove a quantitative stability theorem: small relative slack in the sharp product controls a normalized weighted sum of precisely the two equality defects. Pointwise, edgewise, graphwise, quotient, and frame consequences are stated with the mass, conductance, connectivity, sampling, and conditioning parameters they require. The result is an operator frontier, not a quadrature-design statement or a generic obstruction to signed high-order discretizations. The order is attained by local positive families in dimensions two and three. Regular polygons give the exact product in $d = 2$. In $d = 3$, an explicit reflected adaptive-ring family has

$$r_{\max}(L_h) \leq 64\pi^2 h^{-2}, \quad \mathfrak{D}_2(L_h) \leq \frac{75}{2} h^2,$$

and therefore

$$\frac{3}{32\pi^2} h^2 \leq \inf_{L \in \mathcal{G}_h(64\pi^2)} \mathfrak{D}_2(L) \leq \frac{75}{2} h^2.$$

The three-dimensional robustness proposition is restricted to support-preserving reflected latitude perturbations.

1. Introduction

Positive graph generators are attractive because their semigroups preserve order and mass. On a sampled sphere, exact reproduction of the coordinate functions is also natural: the coordinates form the first nonconstant Laplace–Beltrami eigenspace. These two requirements do not, however, leave arbitrary freedom at the next harmonic degree. This paper identifies the exact finite-dimensional obstruction.

Fix $d \geq 2$. We consider all finite positive reversible generators whose coordinate map is an eigenmap with eigenvalue $-(d-1)$. The next continuous spherical eigenvalue is $-2d$, so the relevant finite residual is

$$R_2 = (L + 2dI)S_2, \quad (S_2 A)_i = \Omega_i^\top A \Omega_i, \quad A \in \text{Sym}_0(d).$$

The essential point is that S_2 need not be injective and its image need not be invariant under L . We therefore take the domain to be the sampled quotient $\text{Sym}_0(d)/\ker S_2$, equipped with the norm induced by S_2 , and allow $R_2 A$ to have components outside $\text{im } S_2$. This convention retains sampling aliases and leakage rather than silently discarding them.

The central theorem is the sharp product inequality

$$\mathfrak{D}_2 r_{\max} \geq d(d-1)$$

together with its equality and stability theory. The proof is finite and deterministic. It uses exact covariance identities forced by the coordinate eigenmap, positivity of the jump rates, and a trace comparison in the sampled metric.

1.1 Contributions

The paper makes the following mathematical contributions.

1. It gives an alias-correct formulation of sampled quadratic fidelity and an exact covariance representation of the residual.
2. It splits the rowwise residual into Frobenius-orthogonal scalar and anisotropic covariance components. The scalar squared-loss moment splits exactly into a rate floor and nonnegative angular-loss variance.
3. It proves the sharp all-dimensional frontier $\mathfrak{D}_2 r_{\max} \geq d(d-1)$, without injectivity of S_2 , connectedness, equal masses, or invariance of the sampled quadratic space.
4. It characterizes equality both algebraically and geometrically. In the nonantipodal branch, each row induces a centered weighted unit-norm tight frame in the tangent space, while reversibility imposes additional global assembly conditions that are not consequences of rowwise feasibility.
5. It exhibits three exact extremal families in every dimension and resolves their sampling kernels explicitly.
6. It proves a normalized near-extremizer budget. Stronger local and global conclusions are derived only under explicit mass, edge-probability, connectivity, sampling-frame, and tangent-frame margins.
7. It constructs matching-order local positive families in $d=2,3$. The circle family is an exact regular-polygon calculation. The spherical family uses shared conductances on adaptive reflected rings, proves every row moment equation at all levels, and gives the explicit constants $R_3 = 64\pi^2$ and $C_3 = 75/2$.

1.2 Priority boundary

The contribution is not the introduction of positive coordinate-exact spherical Laplacians. Izmetiev and Lam already construct local reversible spherical Delaunay Laplacians on geodesic triangulations of $\mathbb{S}^2$, with nonnegative coefficients under their Delaunay condition and exact coordinate eigenvalue -2 (Izmetiev and Lam 2025). Positive meshfree stencils, graphical designs, prescribed graph eigenspaces, eigenpair-preserving Laplacian cones, spherical designs, association schemes, graph-Laplacian convergence, and signed higher-accuracy graph formulas likewise provide important adjacent theory (Seibold 2008; Babecki and Thomas 2022; Babecki and Shiroma 2023; Babecki, Steinerberger, and Thomas 2023; Bannai and Bannai 2009; Martin and Tanaka 2008; García Trillos et al. 2020; Yoon 2025).

The distinction here is the joint theorem: a sampled, alias-correct degree-two operator residual; a sharp universal positivity-rate product; its full equality geometry subject to global detailed balance; and a quantitative stability budget with all transfer parameters exposed. The matching three-dimensional construction additionally derives one globally shared positive stress on an explicit all-level ring family; it is not inferred from local positive-stencil feasibility, spherical quadrature, or a Delaunay transfer. A source-by-source hypothesis comparison and hostile-referee audit is supplied in `PRIORITY_AND_HOSTILE_REFEREE_AUDIT.md`. The priority claim is limited to the focused corpus reviewed there; it is not based on project-specific terminology. Appendix G gives the in-paper variables/hypotheses/conclusions matrix for every compared theorem.

1.3 Organization

Section 2 fixes conventions and the sampled quotient. Section 3 proves the exact covariance and two-defect identities. Section 4 proves the sharp frontier. Section 5 characterizes equality and global assembly. Section 6 gives quantitative stability. Section 7 proves matching constructions in dimensions two and three. Section 8 gives exact extremizers and alias examples. Section 9 records false or overbroad statements ruled out by the theory. The appendices collect the dependency graph, provenance map, falsification protocol, formalization scope, reproducibility record, and limitations.

1.4 Theorem hierarchy

Level	Mathematical role	Principal statement	Status
1	Positive reversible generators and sampled harmonic quotients	Proposition 2.1	accepted P1A
2	Exact covariance and two-defect decomposition	Theorem 3.1 and Proposition 3.2	accepted P1A
3	Sharp positivity–rate–quadratic-fidelity frontier	Theorem 4.1	accepted P1B
4	Equality geometry and exact extremizers	Theorems 5.1–5.3, Proposition 5.4, Corollaries 5.5–5.6, Section 8	accepted P1B–P1C
5	Quantitative near-extremizer stability	Theorem 6.1 and Corollaries 6.2–6.3	accepted P1D
6	Matching-order local positive construction	Theorems 7.1–7.2	accepted P1E for $d=2,3$
7	Generator-lift or negativity consequences	Section 10	deliberately not claimed

2. Positive reversible spherical generators and sampled harmonic quotients

2.1 Notation

Symbol	Meaning	Convention or standing hypothesis
I	finite vertex set	nonempty
d	ambient Euclidean dimension	$d \geq 2$
Ω_i	spherical node	$\Omega_i \in \mathbb{S}^{d-1}$
w_i	stationary mass	$w_i > 0, \sum_i w_i = 1$
γ_{ij}	shared conductance	$\gamma_{ij} = \gamma_{ji} \geq 0, \gamma_{ii} = 0$
a_{ij}	directed jump rate	$a_{ij} = \gamma_{ij}/w_i$
L	negative Markov generator	$(Lf)_i = \sum_{j \neq i} a_{ij}(f_j - f_i)$
$r_i, r_{\max}$	outgoing rate and its maximum	$r_i = \sum_j a_{ij}, r_{\max} = \max_i r_i$
ℓ_{ij}	spherical loss	$1 - \Omega_i \cdot \Omega_j \in [0, 2]$
Δ_{ij}	chord increment	$\Omega_j - \Omega_i$
V	quadratic coefficient space	$\text{Sym}_0(d)$
m	dimension of V	$d(d+1)/2 - 1$
S_2	quadratic sampling map	$(S_2 A)_i = \Omega_i^\top A \Omega_i$
K_X	sampling-alias kernel	$\ker S_2 \subseteq V$
R_2	degree-two residual	$(L + 2dI)S_2$
Q_X	sampled quotient	V/K_X with sampled metric
$\mathfrak{D}_2$	sampled quadratic defect	$\sup_{A \notin K_X} \ R_2 A\ _w / \ S_2 A\ _w$
C_i	row chord covariance	$\sum_j a_{ij} \Delta_{ij} \Delta_{ij}^\top$
Z_i	centered rank-one tensor	$\Omega_i \Omega_i^\top - I/d$
M_i	trace-free residual tensor	$C_i + 2\Omega_i \Omega_i^\top - 2I$
ϵ_i	squared-loss moment	$\sum_j a_{ij} \ell_{ij}^2$
B_i	anisotropy tensor	$M_i - \frac{d}{d-1} \epsilon_i Z_i$
E_ϵ, E_B	weighted defect energies	$\sum_i w_i \epsilon_i^2, \sum_i w_i \ B_i\ _F^2$

The weighted inner product is

$$\langle f, g \rangle_w = \sum_i w_i f_i g_i.$$

Detailed balance implies that L is self-adjoint and negative semidefinite in $\ell^2(w)$. Throughout,

$$L\mathbf{1} = 0, \quad L\Omega = -(d-1)\Omega. \quad (2.1)$$

The second identity is componentwise in $\mathbb{R}^d$.

The quotient $Q_X = V/K_X$ carries the inner product

$$\langle [A], [H] \rangle_S = \langle S_2 A, S_2 H \rangle_w. \quad (2.2)$$

This is a genuine inner product on Q_X . It is not the quotient of the Frobenius metric unless an additional sampling-frame identity happens to hold.

Proposition 2.1 (sampled residual and aliases)

The inclusion $K_X \subseteq \ker R_2$ holds. Hence R_2 induces a well-defined map

$$\bar{R}_2 : Q_X \longrightarrow \ell^2(w), \quad \bar{R}_2[A] = R_2 A,$$

and

$$\mathfrak{D}_2 = \|\bar{R}_2\|_{Q_X \rightarrow \ell^2(w)}. \quad (2.3)$$

Equivalently, $\mathfrak{D}_2^2$ is the largest finite generalized eigenvalue of the deflated Gram pencil

$$G_R = R_2^* R_2, \quad G_S = S_2^* S_2$$

on the Frobenius complement $K_X^{\perp_F} \subseteq V$. No invariance of $\text{im } S_2$ under L is required.

The genuinely sampled exact degree-two space is

$$E_{\text{sample}} = \text{im } S_2 \cap \ker(L + 2dI),$$

while the algebraic exact-form space is $E_{\text{form}} = \ker R_2 \subseteq V$. Then

$$\dim K_X = m - \text{rank } S_2, \quad \dim E_{\text{form}} = m - \text{rank } R_2, \quad \dim E_{\text{sample}} = \text{rank } S_2 - \text{rank } R_2. \quad (2.4)$$

The space E_{form} may contain the alias kernel K_X ; only E_{sample} records genuinely sampled exact modes.

3. Exact covariance and the two-defect decomposition

The coordinate eigenmap fixes the first loss moment in every row:

$$\sum_j a_{ij} \ell_{ij} = d - 1. \quad (3.1)$$

This elementary identity is the point at which the continuous coordinate eigenvalue enters all subsequent estimates.

Theorem 3.1 (exact covariance representation)

For every $A \in \text{Sym}_0(d)$ and every vertex i ,

$$(R_2 A)_i = \langle A, M_i \rangle_F. \quad (3.2)$$

Moreover,

$$\langle M_i, Z_i \rangle_F = \epsilon_i, \quad \|Z_i\|_F^2 = \frac{d-1}{d}, \quad \langle B_i, Z_i \rangle_F = 0, \quad (3.3)$$

and therefore

$$M_i = \frac{d}{d-1} \epsilon_i Z_i + B_i, \quad \|M_i\|_F^2 = \frac{d}{d-1} \epsilon_i^2 + \|B_i\|_F^2. \quad (3.4)$$

If

$$V_i = \sum_j a_{ij} \left(\ell_{ij} - \frac{d-1}{r_i} \right)^2,$$

then

$$\epsilon_i = \frac{(d-1)^2}{r_i} + V_i. \quad (3.5)$$

Thus the scalar part of the quadratic residual is the unavoidable rate term $(d-1)^2/r_i$ plus active-loss variance V_i , while B_i is the orthogonal covariance-anisotropy part.

Proposition 3.2 (weighted adjoints and exact traces)

Use the Frobenius inner product on V and the weighted inner product on samples. The adjoints are

$$S_2^* f = \sum_i w_i f_i Z_i, \quad R_2^* f = \sum_i w_i f_i M_i. \quad (3.6)$$

Consequently,

$$\text{tr } G_S = \frac{d-1}{d}, \quad \text{tr } G_R = \frac{d}{d-1} E_\epsilon + E_B. \quad (3.7)$$

These trace identities use the sampled metric and remain valid when S_2 has a nontrivial kernel.

Proof architecture

Equation (3.2) follows by expanding each sampled quadratic along the chord Δ_{ij} and applying (2.1). Equation (3.3) follows by contracting with Z_i , after which (3.4) is an orthogonal projection. Equation (3.5) is the weighted mean–variance identity with fixed mean (3.1). Finally, (3.6)–(3.7) follow by weighted duality and the row-frame trace formula. Complete proofs are assembled from the P1A source identified in Appendix B.

4. The sharp positivity–rate–quadratic-fidelity frontier

Theorem 4.1 (sharp trace and product bounds)

Every generator satisfying the hypotheses of Section 2 obeys

$$\mathfrak{D}_2^2 \geq \frac{d^2}{(d-1)^2} E_\epsilon + \frac{d}{d-1} E_B. \quad (4.1)$$

In particular,

$$\mathfrak{D}_2 \geq \frac{d}{d-1} \sqrt{E_\epsilon} \geq \frac{d(d-1)}{r_{\max}}, \quad (4.2)$$

and hence

$$\boxed{\mathfrak{D}_2 r_{\max} \geq d(d-1)}. \quad (4.3)$$

The constant is attained for every $d \geq 2$.

Proof architecture

On $K_X^{\perp F}$, compare the trace of the positive Gram operator G_R with $\mathfrak{D}_2^2 G_S$. The exact traces (3.7) give (4.1). Next use (3.5), $V_i \geq 0$, $r_i \leq r_{\max}$, and $\sum_i w_i = 1$ to obtain (4.2). The regular-simplex family in Section 8 attains equality, so the constant cannot be improved.

Remark 4.2 (scope of the frontier)

The theorem is neither a general no-go theorem for high-order graph Laplacians nor a statement about quadrature exactness. It applies to the specified class of nonnegative reversible generators, with exact coordinate eigenvalue, the sampled degree-two quotient norm, and the maximum outgoing rate. Signed formulas and other consistency norms lie outside its hypotheses.

5. Equality geometry and exact extremizers

Theorem 5.1 (algebraic equality characterization)

Equality holds in (4.3) if and only if, for every $i \in I$,

$$r_i = r_{\max}, \quad V_i = 0, \quad B_i = 0. \quad (5.1)$$

Equivalently, with

$$c_* = \frac{d(d-1)}{r_{\max}},$$

one has

$$\epsilon_i = \frac{(d-1)^2}{r_{\max}}, \quad M_i = c_* Z_i \quad \text{for every } i, \quad (5.2)$$

and

$$R_2 = c_* S_2 \quad (5.3)$$

on the full coefficient space and therefore on the sampled quotient.

At equality,

$$\ker R_2 = \ker S_2 = K_X, \quad E_{\text{sample}} = \{0\}. \quad (5.4)$$

Thus equality does not reproduce the continuous degree-two eigenvalue; it produces the smallest positive residual allowed by positivity and the rate cap.

5.1 Tangent decomposition

Fix i , write $u = \Omega_i$, and let

$$P_i = I - uu^\top, \quad \tau_{ij} = P_i \Omega_j.$$

Then

$$\Delta_{ij} = -\ell_{ij}u + \tau_{ij}, \quad \|\tau_{ij}\|^2 = \ell_{ij}(2 - \ell_{ij}), \quad \sum_j a_{ij}\tau_{ij} = 0. \quad (5.5)$$

Define

$$h_i = \sum_j a_{ij}\ell_{ij}\tau_{ij}, \quad T_i = \sum_j a_{ij}\tau_{ij}\tau_{ij}^\top.$$

Theorem 5.2 (local block identity and tight-frame form)

The anisotropy tensor has the exact orthogonal block form

$$B_i = -uh_i^\top - h_iu^\top + T_i - \left(2 - \frac{\epsilon_i}{d-1}\right) P_i, \quad (5.6)$$

and

$$\|B_i\|_F^2 = 2\|h_i\|^2 + \left\|T_i - \left(2 - \frac{\epsilon_i}{d-1}\right) P_i\right\|_F^2. \quad (5.7)$$

If $V_i = 0$, all active edges in row i have the common loss

$$\bar{\ell}_i = \frac{d-1}{r_i}.$$

In the nonantipodal case $0 < \bar{\ell}_i < 2$, let

$$p_{ij} = \frac{a_{ij}}{r_i}, \quad y_{ij} = \frac{\tau_{ij}}{\sqrt{\bar{\ell}_i(2 - \bar{\ell}_i)}}.$$

Then $B_i = 0$ if and only if

$$\sum_j p_{ij}y_{ij} = 0, \quad \sum_j p_{ij}y_{ij}y_{ij}^\top = \frac{1}{d-1}P_i. \quad (5.8)$$

Thus the active tangent directions form a centered weighted unit-norm tight frame. The antipodal case $\bar{\ell}_i = 2$ is governed by the division-free identities (5.6)–(5.7), not by the normalization in (5.8).

Theorem 5.3 (global reversible assembly)

In the nonantipodal branch, put

$$t = 1 - \ell_* \in (-1, 1), \quad p_{ij} = a_{ij}/r_*.$$

Frontier equality is equivalent to one common outgoing rate $r_* = (d-1)/(1-t)$ together with

$$\sum_j p_{ij} = 1, \quad w_i p_{ij} = w_j p_{ji}, \quad p_{ij} > 0 \implies \Omega_i \cdot \Omega_j = t, \quad (5.9)$$

$$\sum_j p_{ij} \Omega_j = t \Omega_i, \quad (5.10)$$

and

$$\sum_j p_{ij} \Omega_j \Omega_j^\top = t^2 \Omega_i \Omega_i^\top + \frac{1-t^2}{d-1} (I - \Omega_i \Omega_i^\top) \quad (5.11)$$

for every vertex i . Conversely, (5.9)–(5.11), with $a_{ij} = r_* p_{ij}$, reconstruct the coordinate eigenmap and all equality conditions.

Equivalently, retaining the Markov and detailed-balance conditions in (5.9), the Gram formulation requires a single matrix $G \succeq 0$ with $G_{ii} = 1$, $\text{rank } G = d$, and $p_{ij} > 0 \implies G_{ij} = t$, together with

$$\sum_j p_{ij} G_{jk} = t G_{ik} \quad (5.12)$$

and, for all i, k, l ,

$$\sum_j p_{ij} G_{jk} G_{jl} = \alpha G_{kl} + (t^2 - \alpha) G_{ik} G_{il}, \quad \alpha = \frac{1-t^2}{d-1}. \quad (5.13)$$

On a connected bidirected support, positive reversible masses exist exactly when the directed rates satisfy the Kolmogorov identity on every oriented cycle. In particular, local row feasibility does not imply the existence of a global reversible spherical generator.

Proposition 5.4 (connected antipodal branch)

If the common equality loss is $\ell_* = 2$ and the active graph is connected, all indexed nodes occupy two antipodal positions $\pm u$, every active edge joins opposite positions, and

$$r_* = \frac{d-1}{2}, \quad R_2 = 2dS_2. \quad (5.14)$$

Conversely, every reversible connected constant-row-rate generator with this geometry is an equality generator. Repeated indices at either antipode are allowed.

Corollary 5.5 (degree and classification consequences)

Every nonantipodal equality row has at least d active neighbors. If it has exactly d , its normalized tangent directions form a regular simplex with equal weights. A distinct nonantipodal complete-support equality configuration is the ambient regular simplex.

For $d = 3$, assume additionally that $0 < \ell_* < 2$, the nodes are the distinct vertices of a strictly convex inscribed polyhedron, the active graph is exactly its one-skeleton, all active directed rates have one common value, and the common vertex degree q satisfies $3 \leq q \leq 5$. Then the configuration is tetrahedral, octahedral, cubical, icosahedral, or dodecahedral. No unrestricted Platonic classification is asserted.

Corollary 5.6 (global moment consequence)

Every connected nonantipodal equality generator satisfies

$$\sum_i w_i \Omega_i = 0, \quad \sum_i w_i \Omega_i \Omega_i^\top = \frac{1}{d} I. \quad (5.15)$$

Thus its stationary node measure is a weighted spherical 2-design. This is a consequence of the reversible equality system; it is neither a standing design hypothesis nor an identification of operator fidelity with quadrature exactness.

6. Quantitative near-extremizer stability

Assume

$$\mathfrak{D}_2 r_{\max} \leq d(d-1)(1+\delta), \quad \delta \geq 0. \quad (6.1)$$

Set

$$n = d-1, \quad a_0 = \frac{n^2}{r_{\max}}, \quad c_0 = \frac{dn}{r_{\max}}, \quad \eta = 2\delta + \delta^2,$$

and define

$$x_i = \frac{\epsilon_i}{a_0}, \quad s_i = \frac{r_{\max}}{r_i} - 1, \quad v_i = \frac{V_i}{a_0}, \quad q_i = x_i - 1 = s_i + v_i. \quad (6.2)$$

The defects s_i, v_i, q_i are nonnegative, and $x_i = 1 + q_i \geq 1$.

Theorem 6.1 (master stability budget)

Under (6.1),

$$\sum_i w_i(2q_i + q_i^2) + \frac{d-1}{da_0^2} \sum_i w_i \|B_i\|_F^2 \leq \eta. \quad (6.3)$$

Consequently,

$$\sum_i w_i(x_i - 1)^2 \leq \eta, \quad \sum_i w_i \left(\frac{r_{\max}}{r_i} - 1 \right)^2 \leq \eta, \quad (6.4)$$

$$\sum_i w_i \|B_i\|_F^2 \leq \frac{d(d-1)^3}{r_{\max}^2} \eta, \quad (6.5)$$

and

$$\sum_i w_i q_i \leq \delta, \quad \sum_i w_i s_i \leq \delta, \quad \sum_i w_i v_i \leq \delta, \quad (6.6)$$

$$\sum_i w_i V_i \leq a_0 \delta, \quad \sum_i w_i V_i^2 \leq a_0^2 \eta. \quad (6.7)$$

The estimates are valid without connectedness, injectivity of S_2 , equal masses, or a uniform lower mass bound.

Corollary 6.2 (exceptional mass and pointwise transfer)

Let

$$H = \eta - \frac{d-1}{da_0^2} \sum_i w_i \|B_i\|_F^2.$$

Then $0 \leq H \leq \eta$ and $\sum_i w_i(2q_i + q_i^2) \leq H$. For every $\rho > 0$,

$$\sum_{i: q_i \geq \rho} w_i \leq \min \left\{ 1, \frac{H}{\rho(2+\rho)} \right\}. \quad (6.8)$$

At an individual vertex,

$$q_i \leq \sqrt{1 + \frac{H}{w_i}} - 1. \quad (6.9)$$

Thus a uniform pointwise conclusion requires an explicit lower bound on the stationary masses.

Corollary 6.3 (conditional geometric transfers)

The master budget admits the following further consequences, each under the named quantitative hypotheses.

- The global shell loss has an explicit second-moment bound in the symmetric directed conductance measure. To control every active edge, one assumes $p_{ij} = a_{ij}/r_i \geq \kappa > 0$; the pointwise bound still displays w_i , and a uniform version requires $w_{\min} > 0$.

- Graphwise propagation continues the hypotheses $p_{ij} \geq \kappa > 0$ and $w_i \geq w_{\min} > 0$ on the connected support. Multiplicative path bounds require the stated small-defect condition; additive path, effective-resistance, Poincaré-gap, and congestion variants retain their corresponding graph parameter.
- The orthogonal block identity (5.7) converts the tensor term of the master budget into weighted control of mixed radial–tangent covariance and tangent anisotropy. Together with the scalar term, it controls the full covariance relative to its stated target and its eigenvalue splitting. Eigenvector control further requires a spectral gap.
- Let $W_X = K_X^{\perp F}$, $E = \text{im } S_2$, and

$$\alpha_X = \inf_{\substack{A \in W_X \\ \|A\|_F = 1}} \|S_2 A\|_w^2 > 0.$$

Define $U : E \rightarrow \ell^2(w)$ by $U(S_2 A) = R_2 A$, and let $\iota_E : E \hookrightarrow \ell^2(w)$ denote inclusion. The lower sampling-frame bound converts the full master budget into quotient-correct operator scalarity, including leakage:

$$\|U - c_0 \iota_E\|_{\text{op}} \leq \sqrt{\frac{K_0}{\alpha_X}}, \quad K_0 = \frac{d(d-1)^3}{r_{\max}^2} \eta. \quad (6.10)$$

- A tangent lower-frame bound gives rotation-quotiented distance to an exact centered weighted tight frame. A positive unit-frame and spherical-neighbor repair further requires nonantipodal shell margins, a lower active probability, and a feature-surjectivity bound.

No parameter-free pointwise, edgewise, graphwise, quotient, or frame-repair statement is inferred from (6.3).

Proof architecture

Insert the near-frontier assumption into the trace inequality (4.1), normalize by a_0 , and use $x_i = 1 + q_i$. This gives (6.3) exactly. The global consequences follow by positivity and elementary moment inequalities. Each stronger transfer in Corollary 6.3 is proved separately using the stated lower margin; the counterexamples in Section 9 show why those margins cannot simply be omitted.

7. Matching-order positive constructions in dimensions two and three

Write

$$d_{\mathbb{S}}(x, y) = \arccos(x \cdot y), \quad h_{\text{fill}}(X) = \sup_{x \in \mathbb{S}^{d-1}} \min_i d_{\mathbb{S}}(x, \Omega_i),$$

$$q_{\text{sep}}(X) = \frac{1}{2} \min_{i \neq j} d_{\mathbb{S}}(\Omega_i, \Omega_j), \quad \rho_X = \frac{h_{\text{fill}}(X)}{q_{\text{sep}}(X)}.$$

For a mesh X_h , let $\mathcal{G}_h(\mathbb{R})$ denote the positive reversible generators on X_h that reproduce constants and coordinates exactly and satisfy $r_{\max} \leq Rh^{-2}$. Any further support restriction will be stated with the construction. The lower half of each estimate below is Theorem 4.1; the work in this section is the matching upper half.

Theorem 7.1 (regular polygons on $\mathbb{S}^1$)

Let $N \geq 5$, put $h = \pi/N$, and set

$$\Omega_k = (\cos(2kh), \sin(2kh)), \quad k \in \mathbb{Z}/N\mathbb{Z}.$$

Then $h_{\text{fill}}(X_h) = q_{\text{sep}}(X_h) = h$ and $\rho_{X_h} = 1$. Take $w_k = 1/N$ and the two nearest-neighbor rates

$$a_{k,k-1} = a_{k,k+1} = \frac{1}{2\{1 - \cos(2h)\}}. \quad (7.1)$$

Then the graph has degree two and active edge angle $2h$, and

$$L\mathbf{1} = 0, \quad L\Omega = -\Omega,$$

$$r_{\max} = \frac{1}{1 - \cos(2h)}, \quad \mathfrak{D}_2 = 2\{1 - \cos(2h)\} = 4\sin^2 h, \quad (7.2)$$

so $\mathfrak{D}_2 r_{\max} = 2$. In particular, with

$$R_2 = \frac{\pi^2}{8}, \quad C_2 = 4, \quad c_2 = \frac{16}{\pi^2},$$

one has

$$\frac{16}{\pi^2} h^2 \leq \inf_{L \in \mathcal{G}_h(R_2)} \mathfrak{D}_2(L) \leq 4h^2. \quad (7.3)$$

Proof The nearest-neighbor sum sends the frequency- m Fourier mode to $2(\cos(2mh) - 1)$ times that mode. With (7.1), the frequency-one eigenvalue is -1 , proving exact coordinate reproduction. The sampled trace-free quadratics are the frequency-two sine and cosine modes; $N \geq 5$ makes their sampled space two-dimensional. Their generator eigenvalue is

$$-\frac{1 - \cos(4h)}{1 - \cos(2h)} = -2\{1 + \cos(2h)\}.$$

Adding the continuous degree-two eigenvalue 4 gives the scalar residual in (7.2). For $0 < h \leq \pi/5$,

$$\frac{2h}{\pi} \leq \sin h \leq h,$$

and hence $1 - \cos(2h) = 2 \sin^2 h \geq 8h^2/\pi^2$. This proves the rate cap and upper bound. Theorem 4.1 gives $\mathfrak{D}_2 \geq 2/r_{\max} \geq 2h^2/R_2$, which is the lower bound in (7.3). $\square$

7.2 The reflected adaptive-ring construction on $\mathbb{S}^2$

We give the construction data before stating the theorem. Fix

$$M_0 = 2^{80}, \quad a_{\text{pole}} = \frac{4}{3}, \quad g = \sqrt{\frac{29}{32}}. \quad (7.4)$$

At level $J \geq 1$, put $M_m = M_0 2^m$ and $S_0 = M_J/8$. For $0 \leq m < J$ define

$$t_m^0 = \frac{M_J}{4\pi} \arcsin(2^{m-J}), \quad k_m = \text{nint}(t_m^0 - a_{\text{pole}} - mg),$$

and put

$$K = \text{nint}(S_0 - a_{\text{pole}} - Jg), \quad t_m = a_{\text{pole}} + k_m + mg, \quad S = a_{\text{pole}} + K + Jg, \quad h = \frac{\pi}{2S}. \quad (7.5)$$

Nearest-integer ties are broken downward. Starting with the north pole and a first ring at latitude $a_{\text{pole}}h$, insert k_0 gaps of size h , one gap of size gh at which the longitude count doubles, then $k_{m+1} - k_m$ ordinary gaps between successive doublings, and finally $K - k_{J-1}$ ordinary gaps before the equator. Reflect through the equator without duplicating the equatorial ring. A ring in the m th band has M_m equally spaced longitudes. The rounding identities give

$$|t_m - t_m^0| \leq \frac{1}{2}, \quad |S - S_0| \leq \frac{1}{2}, \quad \left| \frac{2S}{M_m} \sin \frac{\pi t_m}{2S} - \frac{1}{4} \right| \leq \frac{3}{M_m}. \quad (7.6)$$

Aligned rings use identity longitude matching. At a doubling interface $M : 2M$, put $\alpha = \pi/M$, $c_\alpha = \cos \alpha$, and

$$p = \frac{4c_\alpha^2 + 2c_\alpha - 1}{4c_\alpha(c_\alpha + 1)}, \quad q = \frac{(2c_\alpha + 1)(4c_\alpha^2 + 2c_\alpha - 1)}{8c_\alpha^2(c_\alpha + 1)}. \quad (7.7)$$

For coarse longitude i and fine offset $r = j - 2i$, the seven nonzero entries of the shared radial mask are

$$\begin{array}{c|ccccccc} r & 0 & 2 & -2 & 1 & -1 & 3 & -3 \\ \hline Q_{i,2i+r} & p/2 & (1-p)/4 & (1-p)/4 & q/4 & q/4 & (1-q)/4 & (1-q)/4. \end{array} \quad (7.8)$$

Its row sum is one, every column sum is one half, and its reverse even/odd conditional laws have the same first two cosine moments.

Preliminary symmetric stresses $\Gamma_{ij} = \Gamma_{ji}$ are fixed recursively from the pole. At every ordinary row, two reflected horizontal pairs bracket the target loss by the integer rule

$$t_- = \max\{1, \lfloor \phi_*/(2\delta) \rfloor\}, \quad t_+ = \lceil 2\phi_*/\delta \rceil, \quad (7.9)$$

where $\delta = 2\pi/M$ and

$$\sin(\phi_*/2) = \frac{\sin(h/2)}{\sin \theta}.$$

Given the incoming radial stress, the outgoing radial stress and the two horizontal stresses are the unique solution of the three even moment equations

$$\sum_j \Gamma_{ij}(\Delta_{ij} \cdot e_\theta) = 0, \quad \sum_j \Gamma_{ij} \ell_{ij}(\Delta_{ij} \cdot e_\theta) = 0, \quad (7.10)$$

$$\sum_j \Gamma_{ij} \{(\Delta_{ij} \cdot e_\theta)^2 - (\Delta_{ij} \cdot e_\phi)^2\} = 0. \quad (7.11)$$

At a doubling, the two endpoint rows and their shared radial stress are solved simultaneously in the corresponding 6×6 system using (7.8). For the first ring, put $\delta_0 = 2\pi/M_0$ and choose the two horizontal jump integers nearest to $(3/8)/\delta_0$ and $(9/8)/\delta_0$, with the same downward tie rule. Set every pole-to-first-ring preliminary stress equal to 1, then solve the resulting 3×3 system for the outgoing radial and two horizontal stresses. At the equator use equal reflected incoming stresses and horizontal jump two; reflection cancels (7.10), and the positive horizontal stress enforces (7.11). Longitude reflection cancels every odd- e_ϕ equation.

The exact transition matrix has a removable $M^{-1} = 0$ limit whose inverse has norm below 3 and whose solution has minimum greater than $1/500$. On the correlated box from (7.6), a rational Cauchy estimate gives

$$\|A - A_0\|_{\max} \leq \frac{10^{12}}{M}, \quad \|b - b_0\|_\infty \leq \frac{10^{12}}{M}. \quad (7.12)$$

The polar inverse has norm below 20,000 and its finite solution differs from its positive limiting solution by less than 10^{-6} . For ordinary rows, (7.9) gives the uniform bracket

$$\frac{1}{16} < x_- < \frac{1}{3}, \quad 2 < x_+ < 7, \quad \frac{1}{100}U_- < H_-, H_+ < 100U_-. \quad (7.13)$$

The exact telescoping radial recurrence, including all count changes, yields the level-independent shared-stress margins

$$\Gamma_- = M_0^{-20} \leq \Gamma_{ij} \leq M_0^{20} = \Gamma_+. \quad (7.14)$$

Thus positivity is certified before any normalization; it is not inferred from the finite generator.

Theorem 7.2 (matching order on $\mathbb{S}^2$)

For the all-level family (7.4)–(7.14), there are positive weights and a local reversible generator with

$$L_h \mathbf{1} = 0, \quad L_h \Omega = -2\Omega, \quad \gamma_{ij} = \gamma_{ji} \geq 0. \quad (7.15)$$

The metrics defined above satisfy

$$h_{\text{fill}}(X_h) \leq 2h, \quad q_{\text{sep}}(X_h) \geq q_* h, \quad \rho_{X_h} \leq \frac{2}{q_*}, \quad q_* = (8M_0)^{-1}.$$

Every active edge lies in the angular window $h/8 \leq d_{\mathbb{S}}(\Omega_i, \Omega_j) \leq 5h$, and the degree is at most $D = M_0$. For every vertex,

$$B_i = 0, \quad (R_2 A)_i = c_i (S_2 A)_i, \quad 0 < c_i \leq \frac{75}{2} h^2. \quad (7.16)$$

Consequently,

$$r_{\max}(L_h) \leq 64\pi^2 h^{-2}, \quad \mathfrak{D}_2(L_h) \leq \frac{75}{2} h^2. \quad (7.17)$$

With

$$R_3 = 64\pi^2, \quad C_3 = \frac{75}{2}, \quad c_3 = \frac{3}{32\pi^2},$$

one therefore has, for every level,

$$\frac{3}{32\pi^2}h^2 \leq \inf_{L \in \mathcal{G}_h(\mathbf{R}_3)} \mathfrak{D}_2(L) \leq \frac{75}{2}h^2. \quad (7.18)$$

For the structured robustness proposition below, set

$$K_* = 2^{28 \cdot 2^{106}}, \quad (7.19)$$

Proof The schedule closes exactly because the terminal dimensionless latitude is $a_{\text{pole}} + k_{J-1} + (J-1)g + g + K - k_{J-1} = S$. Equations (7.6), the ring spacing, and the active jumps prove the stated fill, separation, window, degree, and packing bounds. The exact row systems (7.10)–(7.11), the shared column sum in (7.8), the Cauchy bounds (7.12), the ordinary margins (7.13), and the telescoping product (7.14) prove positivity and assign each undirected edge one preliminary symmetric stress.

Set

$$\mu_i = \frac{1}{2} \sum_j \Gamma_{ij} \ell_{ij}, \quad W = \sum_i \mu_i, \quad w_i = \frac{\mu_i}{W}, \quad \gamma_{ij} = \frac{\Gamma_{ij}}{W}. \quad (7.20)$$

The tangent force equation and radial contraction give

$$\sum_j \Gamma_{ij} \Delta_{ij} = -2\mu_i \Omega_i.$$

Therefore $a_{ij} = \gamma_{ij}/w_i = \Gamma_{ij}/\mu_i$ proves (7.15). Let $R_i = \sum_j \Gamma_{ij} \ell_{ij}^2$. The loss-force and isotropy equations give, in the radial–tangent frame,

$$M_i = \frac{1}{\mu_i} \text{diag}(R_i, -R_i/2, -R_i/2) = \frac{3R_i}{2\mu_i} Z_i. \quad (7.21)$$

This proves $B_i = 0$ and (7.16) with $c_i = 3R_i/(2\mu_i)$. Since every active edge has angle at most $5h$,

$$c_i \leq 3\ell_{\max} \leq \frac{75}{2}h^2.$$

The row multiplier descends directly to the sampled quotient: if $S_2 A = 0$, then (7.16) gives $R_2 A = 0$. Thus no sampling-frame denominator occurs.

Every active edge has loss at least $h^2/(32\pi^2)$, so

$$r_i = \frac{2 \sum_j \Gamma_{ij}}{\sum_j \Gamma_{ij} \ell_{ij}} \leq 64\pi^2 h^{-2},$$

which proves (7.17). The lower half of (7.18) is Theorem 4.1: $\mathfrak{D}_2 \geq 6/r_{\max} \geq 6h^2/\mathbf{R}_3$. $\square$

Proposition 7.3 (structured support-preserving robustness)

At any level, preserve every ring count, longitude phase, radial mask, horizontal jump integer, the pole, the equator, and north–south reflection. Perturb each northern ring latitude by at most h^3/K_* , reflect the perturbation southward, and allow one common ambient rotation. Re-solving the same row systems produces a positive reversible generator with exact H_0, H_1 and

$$r_{\max} \leq \mathbf{R}_{\text{rob}} h^{-2}, \quad \mathfrak{D}_2 \leq C_{\text{rob}} h^2, \quad \mathbf{R}_{\text{rob}} = 256\pi^2, \quad C_{\text{rob}} = 54. \quad (7.22)$$

This proposition does not cover independent longitude perturbations or arbitrary node motion.

Proof The cancellation-free differentiated row programs have derivative bound K_* . The factor K_*^{-1} in the perturbation scale makes ordinary normalized rows change by $O(h^2)$ and transition rows by $O(h)$; there are $O(h^{-1})$ ordinary rows, while $Jh < 14/M_0$. The fixed margins (7.12)–(7.14), the separately guarded first row, and equatorial reflection therefore preserve the shared positive recurrence.

Moving either endpoint by at most h^3/K_* changes an active angular length by at most $2h^3/K_*$. Since $h < 1$ and $K_* > 32$, the perturbed active window is contained in $[h/16, 6h]$. Thus

$$\ell_{\min} \geq \frac{2}{\pi^2} \left(\frac{h}{16} \right)^2 = \frac{h^2}{128\pi^2}, \quad \ell_{\max} \leq \frac{(6h)^2}{2} = 18h^2.$$

Re-solving (7.10)–(7.11) and repeating (7.20)–(7.21) gives exact reproduction and the row multiplier. Hence $r_{\max} \leq 2/\ell_{\min} = 256\pi^2 h^{-2}$ and $\mathfrak{D}_2 \leq 3\ell_{\max} = 54h^2$. $\square$

8. Exact examples

8.1 Three extremal families in every dimension

All entries below are exact. Every family has uniform stationary masses, nonnegative reversible rates, exact coordinate eigenvalue $-(d-1)$, and $\mathfrak{D}_2 r_{\max} = d(d-1)$.

Family	Nodes and active graph	Directed active rate	$r_{\max}$	Common loss	$\mathfrak{D}_2$	rank S_2	dim K_X
Regular simplex	$d+1$ simplex vertices; complete graph	$(d-1)/(d+1)$	$d(d-1)/(d+1)$	$(d+1)/d$	$d+1$	d	$(d+1)(d-2)/2$
Cross- polytope	$\{\pm e_k\}_{k=1}^d$; all nonantipodal pairs	$1/2$	$d-1$	1	d	$d-1$	$d(d-1)/2$
Hypercube	$\{\pm 1/\sqrt{d}\}^d$; cube edges	$(d-1)/2$	$d(d-1)/2$	$2/d$	2	$\binom{d}{2}$	$d-1$

For each family, (5.3) holds with $c_* = d(d-1)/r_{\max}$. Consequently,

$$LS_2 = -d(2 - \ell_*)S_2,$$

where ℓ_* is the common active loss. These examples establish sharpness in every ambient dimension; no extrapolation from fixed-dimensional numerics is used.

8.2 Platonic examples in $d = 3$

With uniform masses, natural shortest-edge graphs, and their exact equality rates, the tetrahedron, octahedron, cube, icosahedron, and dodecahedron all satisfy

$$\mathfrak{D}_2 r_{\max} = 6, \quad E_{\text{sample}} = \{0\}.$$

The tetrahedral, octahedral, and cubical coefficient spaces have alias dimensions 2, 3, and 2, respectively. Thus nonzero trace-free quadratic forms may vanish on every node even at frontier equality. The icosahedral and dodecahedral quadratic sampling maps are injective.

8.3 What the examples do and do not show

The examples prove attainment and demonstrate that sampling aliases are compatible with equality. They do not classify all extremizers. Reversible blowups, covers, long-chord symmetric shells, and higher-dimensional equality families prevent such a classification without further hypotheses. Separately, nonregular weighted tangent tight frames show that even local equality does not force a regular vertex figure.

9. Rejected claims and deterministic counterexamples

The following statements are false or unsupported and are excluded from the paper.


Rejected claim	Obstruction retained in the proof package	Correct replacement
The Frobenius norm on $\text{Sym}_0(d)$ can replace the sampled quotient norm.	Nontrivial K_X in the simplex, cross-polytope, cube, and Platonic examples.	Work on $Q_X = V/K_X$ with metric (2.2).
Frontier equality reproduces H_2 exactly.	At equality $R_2 = c_* S_2$ with $c_* > 0$, while $E_{\text{sample}} = \{0\}$.	Equality minimizes the sampled residual under the positivity-rate constraint.
Trace saturation alone implies full frontier equality.	Unequal-rate and repeated-node exact fixtures separate the trace step from the rate and variance steps.	Require all three conditions in (5.1).
Local positive row stencils automatically assemble into a reversible generator.	Cycle incompatibility and detailed-balance failures.	Impose the global Markov, Gram, and Kolmogorov conditions of Theorem 5.3.
Every connected equality generator is Platonic.	Exact blowups, covers, long-chord icosahedral shells, and all-dimensional families.	Use Corollary 5.5 only under its explicit support and convexity hypotheses.
Small weighted defect implies uniform vertexwise closeness.	An order-one defect can be hidden at a vertex of arbitrarily small stationary mass.	Retain (6.9) or assume $w_{\min} > 0$.
Small global shell variance controls every active edge.	An edge of arbitrarily small directed conductance probability may carry order-one loss defect.	Assume $p_{ij} \geq \kappa$, retain the vertex-mass dependence, and use $w_{\min} > 0$ for a uniform all-edge statement.
Local edge control propagates globally without a graph parameter.	Long paths, cycles, and small-gap graphs accumulate variation.	Retain the $\kappa, w_{\min}$ margins and the applicable path, resistance, gap, or congestion dependence.
Small covariance defect alone yields stable eigenvectors or frame repair.	Isotropic targets, nearly singular tangent frames, antipodal collapse, and zero-loss bridges.	Add spectral-gap, tangent-frame, shell, overlap, and surjectivity margins as appropriate.
A fitted $O(h^2)$ slope proves the matching upper theorem.	Any finite prefix is compatible with different asymptotics and does not certify exactness or positivity at all levels.	Use the all-orders symbolic, Cauchy, recurrence, and positivity proof of Theorem 7.2.
Positivity universally forbids higher-order graph Laplacians.	Signed high-order graph formulas use cancellation outside the present class.	State only the scoped product theorem (4.3).

10. Consequences deliberately outside the flagship theorem

No generator-lift, semigroup comparison, transport-cost improvement, or physical-model performance theorem is claimed here. A signed construction may illustrate that leaving the positive cone changes the obstruction, but it does not strengthen (4.3) unless its normalization and comparison class are made identical. Such material should appear only in a later paper or a clearly separated discussion after independent proof and model audits.

Appendix A. Proof dependency graph

The all-dimensional lower frontier does not depend on the constructions. The matching corollaries use it only after the independent $d=2$ or $d=3$ upper family has verified positivity, exact coordinates, and the applicable rate cap. No construction is asserted for $d>3$.



The all-dimensional lower frontier does not depend on the explicit meshes.

Figure 1: Proof dependency graph. The lower frontier is independent of the explicit meshes; the matching result combines the frontier with the construction upper bounds.

Appendix B. Theorem-to-source-to-test-to-registry map

The proof source is controlling. A test source supplies falsification and regression evidence but is not a substitute for proof.

Manuscript result	Proof source	Test or audit source	Claim-registry entry	Status in this draft
Proposition 2.1, sampled quotient, Gram pencil, sampled exact-space dimension	afp_barrier_gate1/pure_math/covariance/P1A_QUADRATIC_FIDELITY_FOUNDATION.md, Sections 4–6	afp_barrier_gate1/pure_math/covariance/p1a_quadratic_fidelity_audit.py	P1A-QUOT; P1A-GRAM	accepted
Theorem 3.1, exact covariance and two-defect identities	same P1A source, Sections 2–3	same P1A audit	P1A-2DEF	accepted
Proposition 3.2, weighted adjoints and traces	same P1A source, Sections 4–6	same P1A audit	P1A-GRAM	accepted
Theorem 4.1, trace inequality and sharp product	afp_barrier_gate1/pure_math/covariance/P1B_SHARP_QUADRATIC_DEFECT_BOUND.md, Sections 3–13	afp_barrier_gate1/pure_math/covariance/p1b_sharp_quadratic_defect_audit.py	P1B-TRACE; P1B-SHARP	accepted
Theorem 5.1, equality characterization	same P1B source, Section 11; P1C source for geometric equivalences	P1B and P1C audits	P1B-EQUALITY; P1C-GLOBAL	accepted
Theorem 5.2, tangent block and tight-frame equivalence	afp_barrier_gate1/pure_math/covariance/P1C_EQUALITY_GEOMETRY.md	afp_barrier_gate1/pure_math/covariance/p1c_equality_geometry_audit.py	P1C-LOCAL	accepted
Theorem 5.3, global reversible assembly	same P1C source	same P1C audit	P1C-GLOBAL	accepted
Proposition 5.4 and Corollaries 5.5–5.6, antipodal, scoped classification, and global-moment consequences	same P1C source	same P1C audit	P1C-GLOBAL; P1C-CLASS; P1C-PLATONIC	accepted under stated hypotheses
Theorem 6.1, normalized stability budget	afp_barrier_gate1/pure_math/covariance/P1D_QUANTITATIVE_STABILITY.md, Sections 1–3	afp_barrier_gate1/pure_math/covariance/p1d_quantitative_stability_audit.py	P1D-MASTER	accepted
Corollary 6.2, exceptional mass and pointwise transfer	same P1D source, Sections 4–5	same P1D audit	P1D-VERTEX	accepted

Manuscript result	Proof source	Test or audit source	Claim-registry entry	Status in this draft
Corollary 6.3, conductance-shell and edgewise transfer	same P1D source, Section 6	same P1D audit	P1D-EDGE	accepted only with the named probability margin
Corollary 6.3, graphwise propagation	same P1D source, Section 7	same P1D audit	P1D-GRAPH	accepted only with the named mass and connectivity parameters
Corollary 6.3, covariance splitting	same P1D source, Sections 5 and 8	same P1D audit	P1D-COV	accepted; eigenvector transfer requires a spectral gap
Corollary 6.3, sampled-quotient scalarity	same P1D source, Section 8	same P1D audit	P1D-QUOTIENT	accepted only with α_X
Corollary 6.3, tangent-frame stability and repair	same P1D source, Sections 9–10	same P1D audit	P1D-FRAME	accepted only with the named frame and repair margins
Section 8, all-dimensional and Platonic exact families and aliases	P1C source	P1C audit	P1C-FAMILIES; P1C-PLATONIC; P1C-ALIAS	accepted
Section 9, unrestricted-classification rejection	P1C and P1D counterexample sections	P1C and P1D audits	P1C-UNRESTRICTED and caveats attached to P1D entries	rejected claims retained
Theorem 7.1, regular-polygon matching family	Section 7.1, direct Fourier proof	p1e_asymptotic_family_audit.py; QuadraticFidelityConstruction.lean	P1E-POLYGON	accepted for $d=2$
Theorem 7.2, adaptive-ring mesh, positivity and exact H_0, H_1	docs/publication_program/P1E_SHORT_GAP_S2_CONSTRUCTION.md, Sections 1–8	symbolic, Cauchy, polar, family, proof, referee and independent audits listed below	P1E-RING	accepted for $d=3$
Theorem 7.2, sampled quotient multiplier and matching constants	same P1E source, Section 9	p1e_short_gap_proof_audit.py; p1e_short_gap_cauchy_hostile_audit.py	P1E-ROW; F-CONSTRUCT	accepted for $d=3$
Proposition 7.3, structured robustness with $R_{\text{rob}} = 256\pi^2$, $C_{\text{rob}} = 54$	P1E source, Section 10, plus the active-window calculation in Proposition 7.3	p1e_no_guard_ring_independent_audit.py; final hostile guard audit	P1E-RING	accepted only for the fixed-support reflected class

The controlling registry is docs/publication_program/THEOREM_REGISTRY.md. Any change in theorem strength must first be reconciled with the proof source and then recorded in that registry; a passing test alone cannot promote a claim.

Appendix C. Numerical and falsification protocol

The computational appendices serve four limited purposes:

1. verify finite tensor identities and sign conventions on deterministic exact examples;
2. expose sampling aliases, leakage, and weighted-adjoint mistakes;
3. retain deterministic failure cases for omitted hypotheses; and
4. check explicit constants and extreme-scale conditioning with exact, algebraic, or outward-rounded interval arithmetic where appropriate.

The required audit programs are:

- `p1a_quadratic_fidelity_audit.py` for quotient, covariance, weighted Gram, and alias identities;
- `p1b_sharp_quadratic_defect_audit.py` for sharpness, degeneracy, sign, and normalization fixtures;
- `p1c_equality_geometry_audit.py` for tangent blocks, all-dimensional extremizers, Platonic exactness, aliases, antipodal branches, blowups, long-chord shells, weighted frames, and cycle compatibility; and
- `p1d_quantitative_stability_audit.py` for small masses, small active edge probabilities, long paths, small gaps, singular frames, sampling gaps, aliases, and certified extreme-scale inequalities;
- `p1e_short_gap_symbolic_matrix_audit.py`, `p1e_short_gap_cauchy_guard_audit.py`, and `p1e_short_gap_cauchy_hostile_audit.py` for the literal transition matrix and its uniform all-orders guard;
- `p1e_short_gap_polar_guard_audit.py`, `p1e_short_gap_proof_audit.py`, and `p1e_no_guard_ring_independent_audit.py` for the first row, exact constants, recurrence, reachable-domain margins, and perturbation arithmetic; and
- `p1e_short_gap_family_audit.py` and `p1e_short_gap_referee_audit.py` for finite generator reconstruction and deterministic failed mutations. The six alternative-route audits are retained separately and cannot be read as a proof by exhaustion.

No numerical rank threshold is used in a theorem. Exact symbolic minors or algebraic identities certify the ranks quoted in Section 8. Theorem 7.2 is all-orders; its finite generator output and convergence measurements are regressions only.

Appendix D. Formalization

The finite algebraic core is formalized in the following Lean sources:

Formal source	Mathematical scope
<code>AFPBarrier/QuadraticFidelityFoundation.lean</code>	weighted finite identities, sampling quotient foundations, covariance decomposition
<code>AFPBarrier/QuadraticFidelityLowerBound.lean</code>	trace comparison and sharp lower-bound algebra
<code>AFPBarrier/QuadraticEqualityGeometry.lean</code>	division-free tangent blocks and guarded normalized-frame identities
<code>AFPBarrier/QuadraticFidelityStability.lean</code>	normalized master-budget algebra and finite consequences
<code>AFPBarrier/QuadraticFidelityConstruction.lean</code>	shared-stress force and detailed-balance identities, moment normalization, four-point connectors, rate conversion, and regular-polygon residuals

The formalization is a second proof surface for the stated finite identities. It does not by itself formalize every geometric classification argument, external-priority statement, adaptive-ring schedule, or analytic Cauchy guard. The manuscript must not label an unformalized conclusion as machine-checked merely because a nearby algebraic lemma is formalized.

Appendix E. Reproducibility record

From the repository root, the deterministic theorem audits may be run with:

```
python afp_barrier_gate1/pure_math/covariance/p1a_quadratic_fidelity_audit.py
python afp_barrier_gate1/pure_math/covariance/p1b_sharp_quadratic_defect_audit.py
python afp_barrier_gate1/pure_math/covariance/p1c_equality_geometry_audit.py
python afp_barrier_gate1/pure_math/covariance/p1d_quantitative_stability_audit.py
python afp_barrier_gate1/pure_math/covariance/p1e_short_gap_symbolic_matrix_audit.py
python afp_barrier_gate1/pure_math/covariance/p1e_short_gap_cauchy_guard_audit.py
python afp_barrier_gate1/pure_math/covariance/p1e_short_gap_cauchy_hostile_audit.py
python afp_barrier_gate1/pure_math/covariance/p1e_short_gap_polar_guard_audit.py
python afp_barrier_gate1/pure_math/covariance/p1e_short_gap_proof_audit.py
python afp_barrier_gate1/pure_math/covariance/p1e_no_guard_ring_independent_audit.py
python afp_barrier_gate1/pure_math/covariance/p1e_short_gap_family_audit.py
python afp_barrier_gate1/pure_math/covariance/p1e_short_gap_referee_audit.py
```

The Lean surface may be checked from `afp_barrier_gate1` using the repository’s pinned toolchain and build command.

Release-specific workflow identifiers, continuous-integration run identifiers, commit hashes, environment captures, and artifact checksums belong only in a separate reproducibility manifest accompanying a tagged release. They are not part of any hypothesis or mathematical argument and must not appear in Sections 1–10.

The external-source bibliography for the priority audit is `docs/publication_program/p1f_manuscript/priority_sources.bib`. Every imported theorem must be accompanied by the source-specific hypothesis-transfer note in `PRIORITY_AND_HOSTILE_REFeree_AUDIT.md`; the present P1A–P1E proofs are self-contained and use those sources for context rather than as hidden proof lemmas.

Appendix F. Limitations and future work

1. The matching construction is proved only in ambient dimensions two and three. The sharp frontier, equality theory, and quantitative stability are all-dimensional, but no matching local family is claimed for $d > 3$.
2. The stability theorem is fundamentally weighted. Uniform pointwise rigidity requires a lower stationary mass; uniform edgewise rigidity requires a lower active conductance probability.
3. Graphwide rigidity depends on a quantitative connectivity parameter. Sampling-quotient scalarity depends on a lower sampling-frame bound, and tangent-frame repair depends on tangent conditioning and nondegeneracy margins.
4. Equality is not a classification by a short list of symmetric polytopes. The scoped three-dimensional convex classification has explicit hypotheses, while blowups, covers, long-chord shells, and higher-dimensional equality families remain; nonregular local frames separately obstruct local regularity.
5. Sampling aliases can occur for small or symmetric node sets. The theorem controls the genuinely sampled quotient and does not reconstruct coefficients in K_X .
6. No transport, physical-model, or application-level performance consequence follows from the harmonic operator norm alone. Such claims require separate models, normalizations, and audits.
7. Signed higher-accuracy graph operators are outside the positive reversible class. A future comparison should specify a common rate or cost constraint before drawing conclusions about the benefit of leaving the positive cone.
8. The $d=3$ perturbation proposition fixes ring counts, phases, masks, horizontal jumps, pole/equator data, and reflection. It does not cover arbitrary node motion or independent longitude perturbations.

The immediate construction question is now higher-dimensional: whether a different local compiler can produce the matching order for $d > 3$ with shared positive conductances and a quantitative global feasibility margin. A broader robustness result in $d=3$ would likewise require a uniform right inverse for the full shared six-moment edge system.

Appendix G. Related-theorem and hypothesis-transfer matrix

The table distinguishes external results from the theorems proved here. The long-form line-by-line convention audit, including primary-source theorem locations, is `PRIORITY_AND_HOSTILE_REFeree_AUDIT.md`.

Source and variables	External hypotheses	External conclusion used	Overlap	New distinction and transfer guard
Babecki–Thomas: graph eigenspaces, node subsets, quadrature weights	connected regular unweighted graph; normalized-adjacency ordering	graphical designs correspond through Gale duality to eigenpolytope faces, with support consequences	finite eigenspaces and positive weights	their weights average a fixed graph; w_i and γ_{ij} here define the operator. No spherical eigenmap, rate cap, or quadratic frontier transfers
Babecki–Shiroma: positive edge-weighted graphs and eigenpolytopes	connected graph; symmetric combinatorial Laplacian in uniform Euclidean metric	broad eigenspace/eigenpolytope universality for positive weighted graphs	prescribed eigenspaces with positive edges	universality need not preserve unit spherical geometry, locality, degree, nonuniform mass, target eigenvalue, or H_2 residual; flip the positive-semidefinite sign and retain the mass matrix
Steinerberger (Steinerberger 2021): manifold eigenfunctions, n quadrature nodes	compact boundaryless manifold; nonnegative quadrature exact on an initial spectral segment	asymptotic cardinality limitation for positive quadrature	positivity and continuous harmonic modes	integration is not a graph-operator identity. Keep manifold dimension $d - 1$, multiplicity, volume measure, and the asymptotic $o(n)$ term; no finite frontier follows
Izmestiev–Lam: geodesic spherical triangulation, c_{ij}, d_i	nondegenerate closed triangulation; local Delaunay for $c_{ij} \geq 0$	their normalized spherical Laplacian is reversible and satisfies $\Delta_s p = -2p$	local positive reversible coordinate-exact operators on $\mathbb{S}^2$	this paper does not claim to introduce that object. Their theorem supplies neither the sampled quotient/frontier nor the adaptive-ring H_2 and mesh constants; conversion requires $w_i = d_i / \sum d$, $\gamma_{ij} = c_{ij} / \sum d$
Seibold: Euclidean point cloud and row stencil coefficients	polynomial moment equations; local geometric cone conditions	sparse positive Poisson stencils and feasibility criteria	positive local moment cones	a rowwise stencil is generally nonsymmetric and gives no shared γ_{ij} or stationary masses. Curvature, global reconciliation, and a strict margin must be proved separately

Source and variables	External hypotheses	External conclusion used	Overlap	New distinction and transfer guard
Babecki–Steinerberger–Thomas: base graph and first k eigenpairs	connected positive weighted graph; candidate is a reweighted spanning subgraph	eigenpair-preserving Laplacians form a polyhedral/spectrahedral slice	positive eigenpair preservation	here the geometric coordinate module is prescribed, aliases are quotiented, and the next continuous shell is measured under $r_{\max}$. With nonuniform w , the comparison is a generalized eigenproblem
García Trillos–Gerlach–Hein–Slepčev: random manifold sample and kernel bandwidth	i.i.d. density bounds, compact manifold geometry, kernel and bandwidth conditions	high-probability eigenvalue/eigenvector convergence to a weighted Laplace–Beltrami operator	local positive graph Laplacians approaching a continuum operator	probabilistic spectral convergence does not give finite exact H_1 , bounded degree, shared correction, or the deterministic $O(h^2)$ quotient bound. Their bandwidth is not silently identified with this fill parameter
Martin–Tanaka: association scheme adjacency algebra	finite commutative scheme; uniform counting metric	primitive idempotents and nonnegative Krein parameters organize products and embeddings	symmetric examples and harmonic-module products	one must first prove that L lies in the Bose–Mesner algebra and identify its idempotents with sampled spherical harmonics. Unequal masses and aliases are not automatic
Bannai–Bannai: finite spherical sets and polynomial moments	usually uniform spherical t -design measure	harmonic-moment characterization and design bounds	common symmetric node sets and frame moments	design exactness is a measure identity, not a generator. It implies neither reversible local edges, sampled injectivity, nor L -invariance; frontier equality has positive residual, not exact H_2
Ahrens–Beylkin (Ahrens and Beylkin 2009): icosahedral orbits and quadrature weights	rotational invariance; moment equations for targeted polynomial spaces	rotationally invariant spherical quadratures	symmetry-orbit reductions	quadrature weights are not conductances or stationary masses. Symmetry preserves a generator constraint only after support and orientation invariance are proved

Source and variables	External hypotheses	External conclusion used	Overlap	New distinction and transfer guard
Yoon: graph powers/walks and m -Laplacian coefficients	simple graph; signed higher-order finite-difference-inspired operator	high formal order on cycles using alternating coefficients for $m \geq 2$	graph-Laplacian accuracy	the signed class lies outside $\gamma \geq 0$. The present theorem is only the sampled degree-two residual/rate frontier for positive reversible coordinate-exact generators
Benedetto–Fickus (Benedetto and Fickus 2003): unit vectors and frame operator	finite unit-norm frames in a Hilbert space	unit-norm tight frames minimize the frame potential	tangent tight-frame equality language	tight frames themselves are established theory. The new statement is their forced appearance together with common loss/rate and global detailed-balance compatibility in equality for this frontier

No row is imported as a proof of Theorems 3.1, 4.1, 5.1–5.3, 6.1, or 7.1–7.2. External results supply context and established terminology; every normalization used in the new theorem hierarchy is derived internally.

References

- Bibliographic data for the sources cited in the priority discussion are in `priority_sources.bib`. Journal versions are used when available, and the related-work discussion retains the convention-transfer notes needed to keep the comparison classes distinct.
- Ahrens, Cory, and Gregory Beylkin. 2009. “Rotationally Invariant Quadratures for the Sphere.” *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* 465 (2110): 3103–25. <https://doi.org/10.1098/rspa.2009.0104>.
- Babecki, Catherine, and David Shiroma. 2023. “Eigenpolytope Universality and Graphical Designs.” <https://arxiv.org/abs/2209.06349>.
- Babecki, Catherine, Stefan Steinerberger, and Rekha R. Thomas. 2023. “Spectrahedral Geometry of Graph Sparsifiers.” <https://arxiv.org/abs/2306.06204>.
- Babecki, Catherine, and Rekha R. Thomas. 2022. “Graphical Designs and Gale Duality.” <https://arxiv.org/abs/2204.01873>.
- Bannai, Eiichi, and Etsuko Bannai. 2009. “A Survey on Spherical Designs and Algebraic Combinatorics on Spheres.” *European Journal of Combinatorics* 30 (6): 1392–1425. <https://doi.org/10.1016/j.ejc.2008.11.007>.
- Benedetto, John J., and Matthew Fickus. 2003. “Finite Normalized Tight Frames.” *Advances in Computational Mathematics* 18 (2–4): 357–85. <https://doi.org/10.1023/A:1021323312367>.
- García Trillos, Nicolás, Moritz Gerlach, Matthias Hein, and Dejan Slepčev. 2020. “Error Estimates for Spectral Convergence of the Graph Laplacian on Random Geometric Graphs Towards the Laplace–Beltrami Operator.” *Foundations of Computational Mathematics* 20: 827–87. <https://arxiv.org/abs/1801.10108>.
- Izmestiev, Ivan, and Wai Yeung Lam. 2025. “Discrete Laplacians—Spherical and Hyperbolic.” *Journal of the London Mathematical Society*. <https://doi.org/10.1112/jlms.70235>.
- Martin, William J., and Hajime Tanaka. 2008. “Commutative Association Schemes.” <https://arxiv.org/abs/0811.2475>.
- Seibold, Benjamin. 2008. “Minimal Positive Stencils in Meshfree Finite Difference Methods for the Poisson Equation.” *Computer Methods in Applied Mechanics and Engineering* 198 (3–4): 592–601. <https://doi.org/10.1016/j.cma.2008.09.001>.
- Steinerberger, Stefan. 2021. “Spectral Limitations of Quadrature Rules and Generalized Spherical Designs.” *International Mathematics Research Notices*, no. 16: 12265–80. <https://arxiv.org/abs/1708.08736>.
- Yoon, Mary. 2025. “Graph Laplacians with Higher Accuracy.” <https://arxiv.org/abs/2504.04461>.