



+974 50124683



dentalreception@dohahealthcare.com



P.O. Box: 23632,
Doha – Qatar

مجموعة الدوحة للرعاية الصحية
قسم طب الأسنان



مركز المستقبل الطبي

Future
Medical Center

Pre-Operative Investigations for Dental Implant Surgery



To ensure your safety and the success of your dental implant procedure, please complete the following investigations **before** your scheduled surgery:



- Complete Blood Count (CBC)



- HbA1c (Glycated Hemoglobin)



- International Normalized Ratio (INR)
(if indicated or if taking blood-thinning medication)



- CBCT Scan (Cone Beam Computed Tomography)



Note:

- ✓ The CBCT scan will be reviewed by the implant center **before** proceeding with treatment.
- ✓ Additional investigations may be requested depending on your medical history and clinical findings.



+974 50124683



dentalreception@dohahealthcare.com



P.O. Box: 23632,
Doha – Qatar



+974 50124683



dentalreception@dohahealthcare.com



P.O. Box: 23632,
Doha – Qatar

مجموعة الدوحة للرعاية الصحية

قسم طب الأسنان



مركز المستقبل الطبي

Future
Medical Center

Dental Implant Pre-Operative Instructions



Please read and follow these instructions **before** your dental implant appointment:



- **Maintain excellent oral hygiene** by brushing at least twice daily and flossing regularly before your implant procedure.



- **Complete professional dental scaling and polishing at least one week before your implant appointment.**

Your implant procedure may be postponed if oral hygiene is inadequate or if active gingival inflammation or periodontal disease is present.



- **Complete all required pre-operative investigations** before your implant appointment. Please refer to the Pre-Operative Investigations sheet provided by your dentist.



- **Take your prescribed antibiotic** exactly as instructed by your dentist.



- **Take your prescribed anti-inflammatory medication one hour before** your appointment, unless otherwise advised by your dentist.



- **Have a light meal before your appointment**, unless you have been instructed otherwise.



- **Inform your dentist immediately** if you develop fever, cold symptoms, or any change in your medical condition before your scheduled implant surgery.



- **Continue taking your regular medications** unless your dentist or physician has advised you otherwise.



Important Reminder

Please arrive
10–15 minutes
before your
scheduled
appointment and
bring:



Your identification
(QID/Passport
if required)



Your recent
investigation
results



Any current
medications or
updated medical
reports



+974 50124683



dentalreception@dohahealthcare.com



P.O. Box: 23632,
Doha – Qatar



+974 50124683



dentalreception@dohahealthcare.com



P.O. Box: 23632,
Doha – Qatar

مجموعة الدوحة للرعاية الصحية

قسم طب الأسنان



مركز المستقبل الطبي

Future
Medical Center

Dental Implant Post-Operative Instructions



Please read and follow these instructions **after** your dental implant surgery:



- **Continue taking your prescribed antibiotic** exactly as directed until the full course is completed.



- **Continue taking your prescribed anti-inflammatory/pain medication as instructed** by your dentist to help control pain and swelling.



- **Do not touch, pull, or play with the sutures (stitches).**
The sutures will be removed by your dentist at your follow-up appointment unless they are dissolvable.



- **Avoid chewing directly on the implant site** until your dentist advises that it is safe to do so.



- **If a temporary crown has been placed:**
 - Avoid biting hard foods with the temporary crown.
 - Do not chew sticky or very hard foods on the implant side.
 - Contact the clinic immediately if the temporary crown becomes loose or falls off.



- **Maintain good oral hygiene.**

Brush the other teeth normally.
Clean around the implant area gently as instructed by your dentist. If a chlorhexidine mouthwash has been prescribed, use it as directed.



- **Attend all scheduled follow-up appointments.**

These visits are important to monitor healing, remove sutures if required, and ensure successful implant integration.



Contact the Clinic Immediately If You Experience:



- Excessive bleeding that does not stop with pressure.



- Increasing swelling after 2–3 days.



- Severe pain that is not relieved by medication.



- Fever or signs of infection.



- A loose implant or temporary crown.



+974 50124683



dentalreception@dohahealthcare.com



P.O. Box: 23632,
Doha – Qatar