



Cambridge International AS & A Level

CANDIDATE
NAME

Fuzail									
--------	--	--	--	--	--	--	--	--	--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3

May/June 2020

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Blank pages are indicated.



2020 remaining 2020. turn over

sample = turn

2

- 1 Find the quotient and remainder when $6x^4 + x^3 - x^2 + 5x - 6$ is divided by $2x^2 - x + 1$. [3]

$$\begin{array}{r} 3x^2 + 2x - 1 \\ 2x^2 - x + 1 \overline{) 6x^4 + x^3 - x^2 + 5x - 6} \end{array}$$

$$- 6x^4 - 3x^3 + 3x^2$$

$$4x^3 - 4x^2 + 5x$$

$$4x^3 - 2x^2 + 2x$$


$$-2x^2 + 3x - 6$$

$$-2x^2 + x - 1$$

$$0 \quad 2x - 5$$

$$\text{quotient} = 3x^2 + 2x - 1$$

$$\text{remainder} = 2x - 5$$



The variables x and y satisfy the equation $y^2 = Ae^{kx}$, where A and k are constants. The graph of $\ln y$ against x is a straight line passing through the points $(1.5, 1.2)$ and $(5.24, 2.7)$ as shown in the diagram.

Find the values of A and k correct to 2 decimal places.

[5]

$$2 \ln y = \ln A + kx$$

$$\ln y = \frac{\ln A}{2} + \frac{kx}{2}$$

$$y \text{ intercept} = \frac{\ln A}{2}, \quad \frac{k}{2} = \text{gradient}$$

$$y = mx + c$$

$$m = \frac{2.7 - 1.2}{5.24 - 1.5} = 0.4012$$

$$\frac{k}{2} = 0.4012$$

$$k = 0.80$$

$$\begin{aligned} c &= y - mx \\ &= 1.2 - 0.4012(1.5) \\ &= 0.5982 \end{aligned}$$

$$\begin{aligned} \therefore \ln A &= 1.1964 \\ A &= e^{1.1964} \\ A &= 3.31 \end{aligned}$$

3 Find the exact value of

$$\int_1^4 x^{\frac{3}{2}} \ln x \, dx.$$

LIATE

[5]

$$u = \ln x \quad u' = \frac{1}{x}$$

$$v = \frac{2x^{2.5}}{5} \quad v' = x^{1.5}$$

$$\ln x \left(\frac{2x^{2.5}}{5} \right) - \frac{2}{5} \int x^{2.5} \left(\frac{1}{x} \right) dx$$

$$\frac{2}{5} x^{2.5} \ln x - \frac{2}{5} \int x^{1.5} dx$$

$$\frac{2}{5} x^{2.5} \ln x - \frac{2}{5} \left(\frac{2}{5} x^{2.5} \right)$$

$$\frac{2}{5} x^{2.5} \ln x - \frac{4x^{2.5}}{25}$$

$$\frac{2x^{2.5}}{5} \left(\ln x - \frac{2}{5} \right)$$

$$\left[\frac{2(4)^{2.5}}{5} \left(\ln 4 - \frac{2}{5} \right) \right] - \left[\frac{2(1)^{2.5}}{5} \left(-\frac{2}{5} \right) \right]$$

$$\frac{64 \ln 4}{5} - \frac{128}{25} + \frac{4}{25}$$

$$\frac{64 \ln 4}{5} - \frac{124}{25}$$

- 4 A curve has equation $y = \cos x \sin 2x$.

Find the x -coordinate of the stationary point in the interval $0 < x < \frac{1}{2}\pi$, giving your answer correct to 3 significant figures. [6]

$$y = \cos x \sin 2x$$

$$\frac{dy}{dx} = -\sin x \sin 2x + 2 \cos 2x \cos x$$

at stationary point $\frac{dy}{dx} = 0$

$$\therefore 2 \cos 2x \cos x = \sin x \sin 2x$$

$$2(2\cos^2 x - 1)\cos x = \sin x \cdot 2\sin x \cos x$$

$$\frac{4\cos^3 x - 2\cos x}{\cancel{\cos x}} = 2\sin^2 x$$

$$4\cos^2 x - 2 = 2 - 2\cos^2 x$$

$$6\cos^2 x - 4 = 0$$

$$\cos x = \pm \sqrt{\frac{4}{6}}$$

$$x = 0.615$$

- 5 (a) Express $\sqrt{2} \cos x - \sqrt{5} \sin x$ in the form $R \cos(x + \alpha)$, where $R > 0$ and $0^\circ < \alpha < 90^\circ$. Give the exact value of R and the value of α correct to 3 decimal places. [3]

$$R = \sqrt{(\sqrt{2})^2 + (\sqrt{5})^2}$$

$$\alpha = \tan^{-1}\left(\frac{\sqrt{5}}{\sqrt{2}}\right)$$

$$= \sqrt{2 + 5}$$

$$= \sqrt{7}$$

$$= 57.688$$

$$\therefore \sqrt{7} \cos(x + 57.688)$$

- (b) Hence solve the equation $\sqrt{2} \cos 2\theta - \sqrt{5} \sin 2\theta = 1$, for $0^\circ < \theta < 180^\circ$. [4]

$$\sqrt{7} \cos(2x + 57.688) = 1$$

$$2x + 57.688 = \cos^{-1}\left(\frac{1}{\sqrt{7}}\right)$$

$$2x + 57.688 = 67.792, 292.208$$


$$2x + 57.688 = 67.792$$

$$\underline{\underline{x = 5.1}}$$

$$2x + 57.688 = 292.208$$

$$\underline{\underline{x = 117.3}}$$

6



The diagram shows the curve $y = \frac{x}{1 + 3x^4}$, for $x \geq 0$, and its maximum point M .

- (a) Find the x -coordinate of M , giving your answer correct to 3 decimal places.

[4]

$$= \frac{(1+3x^4)(1) - (x)(12x^3)}{(1+3x^4)^2}$$

$$= \frac{1+3x^4-12x^4}{(1+3x^4)^2}$$

$$\frac{1-9x^4}{(1+3x^4)^2} = 0$$

$$1-9x^4 = 0$$

$$x^4 = \frac{1}{9}$$

$$x = 0.577$$

- (b) Using the substitution $u = \sqrt{3}x^2$, find by integration the exact area of the shaded region bounded by the curve, the x -axis and the line $x = 1$. [5]

$$y = \frac{x}{1+3x^2}$$

$$u = \sqrt{3}x^2$$

$$du = 2\sqrt{3}x \, dx$$

$$\int_0^1 \frac{x}{1+u^2} \times \frac{du}{2\sqrt{3}x}$$

$$\text{when } x=0 \quad u=0$$

$$x=1 \quad u=\sqrt{3}$$

$$\frac{1}{2\sqrt{3}} \times \int_0^1 \frac{1}{u^2+1} du$$

$$\frac{1}{2\sqrt{3}} \times \tan^{-1}\left(\frac{u}{1}\right)$$

$$\left[\frac{\tan^{-1}\left(\frac{u}{1}\right)}{2\sqrt{3}} \right]_0^{\sqrt{3}}$$

$$\left(\frac{1}{2\sqrt{3}} \times \frac{1}{3} \right) - (0)$$

$$\frac{\pi}{6\sqrt{3}}$$

- 7 The variables x and y satisfy the differential equation

$$\frac{dy}{dx} = \frac{y-1}{(x+1)(x+3)}.$$

It is given that $y = 2$ when $x = 0$.

Solve the differential equation, obtaining an expression for y in terms of x .

[9]

$$\int \frac{1}{y-1} dy = \int \frac{1}{(x+1)(x+3)} dx$$

$$\ln(y-1) = \frac{1}{2} \left[\int \frac{1}{x+1} dx - \int \frac{1}{x+3} dx \right] = \frac{A}{x+1} + \frac{B}{x+3}$$

$$\ln(y-1) = \frac{1}{2} [\ln(x+1) - \ln(x+3)] + C$$

$$\ln(y-1) = \frac{1}{2} \ln \left(\frac{x+1}{x+3} \right) + C$$

$$C = \ln(1) - \frac{1}{2} \ln \left(\frac{1}{3} \right)$$

$$C = -\frac{1}{2} \ln \frac{1}{3}$$

$$\therefore y-1 = e^{\frac{1}{2} \ln \left(\frac{x+1}{x+3} \right) - \frac{1}{2} \ln \left(\frac{1}{3} \right)} + 1$$

$$y = e^{\frac{1}{2} \ln \left(\frac{3x+3}{x+3} \right)} + 1$$

$$y = \sqrt{\frac{3x+3}{x+3}} + 1$$

- 8 (a) Solve the equation $(1 + 2i)w + iw^* = 3 + 5i$. Give your answer in the form $x + iy$, where x and y are real. [4]

$$\text{let } w = x + iy$$

$$(1 + 2i)(x + iy) + i(x - iy) = 3 + 5i$$

$$x + iy + 2xi + 2y(-1) + xi - (-1)y = 3 + 5i$$

$$x + iy + 3xi - 2y + y = 3 + 5i$$

$$3 = x - y$$

$$5 = y + 3x \quad (2)$$

$$\therefore x = 3 + y \quad (1)$$

sub (1) in (2)

$$5 = y + 3(3 + y)$$

$$5 = y + 9 + 3y$$

$$4y = -4$$


$$y = -1$$

$$\therefore x = 3 + (-1)$$

$$x = 2$$

$$\underline{\underline{z = 2 - i}}$$

- (b) (i) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 2 - 2i| \leq 1$ and $\arg(z - 4i) \geq -\frac{1}{4}\pi$. [4]



- (ii) Find the least value of $\text{Im } z$ for points in this region, giving your answer in an exact form. [2]

$$(x-2)^2 + (y-2)^2 = 1^2 \quad y = -x + 4$$

$$x^2 - 4x + 4 + (-x + 2)^2 = 1$$


$$x^2 - 4x + 4 + x^2 - 4x + 4 = 1$$

$$2x^2 - 8x + 7 =$$

$$x = 2 + \frac{1}{\sqrt{2}}$$

$$\therefore y = 2 - \frac{1}{\sqrt{2}} \quad \checkmark$$

9



The diagram shows the curves $y = \cos x$ and $y = \frac{k}{1+x}$, where k is a constant, for $0 \leq x \leq \frac{1}{2}\pi$. The curves touch at the point where $x = p$.

- (a) Show that p satisfies the equation $\tan p = \frac{1}{1+p}$. [5]

$$\cos p = \frac{k}{1+p}$$

$$-\sin p = \frac{(1+p)(0) - k(1)}{(1+p)^2}$$

$$-\sin p = \frac{-k}{(1+p)^2}$$

$$k = (\sin p)(1+p)^2 \quad \text{--- ①}$$

$$k = (\cos p)(1+p) \quad \text{②}$$

$$\begin{array}{l} \text{①} : k = (\sin p)(1+p)^2 \\ \text{②} : k = (\cos p)(1+p) \\ \hline 1 = \tan p (1+p) \end{array}$$

$$\tan p = \frac{1}{1+p}$$

proved

- (b) Use the iterative formula $p_{n+1} = \tan^{-1}\left(\frac{1}{1+p_n}\right)$ to determine the value of p correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

$$p_{n+1} = \tan^{-1}\left(\frac{1}{1+p_n}\right)$$

$$p_1 = \frac{1}{4}\pi$$

$$p_2 = 0.51056$$

$$p_3 = 0.58978$$

$$p_4 = 0.56291$$

$$p_5 = 0.56920$$

$$p_6 = 0.56737$$

$$p_7 = 0.56790$$

$$p_8 = 0.56774$$

$$p_9 = 0.56779$$

$$\therefore p = 0.568$$

- (c) Hence find the value of k correct to 2 decimal places. [2]


$$k = 1 + 0.568 \times (1 + 0.568) \\ = 1.32$$

- 10 With respect to the origin O , the points A and B have position vectors given by $\vec{OA} = 6\mathbf{i} + 2\mathbf{j}$ and $\vec{OB} = 2\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$. The midpoint of OA is M . The point N lying on AB , between A and B , is such that $AN = 2NB$.

(a) Find a vector equation for the line through M and N .

[5]

$$\vec{OM} = \frac{1}{2}\vec{OA} = \frac{1}{2}\begin{pmatrix} 6 \\ 2 \\ 0 \end{pmatrix} = \begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix}$$



$$\vec{MN} = \vec{MA} + \vec{AN}$$

$$= \begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix} + \frac{2}{3}\vec{AB}$$

$$= \begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix} + \frac{2}{3}\begin{pmatrix} -4 \\ 0 \\ 3 \end{pmatrix}$$

$$\vec{AB} = \vec{OB} - \vec{OA}$$

$$= \begin{pmatrix} 2 \\ 2 \\ 3 \end{pmatrix} - \begin{pmatrix} 6 \\ 2 \\ 0 \end{pmatrix} = \begin{pmatrix} -4 \\ 0 \\ 3 \end{pmatrix}$$

$$\vec{MN} = \begin{pmatrix} 1/3 \\ 1 \\ 2 \end{pmatrix}$$

$$\mathbf{r} = \begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 1/3 \\ 1 \\ 2 \end{pmatrix}$$

$$\mathbf{r} = 3\mathbf{i} + \mathbf{j} + \lambda\left(\frac{1}{3}\mathbf{i} + \mathbf{j} + 2\mathbf{k}\right)$$

The line through M and N intersects the line through O and B at the point P .

(b) Find the position vector of P .

[3]

$$MN: \begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 1/3 \\ 1 \\ 2 \end{pmatrix}$$

$$OB: \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + \mu \begin{pmatrix} 2 \\ 2 \\ 3 \end{pmatrix}$$

$$\textcircled{1} \quad 3 + \frac{1}{3}\lambda = 2\mu$$

$$\textcircled{2} \quad 1 + \lambda = 2\mu$$

$$\hookrightarrow \therefore \lambda = 2\mu - 1 \quad \textcircled{3}$$

sub $\textcircled{3}$ into $\textcircled{1}$

$$3 + \frac{1}{3}(2\mu - 1) = 2\mu$$

$$3 = \frac{2}{3}\mu - \frac{1}{3} = 2\mu$$

$$\frac{8}{3} = \frac{4}{3}\mu$$

$$\mu = 2$$

$$\therefore P = 2 \begin{pmatrix} 2 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \\ 6 \end{pmatrix}$$

$$P = 4i + 4j + 6k$$

(c) Calculate angle OPM , giving your answer in degrees.

[3]

$$PO = \begin{pmatrix} -4 \\ -4 \\ -6 \end{pmatrix}$$

$$PM = 3 \begin{pmatrix} -1/3 \\ -1/3 \\ 1 \end{pmatrix} = \begin{pmatrix} -1 \\ -3 \\ 3 \end{pmatrix}$$

$$\lambda = \frac{4-1}{3}$$

$$\therefore \cos \theta = \frac{4 + 12 + 36}{\sqrt{4^2 + 4^2 + 6^2} \cdot \sqrt{1^2 + 3^2 + 6^2}}$$

$$\cos \theta = \frac{52}{2\sqrt{78}}$$

$$\cos \theta = 0.92976$$

$$\theta = 21.6$$

This image shows a full page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page, typical of notebook or legal stationery. There are no margins, text, or other markings on the page.

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.