

**[Turn over**

1 Solve the inequality  $|2x - 1| > 3|x + 2|$ .

[4]

① ✗

$$(2x-1)^2 > [3(x+2)]^2$$

$$4x^2 - 4x + 1 > 9(x^2 + 4x + 4)$$


$$4x^2 - 4x + 1 > 9x^2 + 36x + 36$$

$$5x^2 + 40x + 35 < 0$$

$$x_{1,2} = \frac{-40 \pm \sqrt{1600 - 700}}{10} = \frac{-40 \pm \sqrt{900}}{10} = \frac{-40 \pm 30}{10}$$

$$x < -1$$

$$x > -\frac{3}{5}$$



checking  $x < -1$

$$|2(-2) - 1| > 3|-1 + 2|$$

$$5 > 3$$

✓

checking  $x > -\frac{3}{5}$  ✗

$$|2(0) - 1| > 3|0 + 2|$$

$$1 \not> 6$$

$$\therefore x < -1$$

- 2 Find the exact value of  $\int_0^1 (2-x)e^{-2x} dx$ .

3 [5]

L I A I E

$$\int_0^1 (2-x) e^{-2x}$$

$$u = 2-x \quad u' = -1$$

$$v = -\frac{1}{2}e^{-2x} \quad v' = e^{-2x}$$

$$(2-x) \left(-\frac{1}{2}e^{-2x}\right) - \int \left(-\frac{1}{2}e^{-2x}\right)(-1) dx$$

$$\frac{x}{2}e^{-2x} - e^{-2x} - \frac{1}{2} \int e^{-2x} dx$$

$$\frac{x}{2}e^{-2x} - e^{-2x} - \frac{1}{2} \left(-\frac{1}{2}\right) e^{-2x}$$

$$\frac{x}{2}e^{-2x} - e^{-2x} + \frac{1}{4}e^{-2x}$$

$$\left[ \frac{e^{-2x}}{4} (2x - 3) \right]_0^1$$

$$\left( \frac{e^{-2}}{4} \cdot 2 - \frac{3e^{-2}}{4} \right) - \left( \frac{1}{4} \cdot 0 - \frac{3}{4} \right)$$

$$\frac{e^{-2}}{2} + \frac{3}{4} - \left( -\frac{e^{-2}}{4} + \frac{3}{4} \right)$$

$$\frac{1}{8} (e^{-2} + 3) - \frac{1}{4} (3 - e^2)$$

- 3 (a) Show that the equation

$$\ln(1 + e^{-x}) + 2x = 0$$

\*

can be expressed as a quadratic equation in  $e^x$ . ?

[2]

$$\ln(1 + e^{-x}) = -2x$$

$$1 + e^{-x} = e^{-2x}$$

$$\frac{1}{e^{2x}} = 1 + e^{-x}$$

$$e^{2x} + e^{2x}(e^{-x}) = 1$$

$$e^{2x} + e^x - 1 = 0 \quad \checkmark$$

- (b) Hence solve the equation  $\ln(1 + e^{-x}) + 2x = 0$ , giving your answer correct to 3 decimal places.

[4]

$$\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = e^x$$

(2)

$$x^2 + x - 1$$

$$a = 1, b = 1, c = -1$$

$$\frac{-1 \pm \sqrt{1 - 4(1)(-1)}}{2}$$

$$\frac{-1 \pm \sqrt{1 + 4}}{2}$$

$$\frac{-1 + \sqrt{5}}{2}, \frac{-1 - \sqrt{5}}{2}$$

$$e^x = -1.618, e^x = 0.618$$

$$x = \ln 0.618 = -0.481$$

4 The equation of a curve is  $y = x \tan^{-1}(\frac{1}{2}x)$ .

(a) Find  $\frac{dy}{dx}$ .

[3]

$$\frac{dy}{dx} = \tan^{-1}(\frac{1}{2}x)(1) + x \left( \frac{1}{1+(\frac{x}{2})^2} \right)$$

$$= \tan^{-1}(\frac{1}{2}x) + \left( \frac{x}{1+\frac{x^2}{4}} \right)$$

$$= \tan^{-1}(\frac{1}{2}x) + \left( x \times \frac{4}{4+x^2} \right)$$

$$= \tan^{-1}(\frac{1}{2}x) + \frac{4x}{x^2+4} \times \frac{1}{2}$$

$$\tan^{-1}(\frac{1}{2}x) + \frac{4x}{2x^2+8}$$


$$+ \frac{2x}{x^2+4} = \frac{2x}{x^2+4}$$

(b) The tangent to the curve at the point where  $x = 2$  meets the  $y$ -axis at the point with coordinates  $(0, p)$ .

\*

Find  $p$ .

[3]



$$m = \tan^{-1}(1) + \frac{4}{6} = 1.452$$

$$y = 1.452x + c$$

$$y = c = p$$

$$p = 1.452(0) + p$$

$$0 = 0$$

5 By first expressing the equation

$$\tan \theta \tan(\theta + 45^\circ) = 2 \cot 2\theta$$

as a quadratic equation in  $\tan \theta$ , solve the equation for  $0^\circ < \theta < 90^\circ$ .

[6]

$$\tan \theta \left( \frac{\tan \theta + 1}{1 - \tan \theta} \right) = 2 \times \frac{1}{\tan 2\theta}$$

$$\frac{\tan^2 \theta + \tan \theta}{1 - \tan \theta} = 2 \times \left( \frac{1 - \tan^2 \theta}{2 \tan \theta} \right)$$

$$\frac{\tan^2 \theta + \tan \theta}{1 - \tan \theta} = \frac{1 - \tan^2 \theta}{\tan \theta}$$

$$\tan^3 \theta + \tan^2 \theta = (1 - \tan^2 \theta)(1 - \tan \theta)$$

$$\cancel{\tan^3 \theta} + \tan^2 \theta = 1 - \tan^2 \theta - \tan \theta + \cancel{\tan^3 \theta}$$

$$2 \tan^2 \theta + \tan \theta - 1 = 0$$

$$\frac{-1 \pm \sqrt{1 - 4(-1)(2)}}{4}$$

$$\frac{-1 \pm \sqrt{9}}{4}$$

$$\frac{-1 \pm 3}{4} = \frac{1}{2}, -1$$

$$\theta_1 = \tan^{-1}\left(\frac{1}{2}\right) = 26.6$$

$$\theta_2 = \tan^{-1}(1) = 45, 180 - 45 = 135 \times$$

$$\theta = 26.6$$


$$\begin{array}{c|ccc} x & 1 & 2 & 3 \\ \hline y & 1 & 32 & 243 \end{array}$$

$$y = 2 + a$$

$$\begin{array}{c|ccc} x & 1 & 2 & 3 \\ \hline y & 3 & 4 & 5 \end{array}$$

- 6 (a) By sketching a suitable pair of graphs, show that the equation  $x^5 = 2 + x$  has exactly one real root. [2]

$$y = x^5$$



$$x^5 - x - 2 = 0$$

Intersect at 1 point only, between 1 and 2

- (b) Show that if a sequence of values given by the iterative formula

$$x_{n+1} = \frac{4x_n^5 + 2}{5x_n^4 - 1}$$

converges, then it converges to the root of the equation in part (a).

[2]



- (c) Use the iterative formula with initial value  $x_1 = 1.5$  to calculate the root correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

$$x_1 = \frac{4(1.5)^5 + 2}{5(1.5)^4 - 1} = 1.33162$$

$$x_2 = 1.27352$$

$$x_3 = 1.26724$$

$$x_4 = 1.26717$$

$$x_5 = 1.26717$$

$$\therefore x = 1.267$$

7 Let  $f(x) = \frac{2}{(2x-1)(2x+1)}$ .

(a) Express  $f(x)$  in partial fractions.

[2]

$$\frac{A}{2x-1} + \frac{B}{2x+1}$$

$$2 = A(2x+1) + B(2x-1)$$

$$2 = 2Ax + A + 2Bx - B$$

$$\therefore 2 = A - B$$

$$0 = 2A + 2B$$

$$A = 1$$

$$B = -1$$

$$\frac{1}{2x-1} - \frac{1}{2x+1}$$

(b) Using your answer to part (a), show that

$$(f(x))^2 = \frac{1}{(2x-1)^2} - \frac{1}{2x-1} + \frac{1}{2x+1} + \frac{1}{(2x+1)^2}$$

[2]

$$\left[ (2x-1)^{-1} - (2x+1)^{-1} \right] \times \left[ (2x-1)^{-1} - (2x+1)^{-1} \right]$$

$$(2x-1)^{-2} - 2(2x-1)^{-1}(2x+1)^{-1} + (2x+1)^{-2}$$

$$\frac{1}{(2x-1)^2} - 2\left(\frac{1}{2x-1}\right)\left(\frac{1}{2x+1}\right) + \frac{1}{(2x+1)^2}$$

$$\frac{1}{(2x-1)^2} - \frac{2}{(2x-1)(2x+1)} + \frac{1}{(2x+1)^2}$$

$\rightarrow f(x)$

$$\frac{1}{(2x-1)^2} - \frac{1}{2x-1} + \frac{1}{2x+1} + \frac{1}{(2x+1)^2}$$

(c) Hence show that  $\int_1^2 (f(x))^2 dx = \frac{2}{5} + \frac{1}{2} \ln\left(\frac{5}{3}\right)$ . [5]

$$\int \frac{1}{(2x-1)^2} - \int \frac{1}{2x-1} + \int \frac{1}{2x+1} + \int \frac{1}{(2x+1)^2}$$

$$u = (2x-1)$$

$$\int (2x-1)^{-2} - \frac{1}{2} \ln 2x-1 + \frac{1}{2} \ln 2x+1 + \int (2x+1)^{-2}$$

$$- \frac{1}{2} (2x-1)^{-1} - \frac{1}{2} \ln[(2x-1)(2x+1)] + - \frac{1}{2} (2x+1)^{-1}$$

$$\left[ -\frac{1}{2(2x-1)} - \frac{1}{2} \ln(4x^2-1) - \frac{1}{2(2x+1)} \right]_1^2$$

$$\left( -\frac{1}{6} - \frac{1}{2} \ln 15 - \frac{1}{10} \right) - \left( -\frac{1}{2} + \frac{1}{2} \ln 3 - \frac{1}{6} \right)$$

$$-\frac{4}{15} - \frac{1}{2} \ln 15 - \left( \frac{1}{2} \ln 3 - \frac{2}{3} \right)$$

$$-\frac{4}{15} - \frac{1}{2} \ln 15 - \frac{1}{2} \ln 3 + \frac{2}{3}$$

$$\frac{2}{5} - \frac{1}{2} \ln \frac{15}{3}$$

2,

x

- 8 Relative to the origin  $O$ , the points  $A$ ,  $B$  and  $D$  have position vectors given by


$$\vec{OA} = \mathbf{i} + 2\mathbf{j} + \mathbf{k}, \quad \vec{OB} = 2\mathbf{i} + 5\mathbf{j} + 3\mathbf{k} \quad \text{and} \quad \vec{OD} = 3\mathbf{i} + 2\mathbf{k}.$$

A fourth point  $C$  is such that  $ABCD$  is a parallelogram.

- (a) Find the position vector of  $C$  and verify that the parallelogram is not a rhombus.

[5]

$$\vec{AB} = \begin{pmatrix} 2 \\ 5 \\ 3 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix}$$



$$\vec{BD} = \begin{pmatrix} 3 \\ 0 \\ 2 \end{pmatrix} - \begin{pmatrix} 2 \\ 5 \\ 3 \end{pmatrix} = \begin{pmatrix} 1 \\ -5 \\ -1 \end{pmatrix}$$

$$\begin{aligned} \vec{OC} &= \vec{OA} + \vec{AB} + \vec{BD} - \vec{AB} \\ &= \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ -5 \\ -1 \end{pmatrix} = \begin{pmatrix} 2 \\ -3 \\ 0 \end{pmatrix} \end{aligned}$$

Rhombus, all 4 sides are equal, parallel, 2 pairs of sides are equal

$$\begin{aligned} \therefore AB &= CD \quad \text{and} \quad AC = BD \\ CD &= \begin{pmatrix} 3 \\ 0 \\ 2 \end{pmatrix} - \begin{pmatrix} 2 \\ -3 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix} \\ \sqrt{1^2 + 3^2 + 2^2} &= \sqrt{1^2 + 3^2 + 2^2} \\ \sqrt{1^2 + 5^2 + 1^2} &= \sqrt{1^2 + 5^2 + 1^2} \end{aligned}$$

$$\begin{aligned} AC &= \begin{pmatrix} 2 \\ -3 \\ 0 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 \\ -5 \\ -1 \end{pmatrix} \end{aligned}$$

- (b) Find angle  $BAD$ , giving your answer in degrees.

[3]

$AB, AD$

$$AB = \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix}$$

$$AD = \begin{pmatrix} 3 \\ 0 \\ 2 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}$$

$$Q = \frac{2 - 6 + 2}{\sqrt{1^2 + 3^2 + 2^2} \times \sqrt{2^2 + 2^2 + 1^2}}$$

$$= \frac{-2}{\sqrt{14} \times 3}$$

$$\cos^{-1}\left(\frac{2}{\sqrt{14} \times 3}\right) = 79.736, \quad \theta = 100.26^\circ$$

- (c) Find the area of the parallelogram correct to 3 significant figures.

[2]

$$\text{Area} = x \times AB$$

$$= 2.9547 \times \sqrt{14}$$

$$= 11.0593$$

$$= 11.1$$



$$3 \cos 10^\circ = x$$

$$x = 2.9547$$

SOH CAH TOA

- 9 (a) The complex numbers  $u$  and  $w$  are such that

$$u - w = 2i \quad \text{and} \quad uw = 6.$$

Find  $u$  and  $w$ , giving your answers in the form  $x + iy$ , where  $x$  and  $y$  are real and exact. [5]

$$u \times w = 6$$

0

$$u = \frac{6}{w}$$

$$\frac{6}{w} - w = 2i$$

$$\frac{6 - w^2}{w} = 2i$$

$$6 - w^2 = 2iw$$

$$w(2i + w) = 6$$

$$2i + w = \frac{6}{w}$$

$$w = 6 - 2i$$

$$u = \frac{6}{6 - 2i}$$

$$= \frac{6 \times (6 + 2i)}{(6 - 2i)(6 + 2i)}$$

$$= \frac{36 + 12i}{36 - 4(-1)}$$

$$= \frac{36 + 12i}{40}$$

$$u = \frac{9}{10} + \frac{3i}{10}$$


- (b) On a sketch of an Argand diagram, shade the region whose points represent complex numbers  $z$  satisfying the inequalities

$$|z - 2 - 2i| \leq 2, \quad 0 \leq \arg z \leq \frac{1}{4}\pi \quad \text{and} \quad \operatorname{Re} z \leq 3.$$

[5]

$$|z - (2+2i)| \leq 2$$


(2)



10

$$\frac{dV}{dt} = k\sqrt{h}$$

$$V = \frac{1}{3}\pi(3rh^2 - h^3)$$



A tank containing water is in the form of a hemisphere. The axis is vertical, the lowest point is  $A$  and the radius is  $r$ , as shown in the diagram. The depth of water at time  $t$  is  $h$ . At time  $t = 0$  the tank is full and the depth of the water is  $r$ . At this instant a tap at  $A$  is opened and water begins to flow out at a rate proportional to  $\sqrt{h}$ . The tank becomes empty at time  $t = 14$ .

The volume of water in the tank is  $V$  when the depth is  $h$ . It is given that  $V = \frac{1}{3}\pi(3rh^2 - h^3)$ .

(a) Show that  $h$  and  $t$  satisfy a differential equation of the form

$$\frac{dh}{dt} = -\frac{B}{2rh^{\frac{1}{2}} - h^{\frac{3}{2}}},$$

where  $B$  is a positive constant.

0 [4]

??

$$3rh^2 - h^3 = \frac{3V}{\pi}$$



[8]

This image shows a full page of white paper with horizontal dashed lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

[illegible]

**BLANK PAGE**

**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cambridgeinternational.org](http://www.cambridgeinternational.org) after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.