

58  
75

CANDIDATE  
NAME

Fuzail

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

**MATHEMATICS**

**9709/32**

Paper 3 Pure Mathematics 3 (**P3**)

**February/March 2017**

**1 hour 45 minutes**

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

**DO NOT WRITE IN ANY BARCODES.**

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The total number of marks for this paper is 75.

This document consists of 19 printed pages and 1 blank page.

- 1 Solve the equation  $\ln(1 + 2^x) = 2$ , giving your answer correct to 3 decimal places.

[3]

$$1 + 2^x = e^2$$

$$2^x = e^2 - 1$$

$$\ln 2^x = \ln(e^2 - 1)$$

$$x = \frac{\ln(e^2 - 1)}{\ln 2}$$

$$x = 2.6756$$

$$x = 2.676 \text{ (3 dp)}$$

5.5 min

$$\begin{aligned} (x-4)^2 &< 2(3x+1)^2 \\ (x-4)(x-4) &< 4(3x+1)(3x+1) \\ x^2 - 8x + 16 &< 4(9x^2 + 6x + 1) \\ x^2 - 8x + 16 &< 36x^2 + 24x + 4 \\ 35x^2 + 32x - 12 &> 0 \\ 35x^2 + 32x - 12 &> 0 \\ x = -\frac{6}{5} & \quad x = \frac{2}{5} \end{aligned}$$

$$x < -\frac{6}{5}, \quad x > \frac{2}{5}$$

$$\begin{aligned} \text{Testing } ① & \quad \text{let } x = -2 \\ | -2 - 4 | &< 2 | 3(-2)^2 + 1 | \\ 6 &< 10 \end{aligned}$$


$$\begin{aligned} \text{Testing } ② & \quad \text{let } x = 1 \\ | 1 - 4 | &< 2 | 3(1)^2 + 1 | \\ 3 &< 8 \end{aligned}$$

$$\begin{aligned} x > \frac{2}{5} & \\ x < -\frac{6}{5} & \end{aligned}$$

$$y = |x - 4|$$

$$y = 2(3x^3 + 1)$$

- 2 Solve the inequality  $|x - 4| < 2|3x + 1|$ .



$$2(1) = 2$$

$$2|3x + 1| = 0$$

$$3x + 1 = 0$$

$$3x = -1$$

$$x = -\frac{1}{3}$$

$$y = -x + 4 \quad ? \quad 6x + 2$$

$$2 \quad ? \quad 7x$$

$$(x > \frac{2}{7})$$

$$y = -x + 4 \quad = -6x - 2$$


$$\therefore x = -6$$

$$x = -\frac{6}{9}$$

$$(x < -\frac{6}{9})$$

- 3 (i) By sketching suitable graphs, show that the equation  $e^{-\frac{1}{2}x} = 4 - x^2$  has one positive root and one negative root.

[2]



- (ii) Verify by calculation that the negative root lies between  $-1$  and  $-1.5$ .

[2]

$$\text{Let } f(x) = 4 - x^2 - e^{\frac{1}{2}x}$$

$$f(-1) = 4 - (-1)^2 - e^{\frac{1}{2}} = 1.35$$

$$f(-1.5) = -0.367$$

change of sign,  $\therefore$  root lies between  $-1$  and  $-1.5$

- (iii) Use the iterative formula  $x_{n+1} = -\sqrt{(4 - e^{-\frac{1}{2}x_n})}$  to determine this root correct to 2 decimal places.  
 Give the result of each iteration to 4 decimal places.

[3]

2

$$\text{let } x_1 = -1.25$$

$$\therefore x_2 = -\sqrt{4 - e^{-\frac{1}{2}(-1.25)}} = 1.4601$$

$$x_3 = -1.3874$$

$$x_4 = -1.4138$$

$$x_5 = -1.4044$$

$$x_6 = -1.4078$$

$$x_7 = -1.4066$$

$$x_8 = -1.4070$$

-1.41  
 ↓

$$\therefore x = -1.40$$

- 4 (i) Express  $8 \cos \theta - 15 \sin \theta$  in the form  $R \cos(\theta + \alpha)$ , where  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ , stating the exact value of  $R$  and giving the value of  $\alpha$  correct to 2 decimal places. [3]

$$R = \sqrt{(-15)^2 + (8)^2} = 17$$

$$\alpha = \tan^{-1}\left(\frac{15}{8}\right) = 61.9275^\circ \approx 61.93^\circ$$

$$17 \cos(\theta + 61.93)$$

(ii) Hence solve the equation

$$8 \cos 2x - 15 \sin 2x = 4,$$

for  $0^\circ < x < 180^\circ$ .

[4]

$$17 \cos(2x + 61.93) = 4$$

$$2x + 61.93 = \cos^{-1}\left(\frac{4}{17}\right)$$

*we use remode?*

$$2x + 61.93 = 76.391 \quad \text{or} \quad 2x + 61.93 = 283.609$$

$$x = \underline{\underline{7.23}}$$

$$\text{or} \quad x = \underline{\underline{110.8395}}$$

$$x \approx \underline{\underline{110.84}}$$

red

- 5 The curve with equation  $y = e^{-ax} \tan x$ , where  $a$  is a positive constant, has only one point in the interval  $0 < x < \frac{1}{2}\pi$  at which the tangent is parallel to the  $x$ -axis. Find the value of  $a$  and state the exact value of the  $x$ -coordinate of this point. [7]

(2)

$$y = e^{-ax} \tan x$$

$$\frac{dy}{dx} = (-a e^{-ax}) (\tan x) + (e^{-ax}) (\sec^2 x)$$

$$= -a e^{-ax} \tan x + e^{-ax} \sec^2 x = 0$$

$$e^{-ax} (-a \tan x + \sec^2 x) = 0$$

$$e^{-ax} = 0$$

X

$$\sec^2 x = a \tan x$$

$$\frac{1}{\cos^2 x} = a \frac{\sin x}{\cos x}$$

$$a = \frac{\sec^2 x}{\tan x}$$

$$\begin{aligned} & a \sin x \cos x = 1 \\ & a 2 \sin x \cos x = 2 \quad \times 2 \end{aligned}$$

$$a \sin 2x = 2$$

$$\sin 2x = \frac{2}{a}$$

$$2x = \sin^{-1} \left( \frac{2}{a} \right)$$

$$x = \frac{1}{2} \sin^{-1} \left( \frac{2}{a} \right)$$




- 6 The line  $l$  has equation  $\mathbf{r} = \mathbf{i} + 2\mathbf{j} - 3\mathbf{k} + \lambda(2\mathbf{i} - \mathbf{j} + \mathbf{k})$ . The plane  $p$  has equation  $3x + y - 5z = 20$ .

(i) Show that the line  $l$  lies in the plane  $p$ .

[3]

X

- (ii) A second plane is parallel to  $l$ , perpendicular to  $p$  and contains the point with position vector  $3\mathbf{i} - \mathbf{j} + 2\mathbf{k}$ . Find the equation of this plane, giving your answer in the form  $ax + by + cz = d$ . [5]



$$2xy = 2$$

A water tank has vertical sides and a horizontal rectangular base, as shown in the diagram. The area of the base is  $2 \text{ m}^2$ . At time  $t = 0$  the tank is empty and water begins to flow into it at a rate of  $1 \text{ m}^3$  per hour. At the same time water begins to flow out from the base at a rate of  $0.2\sqrt{h} \text{ m}^3$  per hour, where  $h \text{ m}$  is the depth of water in the tank at time  $t$  hours.

- (i) Form a differential equation satisfied by  $h$  and  $t$ , and show that the time  $T$  hours taken for the depth of water to reach 4 m is given by

$$T = \int_0^4 \frac{10}{5 - \sqrt{h}} dh.$$

[3]

$1 \text{ m}^3$ <span style="margin-left: 20px;">in</span> <span style="margin-left: 20px;">out</span> $\left[ 0.2h^{\frac{1}{2}}$	$2xy \times h = V$ $2h = V$
<span style="font-size: 2em;">overall,</span> $\frac{dV}{dt} = 1 - 0.2\sqrt{h}$	$\frac{dV}{dh} = 2$
<hr/>	
$\frac{dt}{dh} = \frac{dV}{dh} \times \frac{dt}{dV}$ $= 2 \times \frac{1}{1 - 0.2\sqrt{h}}$ $= \frac{2}{1 - 0.2\sqrt{h}} \times 5$	$\int dt = \int \frac{10}{5 - \sqrt{h}} dh$ $T = \int_0^4 \frac{10}{5 - \sqrt{h}} dh$
<hr/>	
$\frac{dt}{dh} = \frac{10}{5 - \sqrt{h}}$	

- (ii) Using the substitution  $u = 5 - \sqrt{h}$ , find the value of  $T$ .

[6]

$$u = 5 - h^{\frac{1}{2}}$$

$$\sqrt{h} = 5 - u$$

$$\frac{du}{dh} = -\frac{1}{2}h^{-\frac{1}{2}}$$

$$\frac{du}{dh} = -\frac{1}{2\sqrt{h}}$$

$$dh = -2\sqrt{h} du = -2(5 - u)du$$

$$T = \int_3^5 \frac{(10 + 10 + 2u)du}{u}$$

$$T = \int_3^5 \frac{(-100 + 20u)du}{u}$$

$$T = - \int_3^5 \frac{-100}{u} + \frac{20u}{u} du$$

$$= -[-100\ln u + 20u]$$

$$[100\ln u - 20u]_3^5$$

when  $h=0, u=5$   
 $h=4, u=3$

$$\text{sub } 5: -100 + 100\ln 5$$

$$3: -60 + 100\ln 3$$

$$-100 + 100\ln 5 + 60 - 100\ln 3$$

$$T = -40 + 100\ln 5$$

$$= 11.08256 \text{ hours}^3$$

$$= \underline{\underline{11.1 \text{ hours}}}$$

**8 Throughout this question the use of a calculator is not permitted.**

The polynomial  $z^4 + 3z^2 + 6z + 10$  is denoted by  $p(z)$ . The complex number  $-1 + i$  is denoted by  $u$ .

- (i) Showing all your working, verify that  $u$  is a root of the equation  $p(z) = 0$ . [3]

$$\begin{aligned} (-1+i)^4 &= (-1+i)^2 \times (-1+i)^2 \\ &= (-1+i)(-1+i) \\ &= 1 - 2i + i^2 \\ &= -2i \quad \therefore (-1+i)^4 = -2i \times -2i = 4i^2 = -4 \end{aligned}$$

$$\begin{aligned} -4 + 3(-2i) + 6(-1+i) + 10 \\ -4 - 6i - 6 + 6i + 10 \\ \underline{-10 + 10 - 6i + 6i = 0} \end{aligned}$$

- (ii) Find the other three roots of the equation  $p(z) = 0$ . [7]

if  $(-1+i)$  is a root then  $(-1-i)$  is also a root.

$$(x - (-1+i))(x - (-1-i)) ( \ ? \ ) = x^4 + 3x^2 + 6x + 10$$

$$\begin{aligned} (x+1-i)(x+1+i), (x^2 - x^4) &= x^4 + 3x^2 + 6x + 10 \\ x^4 = x \times x \times x \times ? & \\ \therefore ? = x^2 & \\ (x^2 + 2x + 2)(x^2 + 2 + 2) &= x^4 + 3x^2 + 6x + 10 \end{aligned}$$

$$\begin{aligned} x^2 + x + x + x + 1 + 1 - x^2 - i^2 \\ (x^2 + 2x + 2)(x^2 + 2 + 2) &= x^4 + 3x^2 + 6x + 10 \end{aligned}$$

$$x^2 - 2x + 5$$

$$\underline{x^4 + 2x^3 + 3x^2 + 6x + 10} \quad |$$

$$- \underline{x^4 + 2x^3 + 2x^2} \quad |$$

$$0 - 2x^3 + x^2 + 6x \quad |$$

$$- \underline{-2x^3 - 4x^2 - 4x} \quad |$$

$$0 + 5x^2 + 10x + 10 \quad |$$

$$- \underline{5x^2 + 10x + 10} \quad |$$

0

$\therefore x^2 - 2x + 5$  is the root

$$\frac{2 \pm \sqrt{4 - 4(1)(5)}}{2}$$

$$\frac{2 \pm \sqrt{-16}}{2} \Rightarrow \frac{2 \pm \sqrt{16} \times \sqrt{-1}}{2}$$

$$\frac{2 \pm 4i}{2} = 1 \pm 2i$$

roots are  $-1+i, -1-i, 1+2i, 1-2i$

The three roots we found

$\frac{6x-x^2}{(2+x)(4+x^2)}$

16

9 Let  $f(x) = \frac{x(6-x)}{(2+x)(4+x^2)} = \frac{A}{2+x} + \frac{Bx+C}{4+x^2}$

(i) Express  $f(x)$  in partial fractions.

[5]

$$6x - x^2 = 4A + Ax^2 + (Bx+C)(2+x)$$

$$6x - x^2 = 4A + Ax^2 + 2Bx + Bx^2 + 2C + Cx$$

$$\textcircled{1} \quad A+B = -1 \Rightarrow B = -A-1$$

$$\textcircled{2} \quad 2B+C = 6$$

$$\textcircled{3} \quad 4A+2C = 0 \Rightarrow C = -\frac{4A}{2} = -2A$$

$$2(-A-1) + (-2A) = 6$$

$$-2A-2 - 2A = 6$$

$$-4A = 8$$

$$\underline{\underline{A = -2}}$$

$$C = -2(-2) = 4$$

$$B = -(-2)-1 = 1$$

$$\frac{-2}{2+x} + \frac{x+4}{4+x^2}$$

(ii) Hence obtain the expansion of  $f(x)$  in ascending powers of  $x$ , up to and including the term in  $x^2$ . [5]

$$\frac{-2}{2+x} + \frac{x+4}{4+x^2}$$

(3)

$$\begin{aligned}
 @ & -2(2+x)^{-1} \\
 & = -2(2)^{-1} \left(1 + \frac{x}{2}\right)^{-1} \\
 & = -1 \left(1 + \frac{x}{2}\right)^{-1} \\
 & -1 \left[ 1 + (-1)\left(\frac{x}{2}\right) + \frac{(-1)(-2)}{2!} \left(\frac{x}{2}\right)^2 \right] = -1 \left[ 1 - \frac{x}{2} + \frac{1}{4}x^2 \right] \\
 & = -1 + \frac{1}{2}x - \frac{1}{4}x^2
 \end{aligned}$$


$$\begin{aligned}
 @ & (x+4)(4+x^2)^{-1} \\
 & (x+4) \times 4^{-1} \left(1 + \frac{x^2}{4}\right)^{-1} \\
 & (x+4) \times \frac{1}{4} \left[ 1 + (-1)\left(\frac{x^2}{4}\right)^1 \right] \Rightarrow (x+4) \times \frac{1}{4} \left(1 - \frac{1}{4}x^2\right) \\
 & = (x+4) \left( \frac{1}{4} - \frac{1}{4}x^2 \right) \Rightarrow \frac{1}{4}x - \frac{1}{4}x^3 + 1 - x^2
 \end{aligned}$$

$$\therefore -\cancel{\frac{1}{2}x} - \cancel{\frac{1}{4}x^2} + \frac{1}{4}x - \cancel{\frac{1}{4}x^3} + 1 - x^2$$

$$\frac{3}{4}x - \frac{5}{4}x^2 - \frac{1}{4}x^3$$

$$\therefore \frac{3}{4}x - \frac{5}{4}x^2$$

10



2.7.8..

The diagram shows the curve  $y = (\ln x)^2$ . The  $x$ -coordinate of the point  $P$  is equal to  $e$ , and the normal to the curve at  $P$  meets the  $x$ -axis at  $Q$ .

- (i) Find the  $x$ -coordinate of  $Q$ .

[4]

$$y = (\ln x)^2$$

$$\frac{dy}{dx} = 2 \ln x \left(\frac{1}{x}\right) = \frac{2 \ln x}{x}$$

$$\therefore \text{gradient of normal} = -\frac{1}{2 \ln x}$$

$$y = -\frac{1}{2 \ln x}(x) + C$$

$$C = y + \frac{x^2}{2 \ln x} \Rightarrow 1 + \frac{e^2}{2} = 4.6945$$

Geo-ordinate of  $P$   
 $y = (\ln e)^2$   
 $= 1$

 $(e, 1)$ 

$$\frac{x^2}{2 \ln x} = C - y$$

$$x^2 (2 \ln x)^{-1} = C - y$$

$$x^2 \times \frac{1}{2} \times \ln x^{-1} = C - y$$

$$x^2 x - \ln x = 2(C - y)$$

$$y = 0$$

$$-x^2 \ln x = 2C$$

$$-\ln x^{x^2} = 2C$$

$$x^{x^2} = 9.389$$

$$e^{\frac{3}{2}} \\ \sqrt{e}$$

- (ii) Show that  $\int \ln x dx = x \ln x - x + c$ , where  $c$  is a constant.

[1]

- (iii) Using integration by parts, or otherwise, find the exact value of the area of the shaded region between the curve, the  $x$ -axis and the normal  $PQ$ . ~~[5]~~

- [5] -

**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cie.org.uk](http://www.cie.org.uk) after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.