

## Open Mathematics

## Research Article

Wang Xiao\*

# On the power sum problem of Lucas polynomials and its divisible property

<https://doi.org/10.1515/math-2018-0063>

Received November 15, 2017; accepted May 15, 2018.

**Abstract:** The main purpose of this paper is to use the mathematical induction and the properties of Lucas polynomials to study the power sum problem of Lucas polynomials. In the end, we obtain an interesting divisible property.

**Keywords:** Lucas polynomials, Power sum problem, Mathematical induction, Divisible property

**MSC:** 11B39

## 1 Introduction

For any integer  $n \geq 0$ , the famous Fibonacci polynomials  $\{F_n(x)\}$  and Lucas polynomials  $\{L_n(x)\}$  are defined by  $F_0(x) = 0$ ,  $F_1(x) = 1$ ,  $L_0(x) = 2$ ,  $L_1(x) = x$  and  $F_{n+2}(x) = xF_{n+1}(x) + F_n(x)$ ,  $L_{n+2}(x) = xL_{n+1}(x) + L_n(x)$ . The general terms of  $F_n(x)$  and  $L_n(x)$  are given by

$$F_{n+1}(x) = \sum_{k=0}^{\left[\frac{n}{2}\right]} \binom{n-k}{k} x^{n-2k}$$

and

$$L_n(x) = \sum_{k=0}^{\left[\frac{n}{2}\right]} \frac{n}{n-k} \binom{n-k}{k} x^{n-2k}, \quad (1)$$

where  $\binom{m}{n} = \frac{m!}{n!(m-n)!}$ , and  $[x]$  denotes the greatest integer  $\leq x$ .

It is easy to prove the identities

$$F_n(x) = \frac{1}{\sqrt{x^2+4}} \left[ \left( \frac{x+\sqrt{x^2+4}}{2} \right)^n - \left( \frac{x-\sqrt{x^2+4}}{2} \right)^n \right]$$

and

$$L_n(x) = \left( \frac{x+\sqrt{x^2+4}}{2} \right)^n + \left( \frac{x-\sqrt{x^2+4}}{2} \right)^n. \quad (2)$$

If we take  $x = 1$ , then  $\{F_n(x)\}$  becomes Fibonacci sequences  $\{F_n(1)\}$ , and  $\{L_n(x)\}$  becomes Lucas sequences  $\{L_n(1)\}$ .

Since these sequences and polynomials have very important positions in the theory and application of mathematics, many scholars have studied their various properties, and obtained a series of important results,

\*Corresponding Author: Wang Xiao: School of Mathematics, Northwest University, Xi'an, Shaanxi, China,  
E-mail: wangxiao\_0606@stumail.nwu.edu.cn

some of which can be found in references [1-5], and some other related papers can also be found in [6-16]. For example, in a private communication with Curtis Cooper, R. S. Melham suggested that it would be interesting to discover an explicit expansion for

$$L_1 L_2 \cdots L_{2m+1} \sum_{k=1}^n F_{2k}^{2m+1}$$

as a polynomial in  $F_{2n+1}$ . Wiemann and Cooper [1] reported some conjectures of Melham related to the sum  $\sum_{k=1}^n F_{2k}^{2m+1}$ . Kiyota Ozeki [2] proved that

$$\sum_{k=1}^n F_{2k}^{2m+1} = \frac{1}{5^m} \sum_{j=0}^m \frac{(-1)^j}{L_{2m+1-2j}} \binom{2m+1}{j} (F_{(2m+1-2j)(2n+1)} - F_{2m+1-2j}).$$

Helmut Prodinger [3] studied the more general summation  $\sum_{k=0}^n F_{2k+\delta}^{2m+1+\epsilon}$ , where  $\delta, \epsilon \in \{0, 1\}$ , and obtained many interesting identities.

For the sum of powers of Lucas numbers, Helmut Prodinger [3] also obtained some similar conclusions.

R. S. Melham [4] also proposed the following two conjectures:

**Conjecture A.** *Let  $m$  be a positive integer. Then the sum*

$$L_1 L_3 L_5 \cdots L_{2m+1} \sum_{k=1}^n F_{2k}^{2m+1}$$

*can be expressed as  $(F_{2n+1} - 1)^2 P_{2m-1}(F_{2n+1})$ , where  $P_{2m-1}(x)$  is a polynomial of degree  $2m - 1$  with integer coefficients.*

**Conjecture B.** *Let  $m \geq 0$  be an integer. Then the sum*

$$L_1 L_3 L_5 \cdots L_{2m+1} \sum_{k=1}^n L_{2k}^{2m+1}$$

*can be expressed as  $(L_{2n+1} - 1) Q_{2m}(L_{2n+1})$ , where  $Q_{2m}(x)$  is a polynomial of degree  $2m$  with integer coefficients.*

Wang Tingting and Zhang Wenpeng [5] studied these problems, and proved several conclusions for  $\sum_{k=1}^h F_k^n(x)$  and  $\sum_{k=1}^h L_k^n(x)$ , where  $h$  and  $n$  are any positive integers.

As some applications of Theorem 2 in reference [5], they deduced the following:

**Corollary A.** *Let  $h \geq 1$  and  $n \geq 0$  be two integers. Then the sum*

$$L_1(x) L_3(x) L_5(x) \cdots L_{2n+1}(x) \sum_{m=1}^h L_{2m}^{2n+1}(x)$$

*can be expressed as  $(L_{2h+1}(x) - x) Q_{2n}(x, L_{2h+1}(x))$ , where  $Q_{2n}(x, y)$  is a polynomial in two variables  $x$  and  $y$  with integer coefficients and degree  $2n$  of  $y$ .*

**Corollary B.** *Let  $h \geq 1$  and  $n \geq 0$  be two integers. Then the sum*

$$L_1(x) L_3(x) L_5(x) \cdots L_{2n+1}(x) \sum_{m=1}^h F_{2m}^{2n+1}(x)$$

*can be expressed as  $(F_{2h+1}(x) - 1) H_{2n}(x, F_{2h+1}(x))$ , where  $H_{2n}(x, y)$  is a polynomial in two variables  $x$  and  $y$  with integer coefficients and degree  $2n$  of  $y$ .*

Therefore, Wang Tingting and Zhang Wenpeng [5] solved Conjecture B completely. They also obtained some substantial progress for Conjecture A.

It is easy to see that in the above Conjecture A and Conjecture B, the subscripts of  $F_n$  and  $L_n$  in sums  $\sum_{k=1}^n F_{2k}^{2m+1}$  and  $\sum_{k=1}^n L_{2k}^{2m+1}$  are even numbers. Inspired by the above researches, we naturally ask that for

any positive integers  $n$  and  $h$  with  $h \geq 2$ , whether there exists a polynomial  $U(x)$  with degree  $> 1$  such that the congruence

$$L_1(x)L_3(x)\cdots L_{2n+1}(x) \sum_{m=0}^{h-1} L_{2m+1}^{2n+1}(x) \equiv 0 \pmod{U(x)}. \quad (3)$$

If so, what would  $U(x)$  look like? How is it different from the form in Conjecture A and Conjecture B? Obviously, it is hard to guess the exact form of  $U(x)$  from (d) of Theorem 1 in reference [5]. Maybe that is why there is no such formal conjecture in [4].

In this paper, we shall use the mathematical induction and the properties of Lucas polynomials to study this problem, and give an exact polynomial  $U(x)$  in (3). The result is detailed in the following theorems.

**Theorem A.** For any positive integers  $n$  and  $h$  with  $h \geq 2$ , we have

$$L_1(x)L_3(x)\cdots L_{2n+1}(x) \sum_{m=0}^{h-1} L_{2m+1}^{2n+1}(x) \equiv 0 \pmod{x \cdot (L_{2h}(x) - 2)}.$$

**Theorem B.** For any positive integers  $n$  and  $h$  with  $h \geq 2$ , we have

$$L_1(x)L_3(x)\cdots L_{2n+1}(x) \sum_{m=0}^{h-1} F_{2m+1}^{2n+1}(x) \equiv 0 \pmod{x \cdot F_{2h}(x)}.$$

Taking  $x = 1$ , from Theorem 1 and Theorem 2 we may immediately deduce the following two corollaries:

**Corollary A.** For any positive integers  $n$  and  $h$  with  $h \geq 2$ , we have

$$L_1L_3\cdots L_{2n+1} \sum_{m=0}^{h-1} L_{2m+1}^{2n+1} \equiv 0 \pmod{(L_{2h} - 2)}.$$

**Corollary B.** For any positive integers  $n$  and  $h$  with  $h \geq 2$ , we have

$$L_1L_3\cdots L_{2n+1} \sum_{m=0}^{h-1} F_{2m+1}^{2n+1} \equiv 0 \pmod{F_{2h}}.$$

## 2 Two Lemmas

**Lemma A.** For any non-negative integer  $n$  and  $k$ , we have the identity

$$L_n(L_{2k+1}(x)) = L_{n(2k+1)}(x).$$

*Proof.* Let  $\alpha = \frac{x+\sqrt{x^2+4}}{2}$ ,  $\beta = \frac{x-\sqrt{x^2+4}}{2}$ , and replace  $x$  by  $L_{2k+1}(x)$  in (2). Since we have  $\alpha^{2k+1}\beta^{2k+1} = -1$ ,

$$\begin{aligned} L_{2k+1}(x) + \sqrt{L_{2k+1}^2(x) + 4} &= \alpha^{2k+1} + \beta^{2k+1} + \sqrt{(\alpha^{2k+1} + \beta^{2k+1})^2 + 4} \\ &= \alpha^{2k+1} + \beta^{2k+1} + \sqrt{\alpha^{2(2k+1)} + \beta^{2(2k+1)} - 2 + 4} \\ &= \alpha^{2k+1} + \beta^{2k+1} + \sqrt{(\alpha^{2k+1} - \beta^{2k+1})^2} = 2\alpha^{2k+1} \end{aligned}$$

and

$$\begin{aligned} L_{2k+1}(x) - \sqrt{L_{2k+1}^2(x) + 4} &= \alpha^{2k+1} + \beta^{2k+1} - \sqrt{(\alpha^{2k+1} + \beta^{2k+1})^2 + 4} \\ &= \alpha^{2k+1} + \beta^{2k+1} - \sqrt{(\alpha^{2k+1} - \beta^{2k+1})^2} = 2\beta^{2k+1}, \end{aligned}$$

from (2) we have the identity

$$L_n(L_{2k+1}(x)) = \left( \frac{L_{2k+1} + \sqrt{L_{2k+1}^2 + 4}}{2} \right)^n + \left( \frac{L_{2k+1} - \sqrt{L_{2k+1}^2 + 4}}{2} \right)^n$$

$$= \alpha^{n(2k+1)} + \beta^{n(2k+1)} = L_{n(2k+1)}(x).$$

This completes the proof of Lemma 1.  $\square$

**Lemma B.** For any positive integers  $n$  and  $h$ , we have the congruence

$$L_1(x) (L_{2h(2n+1)}(x) - 2) - (2n+1)L_{2n+1}(x) (L_{2h}(x) - 2) \equiv 0 \pmod{x(L_{2h}(x) - 2)}.$$

*Proof.* From (1) we know that  $x \mid L_{2n+1}(x)$ . Because that  $L_1(x) = x$ , so in order to prove Lemma 2, we only need to prove the congruence

$$L_{2h(2n+1)}(x) - 2 \equiv 0 \pmod{(L_{2h}(x) - 2)}. \quad (4)$$

Next we prove (4) by complete induction. It is clear that (4) is true for  $n = 0$ . If  $n = 1$ , then apply the identity

$$L_{2h}^3(x) = (\alpha^{2h} + \beta^{2h})^3 = L_{6h}(x) + 3L_{2h}(x),$$

we have

$$\begin{aligned} L_{6h}(x) - 2 &= L_{2h}^3(x) - 3L_{2h}(x) - 2 = (L_{2h}(x) - 2)(L_{2h}(x) + 1)^2 \\ &\equiv 0 \pmod{(L_{2h}(x) - 2)}. \end{aligned}$$

That is to say, the congruence (4) is true for  $n = 1$ .

Suppose that the congruence (4) is true for all integers  $0 \leq n \leq s$ . That is,

$$L_{2h(2n+1)}(x) - 2 \equiv 0 \pmod{(L_{2h}(x) - 2)} \quad (5)$$

holds for all integers  $0 \leq n \leq s$ .

Then for positive integer  $n = s + 1$ , we have the identities

$$L_{4h}(x) = L_{2h}^2(x) - 2 \equiv 2 \pmod{(L_{2h}(x) - 2)}$$

and

$$L_{4h}(x)L_{2h(2s+1)}(x) = L_{2h(2s+3)}(x) + L_{2h(2s-1)}(x),$$

from (5) we can deduce the congruence equations as follows

$$\begin{aligned} L_{2h(2s+3)}(x) - 2 &= L_{4h}(x)L_{2h(2s+1)}(x) - L_{2h(2s-1)}(x) - 2 \\ &\equiv 2L_{2h(2s+1)}(x) - L_{2h(2s-1)}(x) - 2 \\ &\equiv 2(L_{2h(2s+1)}(x) - 2) - (L_{2h(2s-1)}(x) - 2) \\ &\equiv 0 \pmod{(L_{2h}(x) - 2)}. \end{aligned}$$

That is to say, the congruence (4) is true for  $n = s + 1$ .

Now Lemma 2 follows from (4) and completes the induction.  $\square$

### 3 Proofs of the theorems

In this section, we shall use mathematical induction to complete the proofs of our theorems. Here we only prove Theorem 1. Similarly, we can also deduce Theorem 2 and thus we omit its proving process here. After replacing  $x$  by  $L_{2m+1}(x)$  in (1), we obtain the following expression with Lemma 1

$$L_{(2n+1)(2m+1)}(x) = L_{2n+1}(L_{2m+1}(x)) = \sum_{k=0}^n \frac{2n+1}{2n+1-k} \binom{2n+1-k}{k} L_{2m+1}^{2n+1-2k}(x)$$

or

$$L_{(2n+1)(2m+1)}(x) - (2n+1)L_{2m+1}(x) = \sum_{k=0}^{n-1} \frac{2n+1}{2n+1-k} \binom{2n+1-k}{k} L_{2m+1}^{2n+1-2k}(x). \quad (6)$$

For any positive integer  $h \geq 2$ , we first introduce that the identities

$$\begin{aligned} \sum_{m=0}^{h-1} L_{(2n+1)(2m+1)}(x) &= \sum_{m=0}^{h-1} \left( \alpha^{(2n+1)(2m+1)} + \beta^{(2n+1)(2m+1)} \right) \\ &= \alpha^{2n+1} \cdot \frac{\alpha^{2h(2n+1)} - 1}{\alpha^{2(2n+1)} - 1} + \beta^{2n+1} \cdot \frac{\beta^{2h(2n+1)} - 1}{\beta^{2(2n+1)} - 1} \\ &= \frac{\alpha^{2h(2n+1)} - 1}{\alpha^{2n+1} + \beta^{2n+1}} + \frac{\beta^{2h(2n+1)} - 1}{\beta^{2n+1} + \alpha^{2n+1}} = \frac{L_{2h(2n+1)}(x) - 2}{L_{2n+1}(x)}, \end{aligned} \quad (7)$$

$$\sum_{m=0}^{h-1} L_{2m+1}(x) = \sum_{m=0}^{h-1} \left( \alpha^{2m+1} + \beta^{2m+1} \right) = \frac{L_{2h}(x) - 2}{L_1(x)}. \quad (8)$$

Then, combining (6), (7) and (8) we have

$$\begin{aligned} \sum_{m=0}^{h-1} [L_{(2n+1)(2m+1)}(x) - (2n+1)L_{2m+1}(x)] &= \frac{L_{2h(2n+1)}(x) - 2}{L_{2n+1}(x)} - (2n+1) \cdot \frac{L_{2h}(x) - 2}{L_1(x)} \\ &= \sum_{k=0}^{n-1} \frac{2n+1}{2n+1-k} \binom{2n+1-k}{k} \sum_{m=0}^{h-1} L_{2m+1}^{2n+1-2k}(x). \end{aligned} \quad (9)$$

Now we apply (9) and mathematical induction to prove the congruence

$$L_1(x)L_3(x) \cdots L_{2n+1}(x) \sum_{m=0}^{h-1} L_{2m+1}^{2n+1}(x) \equiv 0 \pmod{x(L_{2h}(x) - 2)}. \quad (10)$$

If  $n = 1$ , then from (9) we have

$$L_1(x)L_3(x) \left[ \frac{L_{6h}(x) - 2}{L_3(x)} - 3 \cdot \frac{L_{2h}(x) - 2}{L_1(x)} \right] = L_1(x)L_3(x) \sum_{m=0}^{h-1} L_{2m+1}^3(x). \quad (11)$$

From Lemma 2 we know that

$$L_1(x)L_3(x) \left[ \frac{L_{6h}(x) - 2}{L_3(x)} - 3 \cdot \frac{L_{2h}(x) - 2}{L_1(x)} \right] \equiv 0 \pmod{x(L_{2h}(x) - 2)}. \quad (12)$$

Combining (11) and (12) we know that the congruence (10) is true for  $n = 1$ .

Suppose that (10) is true for all  $1 \leq n \leq s$ . That is,

$$L_1(x)L_3(x) \cdots L_{2n+1}(x) \sum_{m=0}^{h-1} L_{2m+1}^{2n+1}(x) \equiv 0 \pmod{x(L_{2h}(x) - 2)} \quad (13)$$

holds for all  $1 \leq n \leq s$ .

Then for  $n = s + 1$ , from (9) we have

$$\begin{aligned} &\frac{L_{2h(2s+3)}(x) - 2}{L_{2s+3}(x)} - (2s+3) \cdot \frac{L_{2h}(x) - 2}{L_1(x)} \\ &= \sum_{k=0}^s \frac{2s+3}{2s+3-k} \binom{2s+3-k}{k} \sum_{m=0}^{h-1} L_{2m+1}^{2s+3-2k}(x) \\ &= \sum_{m=0}^{h-1} L_{2m+1}^{2s+3}(x) + \sum_{k=1}^s \frac{2s+3}{2s+3-k} \binom{2s+3-k}{k} \sum_{m=0}^{h-1} L_{2m+1}^{2s+3-2k}(x). \end{aligned} \quad (14)$$

Applying Lemma 2 we have the congruence

$$L_1(x)L_3(x)\cdots L_{2s+3}(x) \left[ \frac{L_{2h(2s+3)}(x) - 2}{L_{2s+3}(x)} - (2s+3) \cdot \frac{L_{2h}(x) - 2}{L_1(x)} \right] \\ \equiv 0 \pmod{x(L_{2h}(x) - 2)}. \quad (15)$$

If  $1 \leq k \leq s$ , then  $3 \leq 2s+3-2k \leq 2s+1$ . From the inductive assumption (13) we have

$$L_1(x)L_3(x)\cdots L_{2s+1}(x)L_{2s+3}(x) \sum_{k=1}^s \frac{2s+3}{2s+3-k} \binom{2s+3-k}{k} \sum_{m=0}^{h-1} L_{2m+1}^{2s+3-2k}(x) \\ \equiv 0 \pmod{x(L_{2h}(x) - 2)}. \quad (16)$$

Combining (14), (15) and (16) we may immediately deduce the congruence

$$L_1(x)L_3(x)\cdots L_{2s+1}(x)L_{2s+3}(x) \sum_{m=0}^{h-1} L_{2m+1}^{2s+3}(x) \equiv 0 \pmod{x(L_{2h}(x) - 2)}.$$

This completes the proof of our theorem by mathematical induction.

### Competing interests

The author declare that they have no competing interests.

**Acknowledgement:** This work is supported by the N. S. F. (Grant No. 11771351) of P. R. China. The author would like to thank the referees for their very helpful and detailed comments, which have significantly improved the presentation of this paper.

## References

- [1] Wiemann, M, Cooper C: Divisibility of an F-L Type Convolution, Applications of Fibonacci Numbers, Vol. 9, Kluwer Acad. Publ., Dordrecht, 2004, 267-287
- [2] Kiyota Ozeki, On Melham's sum, The Fibonacci Quarterly, 2008/2009, 46/47, 107-110
- [3] Helmut Prodinger, On a sum of Melham and its variants, The Fibonacci Quarterly, 2008/2009, 46/47, 207-215
- [4] Melham, R. S: Some conjectures concerning sums of odd powers of Fibonacci and Lucas numbers, The Fibonacci Quarterly, 2008/2009, 46/47, 312-315
- [5] Wang, T, Zhang, W: Some identities involving Fibonacci. Lucas polynomials and their applications Bull. Math. Soc. Sci. Math. Roumanie, 2012, 55, 95-103
- [6] Li, X: Some identities involving Chebyshev polynomials. Mathematical Problems in Engineering, 2015, 2015, Article ID 950695
- [7] Ma, Y, Lv, X: Several identities involving the reciprocal sums of Chebyshev polynomials. Mathematical Problems in Engineering, 2017, 2017, Article ID 4194579
- [8] Yi, Y, Zhang, W: Some identities involving the Fibonacci polynomials, The Fibonacci Quarterly, 2002, 40, 314-318
- [9] Ma, R, Zhang, W: Several identities involving the Fibonacci numbers and Lucas numbers, The Fibonacci Quarterly, 2007, 45, 164-170
- [10] Kim, D. S, Kim, T, Lee, S: Some identities for Bernoulli polynomials involving Chebyshev polynomials. J. Comput. Anal. Appl., 2014, 16, 172-180
- [11] Kim, D. S, Dolgy, D. V, Kim, T, Rim, S. H: Identities involving Bernoulli and Euler polynomials arising from Chebyshev polynomials. Proc. Jangjeon Math. Soc., 2012, 15, 361-370
- [12] Kim, T, Kim, D. S, Seo, J. J, Dolgy, D. V: Some identities of Chebyshev polynomials arising from non-linear differential equations. J. Comput. Anal. Appl., 2017, 23, 820-832
- [13] He, Y: Some new results on products of Apostol-Bernoulli and Apostol-Euler polynomials, J. Math. Anal. Appl., 2015, 431, 34-46
- [14] He, Y, Wang, C. P: New symmetric identities involving the Eulerian polynomials, J. Comput. Anal. Appl., 2014, 17, 498-504
- [15] He, Y, Kim, D. S: General convolution identities for Apostol-Bernoulli, Euler and Genocchi polynomials, J. Nonlinear Sci. Appl., 2016, 9, 4780-4797
- [16] Chen, L, Zhang, W: Chebyshev polynomials and their some interesting applications, Advances in Difference Equations, 2017, 2017: 303