


On The Coloring of Graphs and Chromatic Polynomials

Ian Cavey, Christian Sprague, Mack Stannard

Boise State University


November 9, 2014

Introduction




- G is an example of a Graph

Introduction




- G is an example of a Graph
- Graphs are made up of:
 - Vertices

Introduction




- G is an example of a Graph
- Graphs are made up of:
 - Vertices
 - Edges

Introduction




- Assigning a color to every vertex in a graph is called graph coloring

Introduction




- Assigning a color to every vertex in a graph is called graph coloring
- A **proper** coloring of a graph is a coloring where any two vertices connected by an edge are not colored identically

Introduction



- Assigning a color to every vertex in a graph is called graph coloring
- A **proper** coloring of a graph is a coloring where any two vertices connected by an edge are not colored identically
- Is the above coloring proper or improper?

Introduction



- Assigning a color to every vertex in a graph is called graph coloring
- A proper coloring of a graph is a coloring where any two vertices connected by an edge are not colored identically
- Is the above coloring proper or improper?
 - Improper!

Introduction

Natural Questions

- What is the fewest number of colors needed to properly color a graph?

Introduction

Natural Questions

- What is the fewest number of colors needed to properly color a graph?
- How many ways can a graph be properly colored with x colors?

Chromatic Polynomials and Numbers

The Chromatic Number is the fewest number of colors needed to properly color a graph.

Chromatic Polynomials and Numbers

The Chromatic Number is the fewest number of colors needed to properly color a graph.

- How many ways are there to properly color a graph G with x colors?

Chromatic Polynomials and Numbers

The Chromatic Number is the fewest number of colors needed to properly color a graph.

- How many ways are there to properly color a graph G with x colors?
- We can create a function of a graph G and a number of colors x which is the number of ways to color G properly with x colors

Chromatic Polynomials and Numbers

The Chromatic Number is the fewest number of colors needed to properly color a graph.

- How many ways are there to properly color a graph G with x colors?
- We can create a function of a graph G and a number of colors x which is the number of ways to color G properly with x colors
- This function is actually a polynomial, called the Chromatic Polynomial, and is denoted $F(G, x)$


Reduction Algorithm

- Finding the Chromatic Polynomial for a graph is not always easy!

Reduction Algorithm


- Finding the Chromatic Polynomial for a graph is not always easy!
- But thanks to Birkhoff and Lewis, we have a simple algorithm for computing $F(G, x)$ given a graph G .

Reduction Algorithm




- Consider two vertices a and b that are connected by edge e .

Reduction Algorithm




- Consider two vertices a and b that are connected by edge e .
- Since we want proper colorings, a and b must be colored differently

Reduction Algorithm




- Consider two vertices a and b that are connected by edge e .
- Since we want proper colorings, a and b must be colored differently
- To determine the number of ways in which vertices a , b can be colored differently:

Reduction Algorithm



- Consider two vertices a and b that are connected by edge e .
- Since we want proper colorings, a and b must be colored differently
- To determine the number of ways in which vertices a , b can be colored differently:
 - Find the number of colorings where a and b are colored the same or different


Reduction Algorithm



- Consider two vertices a and b that are connected by edge e .
- Since we want proper colorings, a and b must be colored differently
- To determine the number of ways in which vertices a , b can be colored differently:
 - Find the number of colorings where a and b are colored the same or different
 - Then subtract the number of ways where a and b are colored identically.


Reduction Algorithm

First find the number of colorings where a and b are colored the same or different.



Reduction Algorithm


First find the number of colorings where a and b are colored the same or different.



- Remove edge e

Reduction Algorithm


First find the number of colorings where a and b are colored the same or different.



- Remove edge e
- Call this new graph G'

Reduction Algorithm


First find the number of colorings where a and b are colored the same or different.



- Remove edge e
- Call this new graph G'
- $F(G', x)$ is the number of (otherwise proper) colorings of G where a and b can have the same or different colors.


Reduction Algorithm

Then subtract the number of ways where a and b are colored identically.



Reduction Algorithm


Then subtract the number of ways where a and b are colored identically.



- Merge a and b into one vertex

Reduction Algorithm


Then subtract the number of ways where a and b are colored identically.



- Merge a and b into one vertex
- Call this graph G''

Reduction Algorithm

Then subtract the number of ways where a and b are colored identically.



- Merge a and b into one vertex
- Call this graph G''
- $F(G'', x)$ is the number of (otherwise proper) colorings of G where a and b are colored identically

Reduction Algorithm

According to the algorithm, the chromatic polynomial of G is equal to the difference of the chromatic polynomials of G' and G'' ,

$$F(G, x) = F(G', x) - F(G'', x).$$

Reduction Algorithm

According to the algorithm, the chromatic polynomial of G is equal to the difference of the chromatic polynomials of G' and G'' ,

$$F(G, x) = F(G', x) - F(G'', x).$$

- This describes one iteration of the algorithm

Reduction Algorithm

According to the algorithm, the chromatic polynomial of G is equal to the difference of the chromatic polynomials of G' and G'' ,

$$F(G, x) = F(G', x) - F(G'', x).$$

- This describes one iteration of the algorithm
- Continue in this manner by selecting edges until all of the obtained graphs are without edges

Reduction Algorithm

According to the algorithm, the chromatic polynomial of G is equal to the difference of the chromatic polynomials of G' and G'' ,

$$F(G, x) = F(G', x) - F(G'', x).$$

- This describes one iteration of the algorithm
- Continue in this manner by selecting edges until all of the obtained graphs are without edges
- For each of these *empty graphs* with n vertices and x colors there are x^n different colorings, since each of the n vertices has x possible colorings

Results

- Finally, we introduce some well known types of graphs


Results

- Finally, we introduce some well known types of graphs
- Applying the Birkhoff-Lewis Reduction Algorithm and induction to these graphs yields general formulas for Chromatic Polynomials

Results

Path


A path graph of n vertices, denoted P_n , is a graph that can be arranged in such a way that vertices are only connected to their adjacent vertices



Results

Path

A path graph of n vertices, denoted P_n , is a graph that can be arranged in such a way that vertices are only connected to their adjacent vertices




- The chromatic polynomial for a path graph has a general form,
$$F(P_n, x) = x(x - 1)^{n-1}$$

Results

Cycle


A cycle graph of n vertices, denoted C_n , is composed of an equal number of vertices and edges such that each vertex has exactly two edges connected with it, creating a closed graph.



Results

Cycle

A cycle graph of n vertices, denoted C_n , is composed of an equal number of vertices and edges such that each vertex has exactly two edges connected with it, creating a closed graph.




- The chromatic polynomial for a cycle graph has a general form,
$$F(C, n) = (x - 1)^n + (-1)^{n+1}(1 - x)$$

Results

Complete


A graph with n vertices where any two vertices are connected by an edge is called a complete graph, denoted K_n .



Results

Complete

A graph with n vertices where any two vertices are connected by an edge is called a complete graph, denoted K_n .



- The chromatic polynomial for a complete graph has a general form,
$$F(K_n, x) = x(x-1)(x-2)\cdots(x-n+1) = \frac{x!}{(x-n)!}.$$

Acknowledgments

Thanks to Dr. Andrés Caicedo, Boise State University, for providing the Sage code, which enabled us to explore and make conjectures

Bibliography

Mount Holyoke College *Laboratories in Mathematical Experimentation: A Bridge to Higher Mathematics* 1997: Springer-Verlag New York, Inc.
www.sagemathcloud.com