



Before storage, the catalyst can be washed with distilled water at ambient temperature to remove remaining sodium aluminate. Oxygen-free (degassed) water is preferred for storage to prevent oxidation of the catalyst, which would accelerate its aging process and result in reduced catalytic activity.<sup>[7]</sup>

## Properties

Macroscopically, Raney nickel is a finely divided, grey powder. Microscopically, each particle of this powder is a three-dimensional mesh, with pores of irregular size and shape of which the vast majority is created during the leaching process. Raney nickel is notable for being thermally and structurally stable, as well as having a large Brunauer-Emmett-Teller (BET ) surface area. These properties are a direct result of the activation process and contribute to a relatively high catalytic activity.

The surface area is typically determined by a BET measurement using a gas that is preferentially adsorbed on metallic surfaces, such as hydrogen. Using this type of measurement, almost all the exposed area in a particle of the catalyst has been shown to have Ni on its surface.<sup>[7]</sup> Since Ni is the active metal of the catalyst, a large Ni surface area implies a large surface is available for reactions to occur simultaneously, which is reflected in an increased catalyst activity. Commercially available Raney nickel has an average Ni surface area of 100 m<sup>2</sup> per gram of catalyst.<sup>[7]</sup>


A high catalytic activity, coupled with the fact that hydrogen is absorbed within the pores of the catalyst during activation, makes Raney nickel a useful catalyst for many hydrogenation reactions. Its structural and thermal stability (i.e., it does not decompose at high temperatures) allows its use under a wide range of reaction conditions.<sup>[10][11]</sup> Additionally, the solubility of Raney nickel is negligible in most common laboratory solvents, with the exception of mineral acids such as hydrochloric acid, and its relatively high density (about 6.5 g cm<sup>−3</sup>)<sup>[1]</sup> also facilitates its separation from a liquid phase after a reaction is completed.

## Applications

Raney nickel is used in a large number of industrial processes and in organic synthesis because of its stability and high catalytic activity at room temperature.<sup>[7][12][13]</sup>

### Industrial applications

A practical example of the use of Raney nickel in industry is shown in the following reaction, where benzene is reduced to cyclohexane. Reduction of the benzene ring is very hard to achieve through other chemical means, but can be effected by using Raney nickel. Other heterogeneous catalysts, such as those using platinum group elements, may be used instead, to similar effect, but these tend to be more expensive to produce than Raney nickel. The cyclohexane thus produced may be used in the synthesis of adipic acid, a raw material used in the industrial production of polyamides such as nylon.<sup>[14]</sup>



Benzene is routinely reduced to cyclohexane using Raney nickel for the production of nylon.


Other industrial applications of Raney nickel include the conversion of:

- Dextrose to sorbitol;
- Nitro compounds to amines, for example, 2,4-dinitrotoluene to 2,4-toluenediamine;
- Nitriles to amines, for example, stearonitrile to stearylamine and adiponitrile to hexamethylenediamine;
- Olefins to paraffins, for example, sulfolene to sulfolane;
- Acetylenes to paraffins, for example, 1,4-butynediol to 1,4-butanediol.

### Applications in organic synthesis


#### Desulfurization

Raney nickel is used in organic synthesis for desulfurization. For example, thioacetals will be reduced to hydrocarbons in the last step of the Mozingo reduction:<sup>[14][15]</sup>




Example of desulfurization of thioacetals using Raney nickel


Thiols,<sup>[16]</sup> and sulfides<sup>[17]</sup> can be removed from aliphatic, aromatic, or heteroaromatic compounds. Likewise, Raney nickel will remove the sulfur of thiophene to give a saturated alkane.<sup>[18]</sup>




Phase diagram of the Ni–Al system, showing relevant phases



SEM of Raney nickel catalyst in which crystals of 1-50μm are seen.



A close-up of Raney nickel. Small cracks of approximately 1-100 nm width are seen within the crystals, causing the increased surface area.



Reduction of thiophene by Raney nickel

### Reduction of functional groups

It is typically used in the reduction of compounds with multiple bonds, such as alkynes, alkenes,<sup>[19]</sup> nitriles,<sup>[20]</sup> dienes, aromatics<sup>[21]</sup> and carbonyl-containing compounds. Additionally, Raney nickel will reduce heteroatom-heteroatom bonds, such as hydrazines,<sup>[22]</sup> nitro groups, and nitrosamines.<sup>[23]</sup> It has also found use in the reductive alkylation of amines<sup>[24]</sup> and the amination of alcohols.

When reducing a carbon-carbon double bond, Raney nickel will add hydrogen in a *syn* fashion.<sup>[14]</sup>

## Safety

Due to its large surface area and high volume of contained hydrogen gas, dry, activated Raney nickel is a pyrophoric material that requires handling under an inert atmosphere. Raney nickel is typically supplied as a 50% slurry in water. Even after reaction, residual Raney nickel contains significant amounts of hydrogen gas and may spontaneously ignite when exposed to air.<sup>[25]</sup>

Additionally, acute exposure to Raney nickel may cause irritation of the respiratory tract and nasal cavities, and causes pulmonary fibrosis if inhaled. Ingestion may lead to convulsions and intestinal disorders. It can also cause eye and skin irritation. Chronic exposure may lead to pneumonitis and other signs of sensitization to nickel, such as skin rashes ("nickel itch").<sup>[26]</sup>

Nickel is also rated as being a possible human carcinogen by the IARC (Group 2B, EU category 3) and teratogen, while the inhalation of fine aluminium oxide particles is associated with Shaver's disease.

The pyrophoric nature of Raney nickel initiated the development of safer nickel silicide-based catalysts with similar catalytic properties.<sup>[27]</sup>



Raney nickel is flammable.



Nickel metal is classified as "Harmful".

## Development

Murray Raney graduated as a mechanical engineer from the University of Kentucky in 1909. In 1915 he joined the Lookout Oil and Refining Company in Tennessee and was responsible for the installation of electrolytic cells for the production of hydrogen which was used in the hydrogenation of vegetable oils. During that time the industry used a nickel catalyst prepared from nickel(II) oxide. Believing that better catalysts could be produced, around 1921 he started to perform independent research while still working for Lookout Oil. In 1924 a 1:1 ratio Ni/Si alloy was produced, which after treatment with sodium hydroxide, was found to be five times more active than the best catalyst used in the hydrogenation of cottonseed oil. A patent for this discovery was issued in December 1925.<sup>[28]</sup>

Subsequently, Raney produced a 1:1 Ni/Al alloy following a procedure similar to the one used for the nickel-silicon catalyst. He found that the resulting catalyst was even more active and filed a patent application in 1926.<sup>[29]</sup> This is now a common alloy composition for modern Raney nickel catalysts.<sup>[2]</sup> Other common alloy compositions include 21:29 Ni/Al and 3:7 Ni/Al. Both the activity and preparation protocols for these catalysts vary.<sup>[2][30]</sup>

Following the development of Raney nickel, other alloy systems with aluminium were considered, of which the most notable include copper, ruthenium and cobalt.<sup>[31]</sup> Further research showed that adding a small amount of a third metal to the binary alloy would promote the activity of the catalyst. Some widely used promoters are zinc, molybdenum and chromium. An alternative way of preparing enantioselective Raney nickel has been devised by surface adsorption of tartaric acid.<sup>[32]</sup>

<div>NFPA 704</div> <div><i>fire diamond</i></div>			
1	3	1	

## See also

- Nickel aluminide
- Urushibara nickel
- Rieke nickel
- Nickel boride catalyst
- Raney cobalt, a similar cobalt/aluminum alloy catalyst which is sometimes more selective for certain hydrogenation products (e.g. primary amines via nitrile reduction).<sup>[2]</sup>

## References

- "Spongy Nickel" (http://www.spaceflight.esa.int/impress/text/education/Catalysis/Spongy.html). European Space Agency.
- Nishimura, Shigeo (2001). *Handbook of Heterogeneous Catalytic Hydrogenation for Organic Synthesis* (https://books.google.com/books?id=RjZRAAAAMAAJ&q=0471396982&dq=0471396982&hl=en&sa=X&ei=BCacVMTgN5LmoASd34KQCQ&ved=0CB8Q6AEwAA) (1st ed.). New York: Wiley-Interscience. pp. 7–19. ISBN 9780471396987.
- Billica, Harry; Adkins, Homer (1949). "Cataylst, Raney Nickel, W6 (with high contents of aluminum and adsorbed hydrogen)" (http://ww.w.orgsyn.org/demo.aspx?prep=CV3P0176). *Organic Syntheses*. **29**: 24. doi:10.15227/orgsyn.029.0024 (https://doi.org/10.15227%2Forgsyn.029.0024).; *Collective Volume*, **3**, p. 176



4. See:
- Raney, Murray, "Method of producing finely-divided nickel," (<http://patimg2.uspto.gov/.piw?Docid=01628190&homeurl=http%3A%2F%2Fpatft.uspto.gov%2Fnetacgi%2Fnph-Parser%3FSect1%3DPTO2%2526Sect2%3DHITOFF%2526u%3D%25252Fmeta.html%25252FPTO%25252Fsearch-adv.htm%2526r%3D456%2526f%3DG%2526l%3D50%2526d%3DPALL%2526s1%3DMurray%2526s2%3DRaney%2526co1%3DAND%2526p%3D10%2526OS%3DMurray%2526BAND%2526Raney%2526RS%3DMurray%2526BAND%2526>) U.S. patent 1,628,190 (filed: 14 May 1926 ; issued: 10 May 1927).
  - M. S. Wainwright, "3.2 Skeletal metal catalysts" in: Gerhard Ertl, Helmut Knözinger, and Jens Weitkamp, ed.s, *Preparaton of Solid Catalysts* (Weinheim, Federal Republic of Germany: Wiley-VCH Verlag, 1999), pages 28–29. (<https://books.google.com/books?id=ev47CMLmM2sC&pg=PA28>)
5. Yang, Teng-Kuei; Lee, Dong-Sheng; Haas, Julia (2005). "Raney Nickel". *Encyclopedia of Reagents for Organic Synthesis*. New York: John Wiley & Sons. doi:10.1002/047084289X.rr001.pub2 (<https://doi.org/10.1002%2F047084289X.rr001.pub2>).
6. "Specialty Catalysts & Process Technologies" (<https://web.archive.org/web/20091021081210/http://www.grace.com/SpecialtyCatalysts/default.aspx>). Grace company. Archived from the original (<http://www.grace.com/SpecialtyCatalysts/default.aspx>) on 21 October 2009.
7. Ertl, Gerhard; Knözinger, Helmut (1997). *Preparation of Solid Catalysts* (<https://books.google.com/books?id=Yb38DM1loAsC&pg=PA30>). Wiley. pp. 30–34. ISBN 3-527-29826-6.
8. Smith, A.J.; Trimm, D.L. (2005). "The preparation of skeletal catalysts". *Annu. Rev. Mater. Res.* **35**: 127. doi:10.1146/annurev.matsci.35.102303.140758 (<https://doi.org/10.1146%2Fannurev.matsci.35.102303.140758>).
9. M. Guisnet, ed. (1993). *Heterogeneous catalysis and fine chemicals III: proceedings of the 3rd international symposium* (<https://books.google.com/books?id=AwyJ6IEtaBsC&pg=PA69>). Elsevier. p. 69. ISBN 0-444-89063-7.
10. Crawford, Gerald (April 2003). "Exotic Alloy Finds Niche" ([http://www.nidi.org/index.cfm/ci\\_id/12317.htm](http://www.nidi.org/index.cfm/ci_id/12317.htm)). *Nickel magazine*. Retrieved 19 December 2006.
11. Carruthers, W (1986). *Some modern methods of organic synthesis* (<https://books.google.com/books?id=ti7yMYYW7CMC&pg=PA413>). Cambridge University Press. pp. 413–414. ISBN 0-521-31117-9.
12. Hauptmann, Heinrich; Walter, Wolfgang Ferdinand (1962). "The Action of Raney Nickel on Organic Sulfur Compounds". *Chem. Rev.* **62** (5): 347. doi:10.1021/cr60219a001 (<https://doi.org/10.1021%2Fcr60219a001>).
13. "Raney nickel usage in Organic Syntheses" (<https://web.archive.org/web/20090605024106/http://www.orgsyn.org/orgsyn/chemname.asp?nameID=33759>). 2005. Archived from the original (<http://www.orgsyn.org/orgsyn/chemname.asp?nameID=33759>) on 5 June 2009. Retrieved 1 August 2009.
14. Solomons, T.W. Graham; Fryhle, Craig B. (2004). *Organic Chemistry* ([https://archive.org/details/organicchemistry00solo\\_0](https://archive.org/details/organicchemistry00solo_0)). Wiley. ISBN 0-471-41799-8.
15. Jonathan Clayden; Nick Greeves; Stuart Warren (2012). *Organic Chemistry* (2 ed.). Oxford University Press. ISBN 9780199270293.
16. Graham, A. R.; Millidge, A. F.; Young, D. P. (1954). "Oxidation products of diisobutylene. Part III. Products from ring-opening of 1,2-epoxy-2,4,4-trimethylpentane". *Journal of the Chemical Society (Resumed)*: 2180. doi:10.1039/JR9540002180 (<https://doi.org/10.1039%2FJR9540002180>).
17. Gassman, P. G.; van Bergen, T. J. (1988). "Indoles from anilines: Ethyl 2-methylindole-5-carboxylate". *Organic Syntheses*. doi:10.15227/orgsyn.056.0072 (<https://doi.org/10.15227%2Forgsyn.056.0072>).; *Collective Volume*, **6**, p. 601
18. Hoegberg, Hans Erik; Hedenstroem, Erik; Faegerhag, Jonas; Servi, Stefano (1992). "Bakers' yeast reduction of thiophenepropaenals. Enantioselective synthesis of (S)-2-methyl-1-alkanols via bakers' yeast mediated reduction of 2-methyl-3-(2-thiophene)propenals". *J. Org. Chem.* **57** (7): 2052–2059. doi:10.1021/jo00033a028 (<https://doi.org/10.1021%2Fjo00033a028>).
19. Page, G. A.; Tarbell, D. S. (1963). "β-(o-Carboxyphenyl)propionic acid" (<http://www.orgsyn.org/demo.aspx?prep=CV4P0136>). *Organic Syntheses*. **34**: 8. doi:10.15227/orgsyn.034.0008 (<https://doi.org/10.15227%2Forgsyn.034.0008>).; *Collective Volume*, **4**, p. 136
20. Robinson, H. C.; Snyder, H. R. (1955). "β-Phenylethylamine" (<http://www.orgsyn.org/demo.aspx?prep=CV3P0720>). *Organic Syntheses*. **23**: 71. doi:10.15227/orgsyn.023.0071 (<https://doi.org/10.15227%2Forgsyn.023.0071>).; *Collective Volume*, **3**, p. 720
21. Schwenk, E.; Papa, D.; Hankin, H.; Ginsberg, H. (1955). "γ-n-Propylbutyrolactone and β-(Tetrahydrofuryl)propionic acid" (<http://www.orgsyn.org/demo.aspx?prep=CV3P0742>). *Organic Syntheses*. **27**: 68. doi:10.15227/orgsyn.027.0068 (<https://doi.org/10.15227%2Forgsyn.027.0068>).; *Collective Volume*, **3**, p. 742
22. Alexakis, Alex; Lensen, Nathalie; Mangeney, Pierre (1991). "Ultrasound-Assisted Cleavage of N-N Bonds in Hydrazines by Raney Nickel". *Synlett*. **1991** (9): 625–626. doi:10.1055/s-1991-20818 (<https://doi.org/10.1055%2Fs-1991-20818>).
23. Enders, D.; Pieter, R.; Renger, B.; Seebach, D. (1988). "Nucleophilic α-sec-aminoalkylation: 2-(diphenylhydroxymethyl)pyrrolidene" (<http://www.orgsyn.org/demo.aspx?prep=CV6P0542>). *Organic Syntheses*. **58**: 113. doi:10.15227/orgsyn.058.0113 (<https://doi.org/10.15227%2Forgsyn.058.0113>).; *Collective Volume*, **6**, p. 542
24. Rice, R. G.; Kohn, E. J. (1963). "*N,N'*-Diethylbenzidine" (<http://www.orgsyn.org/demo.aspx?prep=CV4P0283>). *Organic Syntheses*. **36**: 21. doi:10.15227/orgsyn.036.0021 (<https://doi.org/10.15227%2Forgsyn.036.0021>).; *Collective Volume*, **4**, p. 283
25. Armour, M.-A (2003). *Hazardous laboratory chemicals disposal guide* (<https://books.google.com/books?id=geK5gjEn3hkC&pg=PA331>). CRC Press. p. 331. ISBN 1-56670-567-3.
26. "Nickel aluminide MSDS" (<http://www.msds hazcom.com/MSDS/E/espi/Nickel%20Aluminide%20NiAl.pdf>) (PDF). Electronic Space Products International. 1994. Retrieved 7 July 2009.
27. P. Ryabchuk, G. Agostini, M.-M. Pohl, H. Lund, A. Agapova, H. Junge, K. Junge and M. Beller, Sci. Adv., 2018, 4, eaat0761, <https://advances.sciencemag.org/content/4/6/eaat0761>
28. US 1563587 (<https://worldwide.espacenet.com/textdoc?DB=EPODOC&IDX=US1563587>), Murray Raney, "Method of Preparing Catalytic Material", issued 1925-12-01 (Raney's original nickel-silicon catalyst)
29. US 1628190 (<https://worldwide.espacenet.com/textdoc?DB=EPODOC&IDX=US1628190>), Murray Raney, "Method of Producing Finely-Divided Nickel", issued 1927-05-10
30. Urushibara, Yoshiyuki; Nishimura, Shigeo (1957). "A Method for the Preparation of the Raney Nickel Catalyst with a Greater Activity" (<https://doi.org/10.1246%2Fbcsj.30.199>). *Bull. Chem. Soc. Jpn.* **30** (2): 199. doi:10.1246/bcsj.30.199 (<https://doi.org/10.1246%2Fbcsj.30.199>).
31. Augustine, Robert L. (1996). *Heterogeneous catalysis for the synthetic chemist* (<https://books.google.com/books?id=Nam8OVqKaXIC&pg=PA250>). CRC Press. pp. 248–249. ISBN 0-8247-9021-9.
32. Bakker, M. L.; Young, D. J.; Wainwright, M. S. (1988). "Selective leaching of NiAl<sub>3</sub> and Ni<sub>2</sub>Al<sub>3</sub> intermetallics to form Raney nickels". *J. Mater. Sci.* **23** (11): 3921–3926. doi:10.1007/BF01106814 (<https://doi.org/10.1007%2FBF01106814>).

## External links

- International Chemical Safety Card 0062 (<http://www.inchem.org/documents/icsc/icsc/eics0062.htm>)
- NIOSH Pocket Guide to Chemical Hazards (<https://www.cdc.gov/niosh/npg/npgd0445.html>)
- 1941 paper describing the preparation of W-2 grade Raney nickel: Mozingo, Ralph (1941). "Catalyst, Raney Nickel, W-2" (<http://www.orgsyn.org/demo.aspx?prep=CV3P0181>). *Organic Syntheses*. **21**: 15. doi:10.15227/orgsyn.021.0015 (<https://doi.org/10.15227%2Forgsyn.021.0015>).

Retrieved from "https://en.wikipedia.org/w/index.php?title=Raney\_nickel&oldid=1005520526"

**This page was last edited on 8 February 2021, at 02:48 (UTC).**

Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.