# **Organic synthesis**

**Organic synthesis** is a special branch of <u>chemical synthesis</u> and is concerned with the intentional construction of <u>organic compounds</u>. Organic <u>molecules</u> are often more complex than <u>inorganic compounds</u>, and their synthesis has developed into one of the most important branches of <u>organic chemistry</u>. There are several main areas of research within the general area of organic synthesis: <u>total synthesis</u>, <u>semisynthesis</u>, and <u>methodology</u>.

#### **Contents**

Total synthesis

Methodology and applications

Stereoselective synthesis

Synthesis design

See also

References

**Further reading** 

**External links** 

# **Total synthesis**

A total synthesis is the complete <u>chemical synthesis</u> of complex <u>organic molecules</u> from simple, commercially available <u>petrochemical or natural precursors. [2]</u> Total synthesis may be accomplished either via a linear or convergent approach. In a <u>linear synthesis</u>—often adequate for simple structures—several steps are performed one after another until the molecule is complete; the chemical compounds made in each step are called synthetic intermediates. [2] Most often, each step in a synthesis refers to a separate reaction taking place to modify the starting compound. For more complex molecules, a <u>convergent synthetic</u> approach may be preferable, one that involves individual preparation of several "pieces" (key intermediates), which are then combined to form the desired product. Convergent synthesis has the advantage of generating higher yield, compared to linear synthesis.

Robert Burns Woodward, who received the 1965 Nobel Prize for Chemistry for several total syntheses [3] (e.g., his 1954 synthesis of strychnine [4]), is regarded as the father of modern organic synthesis. Some latter-day examples include Wender's, [5] Holton's, [6] Nicolaou's, [7] and Danishefsky's [8] total syntheses of the anti-cancer therapeutic, paclitaxel (trade name,  $\underline{\text{Taxol}}$ ). [9]

# Methodology and applications

Each step of a synthesis involves a chemical reaction, and reagents and conditions for each of these reactions must be designed to give an adequate yield of pure product, with as few steps as possible. A method may already exist in the literature for making one of the early synthetic intermediates, and this method will usually be used rather than an effort to "reinvent the wheel". However, most intermediates are compounds that have never been made before, and these will

normally be made using general methods developed by methodology researchers. To be useful, these methods need to give high <u>yields</u>, and to be reliable for a broad range of <u>substrates</u>. For practical applications, additional hurdles include industrial standards of safety and purity. [11]

Methodology research usually involves three main stages: <u>discovery</u>, <u>optimisation</u>, and studies of scope and limitations. The discovery requires extensive knowledge of and experience with chemical reactivities of appropriate reagents. *Optimisation* is a process in which one or two starting compounds are tested in the reaction under a wide variety of conditions of <u>temperature</u>, <u>solvent</u>, <u>reaction time</u>, etc., until the optimal conditions for product yield and purity are found. Finally, the researcher tries to extend the method to a broad range of different starting materials, to find the scope and limitations. Total syntheses (see above) are sometimes used to showcase the new methodology and demonstrate its value in a real-world application. Such applications involve major industries focused especially on polymers (and plastics) and pharmaceuticals. Some syntheses are feasible on a research or academic level, but not for industry level production. This may lead to further modification of the process. [13]

# Stereoselective synthesis

Most complex <u>natural products</u> are chiral, <u>[14][15]</u> and the bioactivity of chiral molecules varies with the <u>enantiomer</u>. <u>[16]</u> Historically, total syntheses targeted <u>racemic</u> mixtures, mixtures of both possible enantiomers, after which the racemic mixture might then be separated via chiral resolution.

In the later half of the twentieth century, chemists began to develop methods of stereoselective catalysis and kinetic resolution whereby reactions could be directed to produce only one enantiomer rather than a racemic mixture. Early examples include stereoselective hydrogenations (e.g., as reported by William Knowles<sup>[17]</sup> and Ryōji Noyori, and functional group modifications such as the asymmetric epoxidation of Barry Sharpless; for these specific achievements, these workers were awarded the Nobel Prize in Chemistry in 2001. Such reactions gave chemists a much wider choice of enantiomerically pure molecules to start from, where previously only natural starting materials could be used. Using techniques pioneered by Robert B. Woodward and new developments in synthetic methodology, chemists became more able to take simple molecules through to more complex molecules without unwanted racemisation, by understanding stereocontrol, allowing final target molecules to be synthesised as pure enantiomers (i.e., without need for resolution). Such techniques are referred to as *stereoselective synthesis*.

## Synthesis design

Elias James Corey brought a more formal approach to synthesis design, based on retrosynthetic analysis, for which he won the Nobel Prize for Chemistry in 1990. In this approach, the synthesis is planned backwards from the product, using standard rules. [21] The steps "breaking down" the parent structure into achievable component parts are shown in a graphical scheme that uses retrosynthetic arrows (drawn as  $\Rightarrow$ , which in effect, mean "is made from").

More recently, and less widely accepted, computer programs have been written for designing a synthesis based on sequences of generic "half-reactions". [22]

### See also

- Organic Syntheses (journal)
- Methods in Organic Synthesis (journal)
- Electrosynthesis
- Automated Synthesis

#### References

- 1. Cornforth, JW (1993-02-01). <u>"The Trouble With Synthesis"</u> (https://doi.org/10.1071%2Fch993015 7). *Australian Journal of Chemistry*. **46** (2): 157–170. <u>doi</u>:10.1071/ch9930157 (https://doi.org/10.1071%2Fch9930157).
- 2. Nicolaou, K. C.; Sorensen, E. J. (1996). *Classics in Total Synthesis* (https://archive.org/details/classicsintotals0000nico\_p5l6). New York: VCH.
- 3. "Nobelprize.org" (https://www.nobelprize.org/nobel\_prizes/chemistry/laureates/1965/press.html). www.nobelprize.org. Retrieved 2016-11-20.
- 4. Woodward, R. B.; Cava, M. P.; Ollis, W. D.; Hunger, A.; Daeniker, H. U.; Schenker, K. (1954). "The Total Synthesis of Strychnine". *Journal of the American Chemical Society*. **76** (18): 4749–4751. doi:10.1021/ja01647a088 (https://doi.org/10.1021%2Fja01647a088).
- Wender, Paul A.; Badham, Neil F.; Conway, Simon P.; Floreancig, Paul E.; Glass, Timothy E.; Gränicher, Christian; Houze, Jonathan B.; Jänichen, Jan; Lee, Daesung (1997-03-01). "The Pinene Path to Taxanes. 5. Stereocontrolled Synthesis of a Versatile Taxane Precursor". *Journal of the American Chemical Society*. 119 (11): 2755–2756. doi:10.1021/ja9635387 (https://doi.org/10.1021%2Fja9635387). ISSN 0002-7863 (https://www.worldcat.org/issn/0002-7863).
- Holton, Robert A.; Somoza, Carmen; Kim, Hyeong Baik; Liang, Feng; Biediger, Ronald J.; Boatman, P. Douglas; Shindo, Mitsuru; Smith, Chase C.; Kim, Soekchan (1994-02-01). "First total synthesis of taxol. 1. Functionalization of the B ring". *Journal of the American Chemical Society*. 116 (4): 1597–1598. doi:10.1021/ja00083a066 (https://doi.org/10.1021%2Fja00083a066). ISSN 0002-7863 (https://www.worldcat.org/issn/0002-7863).
- 7. Nicolaou, K. C.; Yang, Z.; Liu, J. J.; Ueno, H.; Nantermet, P. G.; Guy, R. K.; Claiborne, C. F.; Renaud, J.; Couladouros, E. A. (1994-02-17). "Total synthesis of taxol". *Nature*. **367** (6464): 630–634. Bibcode: 1994Natur.367..630N (https://ui.adsabs.harvard.edu/abs/1994Natur.367..630N). doi:10.1038/367630a0 (https://doi.org/10.1038%2F367630a0). PMID 7906395 (https://pubmed.ncbi.nlm.nih.gov/7906395).
- Danishefsky, Samuel J.; Masters, John J.; Young, Wendy B.; Link, J. T.; Snyder, Lawrence B.; Magee, Thomas V.; Jung, David K.; Isaacs, Richard C. A.; Bornmann, William G. (1996-01-01). "Total Synthesis of Baccatin III and Taxol". *Journal of the American Chemical Society.* 118 (12): 2843–2859. doi:10.1021/ja952692a (https://doi.org/10.1021%2Fja952692a). ISSN 0002-7863 (https://www.worldcat.org/issn/0002-7863).
- 9. "Taxol The Drama behind Total Synthesis" (https://web.archive.org/web/20110727152818/http://www.org-chem.org/yuuki/taxol/taxol\_en.html). www.org-chem.org. Archived from the original (http://www.org-chem.org/yuuki/taxol/taxol\_en.html) on 2011-07-27. Retrieved 2016-11-20.
- 10. March, J.; Smith, D. (2001). Advanced Organic Chemistry, 5th ed. New York: Wiley.
- 11. Carey, J.S.; Laffan, D.; Thomson, C. & Williams, M.T. (2006). "Analysis of the reactions used for the preparation of drug candidate molecules" (https://semanticscholar.org/paper/5b27f82f2e7bd7 365491988eb063e78b250e0b1d). Org. Biomol. Chem. 4 (12): 2337–2347. doi:10.1039/B602413K (https://doi.org/10.1039%2FB602413K). PMID 16763676 (https://pubmed.ncbi.nlm.nih.gov/16763676).
- Nicolaou, K. C.; Hale, Christopher R. H.; Nilewski, Christian; Ioannidou, Heraklidia A. (2012-07-09). "Constructing molecular complexity and diversity: total synthesis of natural products of biological and medicinal importance" (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3426871). Chemical Society Reviews. 41 (15): 5185–5238. doi:10.1039/C2CS35116A (https://doi.org/10.1039/C2CS35116A). ISSN 1460-4744 (https://www.worldcat.org/issn/1460-4744). PMC 3426871 (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3426871). PMID 22743704 (https://pubmed.ncbi.nlm.nih.gov/22743704).
- Chen, Weiming; Suo, Jin; Liu, Yongjian; Xie, Yuanchao; Wu, Mingjun; Zhu, Fuqiang; Nian, Yifeng; Aisa, Haji A.; Shen, Jingshan (2019-03-08). "Industry-Oriented Route Evaluation and Process Optimization for the Preparation of Brexpiprazole". *Organic Process Research & Development*.
  (5): 852–857. doi:10.1021/acs.oprd.8b00438 (https://doi.org/10.1021%2Facs.oprd.8b00438). ISSN 1083-6160 (https://www.worldcat.org/issn/1083-6160).

- 14. Blackmond, Donna G. (2016-11-20). "The Origin of Biological Homochirality" (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2857173). Cold Spring Harbor Perspectives in Biology. 2 (5): a002147. doi:10.1101/cshperspect.a002147 (https://doi.org/10.1101%2Fcshperspect.a002147). ISSN 1943-0264 (https://www.worldcat.org/issn/1943-0264). PMC 2857173 (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2857173). PMID 20452962 (https://pubmed.ncbi.nlm.nih.gov/20452962).
- 15. Welch, CJ (1995). Advances in Chromatography. New York: Marcel Dekker, Inc. p. 172.
- 16. Nguyen, Lien Ai; He, Hua; Pham-Huy, Chuong (2016-11-20). "Chiral Drugs: An Overview" (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3614593). International Journal of Biomedical Science. 2 (2): 85–100. ISSN 1550-9702 (https://www.worldcat.org/issn/1550-9702). PMC 3614593 (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3614593). PMID 23674971 (https://pubmed.ncbi.nlm.nih.gov/23674971).
- 17. Knowles, William S. (2002-06-17). "Asymmetric Hydrogenations (Nobel Lecture)". *Angewandte Chemie International Edition*. **41** (12): 1998–2007. <a href="doi:10.1002/1521-3773(20020617)41:12<1998::AID-ANIE1998>3.0.CO;2-8 (https://doi.org/10.1002%2F1521-3773%2820020617%2941%3A12%3C1998%3A%3AAID-ANIE1998%3E3.0.CO%3B2-8). ISSN 1521-3773 (https://www.worldcat.org/issn/1521-3773).
- 18. Noyori, R.; Ikeda, T.; Ohkuma, T.; Widhalm, M.; Kitamura, M.; Takaya, H.; Akutagawa, S.; Sayo, N.; Saito, T. (1989). "Stereoselective hydrogenation via dynamic kinetic resolution". *Journal of the American Chemical Society.* **111** (25): 9134–9135. <a href="https://doi.org/10.1021/ja00207a038">doi:10.1021/ja00207a038</a> (https://doi.org/10.1021%2Fja00207a038).
- 19. Gao, Yun; Klunder, Janice M.; Hanson, Robert M.; Masamune, Hiroko; Ko, Soo Y.; Sharpless, K. Barry (1987-09-01). "Catalytic asymmetric epoxidation and kinetic resolution: modified procedures including in situ derivatization". *Journal of the American Chemical Society*. **109** (19): 5765–5780. doi:10.1021/ja00253a032 (https://doi.org/10.1021%2Fja00253a032). ISSN 0002-7863 (https://www.worldcat.org/issn/0002-7863).
- 20. Service. R.F. (2001). "Science Awards Pack a Full House of Winners". *Science*. **294** (5542, October 19): 503–505. doi:10.1126/science.294.5542.503b (https://doi.org/10.1126%2Fscience.294.5542.503b). PMID 11641480 (https://pubmed.ncbi.nlm.nih.gov/11641480).
- 21. Corey, E. J.; Cheng, X-M. (1995). The Logic of Chemical Synthesis. New York: Wiley.
- 22. Todd, Matthew H. (2005). "Computer-aided Organic Synthesis" (https://semanticscholar.org/paper/28a699880bce80839953ac07c1d1838822a36120). *Chemical Society Reviews*. **34** (3): 247–266. doi:10.1039/b104620a (https://doi.org/10.1039%2Fb104620a). PMID 15726161 (https://pubmed.ncbi.nlm.nih.gov/15726161).

### **Further reading**

 Corey EJ; Cheng, X-M (1995). The Logic of Chemical Synthesis. New York, NY: John Wiley & Sons. ISBN 978-0471115946.

## **External links**

- The Organic Synthesis Archive (http://www.synarchive.com)
- Chemical synthesis database (http://www.chemsynthesis.com)
- https://web.archive.org/web/20070927231356/http://www.webreactions.net/search.html
- https://www.organic-chemistry.org/synthesis/
- Prof. Hans Reich's collection of (http://www.chem.wisc.edu/areas/reich/syntheses/syntheses.htm)natural product syntheses
- Chemical synthesis semantic wiki (https://web.archive.org/web/20141222223356/http://www.wikis ynth.net/)

This page was last edited on 16 July 2021, at 02:35 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.