

Direct Proofs

Two Quick Definitions

- An integer n is **even** if there is some integer k such that $n = 2k$.
 - This means that 0 is even.
- An integer n is **odd** if there is some integer k such that $n = 2k + 1$.
 - This means that 0 is not odd.
- We'll assume the following for now:
 - Every integer is either even or odd.
 - No integer is both even and odd.

Our First Direct Proof

Theorem: If n is an even integer, then n^2 is even.

Proof: Let n be an even integer.

Since n is even, there is some integer k such that $n = 2k$.

This means that $n^2 = (2k)^2 = 4k^2 = 2(2k^2)$.

From this, we see that there is an integer m (namely, $2k^2$) where $n^2 = 2m$.

Therefore, n^2 is even. ■

Our First Direct Proof

Theorem: If n is an even integer, then n^2 is even.


Proof: Let n be an even integer.

Since n is even, there is some integer k such that $n = 2k$.

This means that $n^2 = (2k)^2 = 4k^2 = 2(2k^2)$.

From this, we see that there is an integer m (namely, $2k^2$) where $n^2 = 2m$.

Therefore, n^2 is even. 



This symbol
means "end of
proof"

Our First Direct Proof

Theorem: If n is an even integer, then n^2 is even.

Proof: Let n be an even integer.

Since n is even,
such that

This means

From this
 m (name

Therefore

To prove a statement of the
form

“If P , then Q ”

Assume that **P** is true, then show
that **Q** must be true as well.

Our First Direct Proof

Theorem: If n is an even integer, then n^2 is even.

Proof: Let n be an even integer.

Since n is even, there is some integer k such that $n = 2k$.

This means that

From this, we can write n as $2k$ (namely, $2k$).

Therefore, $n^2 = (2k)^2 = 4k^2$.

This is the definition of an even integer. When writing a mathematical proof, it's common to call back to the definitions.

Our First Direct Proof

Theorem: If n is an even integer, then n^2 is even.

Proof: Let n be an even integer.

Since n is even, there is some integer k such that $n = 2k$.

This means that $n^2 = (2k)^2 = 4k^2 = 2(2k^2)$.

From
m
Th Notice how we use the value of k that we obtained above. Giving names to quantities, even if we aren't fully sure what they are, allows us to manipulate them. This is similar to variables in programs.

Our First Direct Proof

Theorem: If n is even, then n^2 is even.

Proof: Let n be an even integer.

Since n is even,
such that

This means that $n = 2k$ for some integer k .
($n = 2k \implies n^2 = (2k)^2 = 4k^2 = 2(2k^2)$).

From this, we see that there is an integer m (namely, $2k^2$) where $n^2 = 2m$.

Therefore, n^2 is even. ■

Our ultimate goal is to prove that n^2 is even. This means that we need to find some m such that $n^2 = 2m$. Here, we're explicitly showing how we can do that.

Our First Direct Proof

Theorem: If n is an even integer, then n^2 is even.

Proof: Let n be an even integer.

Since n is even, there is some integer k such that $n = 2k$.

This means

From this we get
 m (name

Hey, that's what we were trying to show! We're done now.

Therefore, n^2 is even. ■

Our First Direct Proof

Theorem: If n is an even integer, then n^2 is even.

Proof: Let n be an even integer.


Since n is even, there is some integer k such that $n = 2k$.

This means that $n^2 = (2k)^2 = 4k^2 = 2(2k^2)$.


From this, we see that there is an integer m (namely, $2k^2$) where $n^2 = 2m$.

Therefore, n^2 is even. ■

A Visual Intuition




A Visual Intuition




The diagram shows a 10x10 grid. The top row is divided into two groups of 5 columns each. The first group of 5 columns is colored orange and is labeled with a blue k above it, with a brace spanning the 5 columns. The second group of 5 columns is colored dark blue and is also labeled with a blue k above it, with a brace spanning the 5 columns. The remaining 8 rows of the grid are colored light blue.

A Visual Intuition





A Visual Intuition



That wasn't so bad! Let's do another one.



Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.



Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.



Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.



Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof:

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof:

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof:

How do we prove
that this is true for
any integers?

Proving Something Always Holds

- Many statements have the form

For any x , [some-property] holds of x .

- Examples:

For all integers n , if n is even, n^2 is even.

For any sets A , B , and C , if $A \subseteq B$ and $B \subseteq C$, then $A \subseteq C$.

For all sets S , $|S| < |\wp(S)|$.

Everything that drowns me makes me wanna fly.

- How do we prove these statements when there are (potentially) infinitely many cases to check?

Arbitrary Choices

- To prove that some property holds true for all possible x , show that no matter what choice of x you make, that property must be true.
- Start the proof by choosing x *arbitrarily*:
 - “Let x be an arbitrary even integer.”
 - “Let x be any set containing 137.”
 - “Consider any x .”
 - “Pick an odd integer x .”
- Demonstrate that the property holds true for this choice of x .

ar·bi·trar·y

adjective /'ärbi,trerē/

...not this
one!

1. Based on random choice or personal whim, rather than any reason or system - *“his mealtimes were entirely arbitrary”*

2. (of power or a ruling body) Unrestrained and autocratic in the use of authority - *“arbitrary rule by King and bishops has been made impossible”*

3. (of a constant or other quantity) Of unspecified value

Use this
definition...

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof:

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd.

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd.

By picking m and n arbitrarily, anything we prove about m and n will generalize to all possible choices we could have made.

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd.

To prove a statement of the form

“If P , then Q ”

Assume that **P** is true, then show that **Q** must be true as well.

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd.

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd. Since m is odd, we know that there is an integer k where

$$m = 2k + 1. \quad (1)$$

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd. Since m is odd, we know that there is an integer k where

$$m = 2k + 1. \quad (1)$$

Similarly, because n is odd there must be some integer r such that

$$n = 2r + 1. \quad (2)$$

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd. Since m is odd, we know that there is an integer k where

$$m = 2k + 1. \quad (1)$$

Similarly, because n is odd there must be some integer r such that

$$n = 2r + 1. \quad (2)$$

By adding equations (1) and (2) we learn that

$$m + n = 2k + 1 + 2r + 1$$

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd. Since m is odd, we know that there is an integer k where

$$m = 2k + 1. \quad (1)$$

Similarly, because n is odd there must be some integer r such that

$$n = 2r + 1. \quad (2)$$

By adding equations (1) and (2) we learn that

$$\begin{aligned} m + n &= 2k + 1 + 2r + 1 \\ &= 2k + 2r + 2 \end{aligned}$$

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd. Since m is odd, we know that there is an integer k where

$$m = 2k + 1. \quad (1)$$

Similarly, because n is odd there must be some integer r such that

$$n = 2r + 1. \quad (2)$$

By adding equations (1) and (2) we learn that

$$\begin{aligned} m + n &= 2k + 1 + 2r + 1 \\ &= 2k + 2r + 2 \\ &= 2(k + r + 1). \end{aligned} \quad (3)$$

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd. Since m is odd, we know that there is an integer k where

$$m = 2k + 1. \quad (1)$$

Similarly, because n is odd there must be some integer r such that

$$n = 2r + 1. \quad (2)$$

By adding equations (1) and (2) we learn that

$$\begin{aligned} m + n &= 2k + 1 + 2r + 1 \\ &= 2k + 2r + 2 \\ &= 2(k + r + 1). \end{aligned} \quad (3)$$

Equation (3) tells us that there is an integer s (namely, $k + r + 1$) such that $m + n = 2s$.

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd. Since m is odd, we know that there is an integer k where

$$m = 2k + 1. \quad (1)$$

Similarly, because n is odd there must be some integer r such that

$$n = 2r + 1. \quad (2)$$

By adding equations (1) and (2) we learn that

$$\begin{aligned} m + n &= 2k + 1 + 2r + 1 \\ &= 2k + 2r + 2 \\ &= 2(k + r + 1). \end{aligned} \quad (3)$$

Equation (3) tells us that there is an integer s (namely, $k + r + 1$) such that $m + n = 2s$. Therefore, we see that $m + n$ is even, as required.

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd. Since m is odd, we know that there is an integer k where

$$m = 2k + 1. \quad (1)$$

Similarly, because n is odd there must be some integer r such that

$$n = 2r + 1. \quad (2)$$

By adding equations (1) and (2) we learn that

$$\begin{aligned} m + n &= 2k + 1 + 2r + 1 \\ &= 2k + 2r + 2 \\ &= 2(k + r + 1). \end{aligned} \quad (3)$$

Equation (3) tells us that there is an integer s (namely, $k + r + 1$) such that $m + n = 2s$. Therefore, we see that $m + n$ is even, as required. ■

Numbering these equalities lets us refer back to them later on, making the flow of the proof a bit easier to understand.

$$m = 2k + 1. \quad (1)$$

Similarly, because n is odd there must be some integer r such that

$$n = 2r + 1. \quad (2)$$

By adding equations (1) and (2) we learn that

$$\begin{aligned} m + n &= 2k + 1 + 2r + 1 \\ &= 2k + 2r + 2 \\ &= 2(k + r + 1). \end{aligned} \quad (3)$$

Equation (3) tells us that there is an integer s (namely, $k + r + 1$) such that $m + n = 2s$. Therefore, we see that $m + n$ is even, as required. ■

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd. Since m is odd, we know that there is an integer k where

$$m = 2k + 1. \quad (1)$$

Similarly, because n is odd there must be some integer r such that

$$n = 2r + 1. \quad (2)$$

By adding equations (1) and (2) we learn that

Notice that we use k in the first equality and r in the second equality. That's because we know that n is twice something plus one, but we can't say for sure that it's k specifically.

Equation (3) tells us that $m + n = 2(k + r + 1)$ such that $m + n$ is even, as required. ■

$k + r + 1$)
even, as

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd. Since m is odd, we know that there is an integer k where


$$m = 2k + 1. \quad (1)$$

Similarly, because n is odd there must be some integer r such that

$$n = 2r + 1. \quad (2)$$

This is a grammatically correct and complete sentence! Proofs are expected to be written in complete sentences, so you'll often use punctuation at the end of formulas.

We recommend using the "mugga mugga" test – if you read a proof and replace all the mathematical notation with "mugga mugga," what comes back should be a valid sentence.



Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd. Since m is odd, we know that there is an integer k where

$$m = 2k + 1. \quad (1)$$

Similarly, because n is odd there must be some integer r such that

$$n = 2r + 1. \quad (2)$$

By adding equations (1) and (2) we learn that

$$\begin{aligned} m + n &= 2k + 1 + 2r + 1 \\ &= 2k + 2r + 2 \\ &= 2(k + r + 1). \end{aligned} \quad (3)$$

Equation (3) tells us that there is an integer s (namely, $k + r + 1$) such that $m + n = 2s$. Therefore, we see that $m + n$ is even, as required. ■

Theorem: For any integers m and n , if m and n are odd, then $m + n$ is even.

Proof: Consider any arbitrary integers m and n where m and n are odd. Since m is odd, we know that there is an integer k where

$$m = 2k + 1. \quad (1)$$

Similarly, because n is odd there must be some integer r such that

$$n = 2r + 1.$$

By adding equations (1) and (2) we learn that

$$\begin{aligned} m + n &= 2k + 1 + 2r + 1 \\ &= 2k + 2r + 2 \\ &= 2(k + r + 1). \end{aligned}$$

Trace through this proof if $m = 7$ and $n = 9$. What is the resulting value of s ?

- A. 3
- B. 8
- C. 17

Equation (3) tells us that there is an integer s (namely, $k + r + 1$) such that $m + n = 2s$. Therefore, we see that $m + n$ is even, as required. ■

Answer at **PollEv.com/cs103** or
text **CS103** to **22333** once to join, then **A**, **B**, or **C**.