

# Cardinality

New Stuff!

# Inverse Functions



**Katniss  
Everdeen**



**Elsa**



**Hermione  
Granger**

# Inverse Functions

- In some cases, it's possible to “turn a function around.”
- Let  $f : A \rightarrow B$  be a function. A function  $f^{-1} : B \rightarrow A$  is called an **inverse of  $f$**  if the following first-order logic statements are true about  $f$  and  $f^{-1}$

$$\forall a \in A. (f^{-1}(f(a)) = a) \qquad \forall b \in B. (f(f^{-1}(b)) = b)$$

- In other words, if  $f$  maps  $a$  to  $b$ , then  $f^{-1}$  maps  $b$  back to  $a$  and vice-versa.
- Not all functions have inverses (we just saw a few examples of functions with no inverses).
- If  $f$  is a function that has an inverse, then we say that  $f$  is **invertible**.

# Inverse Functions

- ***Theorem:*** Let  $f : A \rightarrow B$ . Then  $f$  is invertible if and only if  $f$  is a bijection.
- This proof is in the course reader. Feel free to check it out if you'd like!
- ***Really cool observation:*** Look at the formal definition of a function. Look at the rules for injectivity and surjectivity. Do you see why this result makes sense?

# Where We Are

- We now know
  - what an injection, surjection, and bijection are;
  - that the composition of two injections, surjections, or bijections is also an injection, surjection, or bijection, respectively; and
  - that bijections are invertible and invertible functions are bijections.
- You might wonder why this all matters. Well, there's a good reason...

# Cardinality Revisited



# Cardinality

- Recall (*from our first lecture!*) that the **cardinality** of a set is the number of elements it contains.
- If  $S$  is a set, we denote its cardinality by  $|S|$ .
- For finite sets, cardinalities are natural numbers:
  - $|\{1, 2, 3\}| = 3$
  - $|\{100, 200\}| = 2$
- For infinite sets, we introduced **infinite cardinals** to denote the size of sets:

$$|\mathbb{N}| = \aleph_0$$


# Defining Cardinality

- It is difficult to give a rigorous definition of what cardinalities actually are.
  - What is 4? What is  $\aleph_0$ ?
  - (Take Math 161 for an answer!)
- ***Idea:*** Define cardinality as a *relation* between two sets rather than an absolute quantity.

# Comparing Cardinalities

- Here is the formal definition of what it means for two sets to have the same cardinality:


**$|S| = |T|$  if there exists a *bijection*  $f : S \rightarrow T$**



# Comparing Cardinalities

- Here is the formal definition of what it means for two sets to have the same cardinality:

**$|S| = |T|$  if there exists a *bijection*  $f : S \rightarrow T$**



# Fun with Cardinality

# Terminology Refresher

- Let  $a$  and  $b$  be real numbers where  $a \leq b$ .
- The notation  $[a, b]$  denotes the set of all real numbers between  $a$  and  $b$ , inclusive.

$$[a, b] = \{ x \in \mathbb{R} \mid a \leq x \leq b \}$$


- The notation  $(a, b)$  denotes the set of all real numbers between  $a$  and  $b$ , exclusive.

$$(a, b) = \{ x \in \mathbb{R} \mid a < x < b \}$$

# Home on the Range



# Home on the Range



$$f: [0, 1] \rightarrow [0, 2]$$
$$f(x) = 2x$$



***Theorem:***  $|[0, 1]| = |[0, 2]|$

***Theorem:***  $|[0, 1]| = |[0, 2]|$

***Proof:***

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ .

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ .  
We will prove that  $f$  is a bijection.

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ .  
We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function.

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ .  
We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ .

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ .  
We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ .

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ .  
We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .



**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ .  
We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .

Next, we'll show that  $f$  is injective.

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ .  
We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in [0, 1]$  where  $f(x_1) = f(x_2)$ .

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ . We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in [0, 1]$  where  $f(x_1) = f(x_2)$ . We will show that  $x_1 = x_2$ .

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ . We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in [0, 1]$  where  $f(x_1) = f(x_2)$ . We will show that  $x_1 = x_2$ . To see this, notice that since  $f(x_1) = f(x_2)$ , we see that  $2x_1 = 2x_2$ , which in turn tells us that  $x_1 = x_2$ , as required.

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ .

How many of the following are proper ways of setting up the next part of this proof?

Choose any  $x \in [0, 1]$ . We will show there is a  $y \in [0, 2]$  such that  $f(x) = y$ .

Pick any  $y \in [0, 2]$ . We will show there is an  $x \in [0, 1]$  where  $f(x) = y$ .

Assume for the sake of contradiction that, for any  $y \in [0, 2]$  and for any  $x \in [0, 1]$ , we have  $f(x) \neq y$ .

Finally, we will show that  $f$  is surjective.

Answer at **PollEv.com/cs103** or  
text **CS103** to **22333** once to join, then a number between **0** and **3**.

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ . We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in [0, 1]$  where  $f(x_1) = f(x_2)$ . We will show that  $x_1 = x_2$ . To see this, notice that since  $f(x_1) = f(x_2)$ , we see that  $2x_1 = 2x_2$ , which in turn tells us that  $x_1 = x_2$ , as required.

Finally, we will show that  $f$  is surjective. To do so, consider any  $y \in [0, 2]$ .

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ . We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in [0, 1]$  where  $f(x_1) = f(x_2)$ . We will show that  $x_1 = x_2$ . To see this, notice that since  $f(x_1) = f(x_2)$ , we see that  $2x_1 = 2x_2$ , which in turn tells us that  $x_1 = x_2$ , as required.

Finally, we will show that  $f$  is surjective. To do so, consider any  $y \in [0, 2]$ . We'll show that there is some  $x \in [0, 1]$  where  $f(x) = y$ .

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ . We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in [0, 1]$  where  $f(x_1) = f(x_2)$ . We will show that  $x_1 = x_2$ . To see this, notice that since  $f(x_1) = f(x_2)$ , we see that  $2x_1 = 2x_2$ , which in turn tells us that  $x_1 = x_2$ , as required.

Finally, we will show that  $f$  is surjective. To do so, consider any  $y \in [0, 2]$ . We'll show that there is some  $x \in [0, 1]$  where  $f(x) = y$ .

Let  $x = y/2$ .



**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ . We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in [0, 1]$  where  $f(x_1) = f(x_2)$ . We will show that  $x_1 = x_2$ . To see this, notice that since  $f(x_1) = f(x_2)$ , we see that  $2x_1 = 2x_2$ , which in turn tells us that  $x_1 = x_2$ , as required.

Finally, we will show that  $f$  is surjective. To do so, consider any  $y \in [0, 2]$ . We'll show that there is some  $x \in [0, 1]$  where  $f(x) = y$ .

Let  $x = y/2$ . Since  $y \in [0, 2]$ , we know  $0 \leq y \leq 2$ , and therefore that  $0 \leq y/2 \leq 1$ .

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ . We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in [0, 1]$  where  $f(x_1) = f(x_2)$ . We will show that  $x_1 = x_2$ . To see this, notice that since  $f(x_1) = f(x_2)$ , we see that  $2x_1 = 2x_2$ , which in turn tells us that  $x_1 = x_2$ , as required.

Finally, we will show that  $f$  is surjective. To do so, consider any  $y \in [0, 2]$ . We'll show that there is some  $x \in [0, 1]$  where  $f(x) = y$ .

Let  $x = y/2$ . Since  $y \in [0, 2]$ , we know  $0 \leq y \leq 2$ , and therefore that  $0 \leq y/2 \leq 1$ . We picked  $x = y/2$ , so we know that  $0 \leq x \leq 1$ , which in turn means  $x \in [0, 1]$ .

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ . We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in [0, 1]$  where  $f(x_1) = f(x_2)$ . We will show that  $x_1 = x_2$ . To see this, notice that since  $f(x_1) = f(x_2)$ , we see that  $2x_1 = 2x_2$ , which in turn tells us that  $x_1 = x_2$ , as required.

Finally, we will show that  $f$  is surjective. To do so, consider any  $y \in [0, 2]$ . We'll show that there is some  $x \in [0, 1]$  where  $f(x) = y$ .

Let  $x = y/2$ . Since  $y \in [0, 2]$ , we know  $0 \leq y \leq 2$ , and therefore that  $0 \leq y/2 \leq 1$ . We picked  $x = y/2$ , so we know that  $0 \leq x \leq 1$ , which in turn means  $x \in [0, 1]$ . Moreover, notice that

$$f(x) = 2x$$

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ . We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in [0, 1]$  where  $f(x_1) = f(x_2)$ . We will show that  $x_1 = x_2$ . To see this, notice that since  $f(x_1) = f(x_2)$ , we see that  $2x_1 = 2x_2$ , which in turn tells us that  $x_1 = x_2$ , as required.

Finally, we will show that  $f$  is surjective. To do so, consider any  $y \in [0, 2]$ . We'll show that there is some  $x \in [0, 1]$  where  $f(x) = y$ .

Let  $x = y/2$ . Since  $y \in [0, 2]$ , we know  $0 \leq y \leq 2$ , and therefore that  $0 \leq y/2 \leq 1$ . We picked  $x = y/2$ , so we know that  $0 \leq x \leq 1$ , which in turn means  $x \in [0, 1]$ . Moreover, notice that

$$f(x) = 2x = 2(y/2)$$

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ . We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in [0, 1]$  where  $f(x_1) = f(x_2)$ . We will show that  $x_1 = x_2$ . To see this, notice that since  $f(x_1) = f(x_2)$ , we see that  $2x_1 = 2x_2$ , which in turn tells us that  $x_1 = x_2$ , as required.

Finally, we will show that  $f$  is surjective. To do so, consider any  $y \in [0, 2]$ . We'll show that there is some  $x \in [0, 1]$  where  $f(x) = y$ .

Let  $x = y/2$ . Since  $y \in [0, 2]$ , we know  $0 \leq y \leq 2$ , and therefore that  $0 \leq y/2 \leq 1$ . We picked  $x = y/2$ , so we know that  $0 \leq x \leq 1$ , which in turn means  $x \in [0, 1]$ . Moreover, notice that

$$f(x) = 2x = 2(y/2) = y$$

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ . We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in [0, 1]$  where  $f(x_1) = f(x_2)$ . We will show that  $x_1 = x_2$ . To see this, notice that since  $f(x_1) = f(x_2)$ , we see that  $2x_1 = 2x_2$ , which in turn tells us that  $x_1 = x_2$ , as required.

Finally, we will show that  $f$  is surjective. To do so, consider any  $y \in [0, 2]$ . We'll show that there is some  $x \in [0, 1]$  where  $f(x) = y$ .

Let  $x = y/2$ . Since  $y \in [0, 2]$ , we know  $0 \leq y \leq 2$ , and therefore that  $0 \leq y/2 \leq 1$ . We picked  $x = y/2$ , so we know that  $0 \leq x \leq 1$ , which in turn means  $x \in [0, 1]$ . Moreover, notice that

$$f(x) = 2x = 2(y/2) = y,$$

so  $f(x) = y$ , as required.

**Theorem:**  $|[0, 1]| = |[0, 2]|$

**Proof:** Consider the function  $f : [0, 1] \rightarrow [0, 2]$  defined as  $f(x) = 2x$ . We will prove that  $f$  is a bijection.

First, we will show that  $f$  is a well-defined function. Choose any  $x \in [0, 1]$ . This means that  $0 \leq x \leq 1$ , so we know that  $0 \leq 2x \leq 2$ . Consequently, we see that  $0 \leq f(x) \leq 2$ , so  $f(x) \in [0, 2]$ .

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in [0, 1]$  where  $f(x_1) = f(x_2)$ . We will show that  $x_1 = x_2$ . To see this, notice that since  $f(x_1) = f(x_2)$ , we see that  $2x_1 = 2x_2$ , which in turn tells us that  $x_1 = x_2$ , as required.

Finally, we will show that  $f$  is surjective. To do so, consider any  $y \in [0, 2]$ . We'll show that there is some  $x \in [0, 1]$  where  $f(x) = y$ .

Let  $x = y/2$ . Since  $y \in [0, 2]$ , we know  $0 \leq y \leq 2$ , and therefore that  $0 \leq y/2 \leq 1$ . We picked  $x = y/2$ , so we know that  $0 \leq x \leq 1$ , which in turn means  $x \in [0, 1]$ . Moreover, notice that

$$f(x) = 2x = 2(y/2) = y,$$

so  $f(x) = y$ , as required. ■

# Home on the Range



$$f : [0, 1] \rightarrow [0, 2]$$
$$f(x) = 2x$$



# Home on the Range




$$f : [0, 1] \rightarrow [0, 3]$$
$$f(x) = 3x$$

# Home on the Range



$$f : [0, 1] \rightarrow [0, 137]$$
$$f(x) = 137x$$

# Home on the Range




$$f : [0, 1] \rightarrow [0, k]$$
$$f(x) = kx$$

This means that *cardinality* (how many points there are) is a different idea than *mass* (how much those points weight). Look into *measure theory* if you're curious to learn more!

And one more example, just for funzies.

# Put a Ring On It



$$f : (-\pi/2, \pi/2) \rightarrow \mathbb{R}$$
$$f(x) = \tan x$$

$$|(-\pi/2, \pi/2)| = |\mathbb{R}|$$

# Some Properties of Cardinality

***Theorem:*** For any set  $A$ , we have  $|A| = |A|$ .

**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:**

Which of the following is the right high-level way to approach this proof?

- A. Pick an arbitrary set  $A$ , then find a bijection  $f : A \rightarrow A$ .
- B. Pick an arbitrary set  $A$  and show every function  $f : A \rightarrow A$  is bijective.
- C. There's nothing to prove here. Every object is equal to itself.

Answer at **PollEv.com/cs103** or  
text **CS103** to **22333** once to join, then **A**, **B**, or **C**.



**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:** Consider any set  $A$ , and let  $f: A \rightarrow A$  be the function defined as  $f(x) = x$ .

**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:** Consider any set  $A$ , and let  $f: A \rightarrow A$  be the function defined as  $f(x) = x$ . We will prove that  $f$  is a bijection.

**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:** Consider any set  $A$ , and let  $f : A \rightarrow A$  be the function defined as  $f(x) = x$ . We will prove that  $f$  is a bijection.

First, we'll show that  $f$  is a well-defined function.

**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:** Consider any set  $A$ , and let  $f: A \rightarrow A$  be the function defined as  $f(x) = x$ . We will prove that  $f$  is a bijection.

First, we'll show that  $f$  is a well-defined function. To see this, note that for any  $x \in A$ , we have  $f(x) = x \in A$ , as needed.

**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:** Consider any set  $A$ , and let  $f : A \rightarrow A$  be the function defined as  $f(x) = x$ . We will prove that  $f$  is a bijection.

First, we'll show that  $f$  is a well-defined function. To see this, note that for any  $x \in A$ , we have  $f(x) = x \in A$ , as needed.

Next, we'll show that  $f$  is injective.

**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:** Consider any set  $A$ , and let  $f: A \rightarrow A$  be the function defined as  $f(x) = x$ . We will prove that  $f$  is a bijection.

First, we'll show that  $f$  is a well-defined function. To see this, note that for any  $x \in A$ , we have  $f(x) = x \in A$ , as needed.

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in A$  where  $f(x_1) = f(x_2)$ .

**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:** Consider any set  $A$ , and let  $f: A \rightarrow A$  be the function defined as  $f(x) = x$ . We will prove that  $f$  is a bijection.

First, we'll show that  $f$  is a well-defined function. To see this, note that for any  $x \in A$ , we have  $f(x) = x \in A$ , as needed.

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in A$  where  $f(x_1) = f(x_2)$ . We need to show that  $x_1 = x_2$ .

**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:** Consider any set  $A$ , and let  $f: A \rightarrow A$  be the function defined as  $f(x) = x$ . We will prove that  $f$  is a bijection.

First, we'll show that  $f$  is a well-defined function. To see this, note that for any  $x \in A$ , we have  $f(x) = x \in A$ , as needed.

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in A$  where  $f(x_1) = f(x_2)$ . We need to show that  $x_1 = x_2$ . Since  $f(x_1) = f(x_2)$ , we see by definition of  $f$  that  $x_1 = x_2$ , as required.



**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:** Consider any set  $A$ , and let  $f: A \rightarrow A$  be the function defined as  $f(x) = x$ . We will prove that  $f$  is a bijection.

First, we'll show that  $f$  is a well-defined function. To see this, note that for any  $x \in A$ , we have  $f(x) = x \in A$ , as needed.

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in A$  where  $f(x_1) = f(x_2)$ . We need to show that  $x_1 = x_2$ . Since  $f(x_1) = f(x_2)$ , we see by definition of  $f$  that  $x_1 = x_2$ , as required.

Finally, we'll show that  $f$  is surjective.

**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:** Consider any set  $A$ , and let  $f: A \rightarrow A$  be the function defined as  $f(x) = x$ . We will prove that  $f$  is a bijection.

First, we'll show that  $f$  is a well-defined function. To see this, note that for any  $x \in A$ , we have  $f(x) = x \in A$ , as needed.

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in A$  where  $f(x_1) = f(x_2)$ . We need to show that  $x_1 = x_2$ . Since  $f(x_1) = f(x_2)$ , we see by definition of  $f$  that  $x_1 = x_2$ , as required.

Finally, we'll show that  $f$  is surjective. Consider any  $y \in A$ .

**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:** Consider any set  $A$ , and let  $f: A \rightarrow A$  be the function defined as  $f(x) = x$ . We will prove that  $f$  is a bijection.

First, we'll show that  $f$  is a well-defined function. To see this, note that for any  $x \in A$ , we have  $f(x) = x \in A$ , as needed.

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in A$  where  $f(x_1) = f(x_2)$ . We need to show that  $x_1 = x_2$ . Since  $f(x_1) = f(x_2)$ , we see by definition of  $f$  that  $x_1 = x_2$ , as required.

Finally, we'll show that  $f$  is surjective. Consider any  $y \in A$ . We will prove that there is some  $x \in A$  where  $f(x) = y$ .

**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:** Consider any set  $A$ , and let  $f: A \rightarrow A$  be the function defined as  $f(x) = x$ . We will prove that  $f$  is a bijection.

First, we'll show that  $f$  is a well-defined function. To see this, note that for any  $x \in A$ , we have  $f(x) = x \in A$ , as needed.

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in A$  where  $f(x_1) = f(x_2)$ . We need to show that  $x_1 = x_2$ . Since  $f(x_1) = f(x_2)$ , we see by definition of  $f$  that  $x_1 = x_2$ , as required.

Finally, we'll show that  $f$  is surjective. Consider any  $y \in A$ . We will prove that there is some  $x \in A$  where  $f(x) = y$ . Pick  $x = y$ .

**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:** Consider any set  $A$ , and let  $f: A \rightarrow A$  be the function defined as  $f(x) = x$ . We will prove that  $f$  is a bijection.

First, we'll show that  $f$  is a well-defined function. To see this, note that for any  $x \in A$ , we have  $f(x) = x \in A$ , as needed.

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in A$  where  $f(x_1) = f(x_2)$ . We need to show that  $x_1 = x_2$ . Since  $f(x_1) = f(x_2)$ , we see by definition of  $f$  that  $x_1 = x_2$ , as required.

Finally, we'll show that  $f$  is surjective. Consider any  $y \in A$ . We will prove that there is some  $x \in A$  where  $f(x) = y$ . Pick  $x = y$ . Then  $x \in A$  (since  $y \in A$ ) and  $f(x) = x = y$ , as required.

**Theorem:** For any set  $A$ , we have  $|A| = |A|$ .

**Proof:** Consider any set  $A$ , and let  $f : A \rightarrow A$  be the function defined as  $f(x) = x$ . We will prove that  $f$  is a bijection.

First, we'll show that  $f$  is a well-defined function. To see this, note that for any  $x \in A$ , we have  $f(x) = x \in A$ , as needed.

Next, we'll show that  $f$  is injective. Pick any  $x_1, x_2 \in A$  where  $f(x_1) = f(x_2)$ . We need to show that  $x_1 = x_2$ . Since  $f(x_1) = f(x_2)$ , we see by definition of  $f$  that  $x_1 = x_2$ , as required.

Finally, we'll show that  $f$  is surjective. Consider any  $y \in A$ . We will prove that there is some  $x \in A$  where  $f(x) = y$ . Pick  $x = y$ . Then  $x \in A$  (since  $y \in A$ ) and  $f(x) = x = y$ , as required. ■

***Theorem:*** If  $A$ ,  $B$ , and  $C$  are sets where  $|A| = |B|$  and  $|B| = |C|$ , then  $|A| = |C|$ .

***Theorem:*** If  $A$ ,  $B$ , and  $C$  are sets where  $|A| = |B|$  and  $|B| = |C|$ , then  $|A| = |C|$ .

***Proof:***



**Theorem:** If  $A$ ,  $B$ , and  $C$  are sets where  $|A| = |B|$  and  $|B| = |C|$ , then  $|A| = |C|$ .

**Proof:** Consider any sets  $A$ ,  $B$ , and  $C$  where  $|A| = |B|$  and  $|B| = |C|$ .

**Theorem:** If  $A$ ,  $B$ , and  $C$  are sets where  $|A| = |B|$  and  $|B| = |C|$ , then  $|A| = |C|$ .

**Proof:** Consider any sets  $A$ ,  $B$ , and  $C$  where  $|A| = |B|$  and  $|B| = |C|$ . We need to prove that  $|A| = |C|$ .

**Theorem:** If  $A$ ,  $B$ , and  $C$  are sets where  $|A| = |B|$  and  $|B| = |C|$ , then  $|A| = |C|$ .

**Proof:** Consider any sets  $A$ ,  $B$ , and  $C$  where  $|A| = |B|$  and  $|B| = |C|$ . We need to prove that  $|A| = |C|$ . To do so, we need to show that there is a bijection from  $A$  to  $C$ .

**Theorem:** If  $A$ ,  $B$ , and  $C$  are sets where  $|A| = |B|$  and  $|B| = |C|$ , then  $|A| = |C|$ .

**Proof:** Consider any sets  $A$ ,  $B$ , and  $C$  where  $|A| = |B|$  and  $|B| = |C|$ . We need to prove that  $|A| = |C|$ . To do so, we need to show that there is a bijection from  $A$  to  $C$ .

Since  $|A| = |B|$ , we know that there is a some bijection  $f : A \rightarrow B$ .

**Theorem:** If  $A$ ,  $B$ , and  $C$  are sets where  $|A| = |B|$  and  $|B| = |C|$ , then  $|A| = |C|$ .

**Proof:** Consider any sets  $A$ ,  $B$ , and  $C$  where  $|A| = |B|$  and  $|B| = |C|$ . We need to prove that  $|A| = |C|$ . To do so, we need to show that there is a bijection from  $A$  to  $C$ .

Since  $|A| = |B|$ , we know that there is a some bijection  $f : A \rightarrow B$ . Similarly, since  $|B| = |C|$  we know that there is at least one bijection  $g : B \rightarrow C$ .

**Theorem:** If  $A$ ,  $B$ , and  $C$  are sets where  $|A| = |B|$  and  $|B| = |C|$ , then  $|A| = |C|$ .

**Proof:** Consider any sets  $A$ ,  $B$ , and  $C$  where  $|A| = |B|$  and  $|B| = |C|$ . We need to prove that  $|A| = |C|$ . To do so, we need to show that there is a bijection from  $A$  to  $C$ .

Since  $|A| = |B|$ , we know that there is a some bijection  $f : A \rightarrow B$ . Similarly, since  $|B| = |C|$  we know that there is at least one bijection  $g : B \rightarrow C$ .

Consider the function  $g \circ f : A \rightarrow C$ .

**Theorem:** If  $A$ ,  $B$ , and  $C$  are sets where  $|A| = |B|$  and  $|B| = |C|$ , then  $|A| = |C|$ .

**Proof:** Consider any sets  $A$ ,  $B$ , and  $C$  where  $|A| = |B|$  and  $|B| = |C|$ . We need to prove that  $|A| = |C|$ . To do so, we need to show that there is a bijection from  $A$  to  $C$ .

Since  $|A| = |B|$ , we know that there is a some bijection  $f : A \rightarrow B$ . Similarly, since  $|B| = |C|$  we know that there is at least one bijection  $g : B \rightarrow C$ .

Consider the function  $g \circ f : A \rightarrow C$ . Since  $g$  and  $f$  are bijections and the composition of two bijections is a bijection, we see that  $g \circ f$  is a bijection from  $A$  to  $C$ .

**Theorem:** If  $A$ ,  $B$ , and  $C$  are sets where  $|A| = |B|$  and  $|B| = |C|$ , then  $|A| = |C|$ .

**Proof:** Consider any sets  $A$ ,  $B$ , and  $C$  where  $|A| = |B|$  and  $|B| = |C|$ . We need to prove that  $|A| = |C|$ . To do so, we need to show that there is a bijection from  $A$  to  $C$ .

Since  $|A| = |B|$ , we know that there is a some bijection  $f : A \rightarrow B$ . Similarly, since  $|B| = |C|$  we know that there is at least one bijection  $g : B \rightarrow C$ .

Consider the function  $g \circ f : A \rightarrow C$ . Since  $g$  and  $f$  are bijections and the composition of two bijections is a bijection, we see that  $g \circ f$  is a bijection from  $A$  to  $C$ . Thus  $|A| = |C|$ , as required.



**Theorem:** If  $A$ ,  $B$ , and  $C$  are sets where  $|A| = |B|$  and  $|B| = |C|$ , then  $|A| = |C|$ .

**Proof:** Consider any sets  $A$ ,  $B$ , and  $C$  where  $|A| = |B|$  and  $|B| = |C|$ . We need to prove that  $|A| = |C|$ . To do so, we need to show that there is a bijection from  $A$  to  $C$ .

Since  $|A| = |B|$ , we know that there is a some bijection  $f : A \rightarrow B$ . Similarly, since  $|B| = |C|$  we know that there is at least one bijection  $g : B \rightarrow C$ .

Consider the function  $g \circ f : A \rightarrow C$ . Since  $g$  and  $f$  are bijections and the composition of two bijections is a bijection, we see that  $g \circ f$  is a bijection from  $A$  to  $C$ . Thus  $|A| = |C|$ , as required. ■

***Great exercise:*** Prove that if  $A$  and  $B$  are sets where  $|A| = |B|$ , then  $|B| = |A|$ .

Time-Out for Announcements!


# Midterm Exam Logistics

- Our first midterm exam is next **Monday, February 5<sup>th</sup>**, from **7:00PM - 10:00PM**. Locations are divvied up by last (family) name:
  - A - H: Go to Cubberley Auditorium.
  - I - Z: Go to 320-105.
- You're responsible for Lectures 00 – 05 and topics covered in PS1 – PS2. Later lectures (relations forward) and problem sets (PS3 onward) won't be tested here.
- The exam is closed-book, closed-computer, and limited-note. You can bring a double-sided, 8.5" × 11" sheet of notes with you to the exam, decorated however you'd like.
- Students with OAE accommodations: we will be reaching out to you soon with room and time assignments.

# Practice Midterm Exam


- To help you prepare for the midterm, we'll be holding a practice midterm exam **tonight** from **7PM - 10PM** in **Cemex Auditorium**.
  - The exam we'll use isn't one of the ones posted up on the course website, so feel free to use those as practice in the meantime.
- The practice midterm exam is an actual midterm we gave out in a previous quarter. It's probably the best indicator of what you should expect to see.
- Course staff will be on hand to answer your questions.
- Can't make it? We'll release that practice exam and solutions online. Set up your own practice exam time with a small group and work through it under realistic conditions!

# Problem Set 2 Grades




Average grades are *malicious lies*. Ignore them.  
Standard deviations are *malicious lies*. Ignore them.

# Problem Set 2 Grades



“Excellent job! Look over your feedback to find those last few spots to patch up.”


# Problem Set 2 Grades



"Good job! Take a look at your feedback to see what areas you need to focus on for next time."




# Problem Set 2 Grades




"A solid performance! Seems like there are a few spots you may want to get more practice with. Review your feedback, and let us know how we can help!"

# Problem Set 2 Grades



"You're on the right track, but it looks like something hasn't quite clicked yet. Come by office hours with questions – we're happy to help out!"

# Problem Set 2 Grades



"You're not where you need to be now, but we know you can do this. Stop by office hours and let us know what we can do to help out!"

# Next Steps

- Regardless of how you did on the problem set, make sure you understand all the feedback you've received, *especially* on the first-order translations and the proofs.
  - Like, seriously, do this. You don't want to make the same mistakes on the midterm!
- Ask questions on Piazza or stop by office hours if you have questions – we're happy to help out.

# Problem Set Three

- The Problem Set Three checkpoint has been graded.
  - ***Please, please, please review your feedback!*** That problem was tricky and a lot of people had a lot of trouble with it.
- Remaining problems are due on Friday at 2:30PM. Be strategic about taking late days.

Back to CS103!

# Unequal Cardinalities


- Recall:  $|A| = |B|$  if the following statement is true:

**There exists a bijection  $f : A \rightarrow B$**

- What does it mean for  $|A| \neq |B|$  to be true?

**Every function  $f : A \rightarrow B$  is not a bijection.**

- This is a strong statement! To prove  $|A| \neq |B|$ , we need to show that *no possible function* from  $A$  to  $B$  can be injective and surjective.



# Unequal Cardinalities


- Recall:  $|A| = |B|$  if the following statement is true:

**There exists a bijection  $f : A \rightarrow B$**

- What does it mean for  $|A| \neq |B|$  to be true?

**Every function  $f : A \rightarrow B$  is not a bijection.**

- This is a strong statement! To prove  $|A| \neq |B|$ , we need to show that *no possible function* from  $A$  to  $B$  can be injective and surjective.





# Unequal Cardinalities


- Recall:  $|A| = |B|$  if the following statement is true:

**There exists a bijection  $f : A \rightarrow B$**

- What does it mean for  $|A| \neq |B|$  to be true?

**Every function  $f : A \rightarrow B$  is not a bijection.**

- This is a strong statement! To prove  $|A| \neq |B|$ , we need to show that *no possible function* from  $A$  to  $B$  can be injective and surjective.



# Unequal Cardinalities


- Recall:  $|A| = |B|$  if the following statement is true:

**There exists a bijection  $f : A \rightarrow B$**

- What does it mean for  $|A| \neq |B|$  to be true?

**Every function  $f : A \rightarrow B$  is not a bijection.**

- This is a strong statement! To prove  $|A| \neq |B|$ , we need to show that *no possible function* from  $A$  to  $B$  can be injective and surjective.



# Unequal Cardinalities


- Recall:  $|A| = |B|$  if the following statement is true:

**There exists a bijection  $f : A \rightarrow B$**

- What does it mean for  $|A| \neq |B|$  to be true?

**Every function  $f : A \rightarrow B$  is not a bijection.**

- This is a strong statement! To prove  $|A| \neq |B|$ , we need to show that *no possible function* from  $A$  to  $B$  can be injective and surjective.



# Cantor's Theorem Revisited

# Cantor's Theorem

- In our very first lecture, we sketched out a proof of ***Cantor's theorem***, which says that

**If  $S$  is a set, then  $|S| < |\wp(S)|$ .**

- That proof was visual and pretty hand-wavy. Let's see if we can go back and formalize it!

# Where We're Going

- Today, we're going to formally prove the following result:

**If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .**

- We've released an online Guide to Cantor's Theorem, which will go into *way* more depth than what we're going to see here.
- The goal for today will be to see how to start with our picture and turn it into something rigorous.
- On the next problem set, you'll explore the proof in more depth and see some other applications.

# The Roadmap

- We're going to prove this statement:

If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

- Here's how this will work:
  - Pick an arbitrary set  $S$ .
  - Pick an arbitrary function  $f : S \rightarrow \wp(S)$ .
  - Show that  $f$  is not surjective using a diagonal argument.
  - Conclude that there are no bijections from  $S$  to  $\wp(S)$ .
  - Conclude that  $|S| \neq |\wp(S)|$ .

# The Roadmap

We're going to prove this statement:

If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

Here's how this will work:

Pick an arbitrary set  $S$ .

Pick an arbitrary function  $f : S \rightarrow \wp(S)$ .

- Show that  $f$  is not surjective using a diagonal argument.

Conclude that there are no bijections from  $S$  to  $\wp(S)$ .

Conclude that  $|S| \neq |\wp(S)|$ .



$x_0$

$x_1$

$x_2$

$x_3$

$x_4$

$x_5$

$\dots$

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

$$x_0 \longrightarrow \{ x_0, x_2, x_4, \dots \}$$

$$x_1 \longrightarrow \{ x_0, x_3, x_4, \dots \}$$

$$x_2 \longrightarrow \{ x_4, \dots \}$$

$$x_3 \longrightarrow \{ x_1, x_4, \dots \}$$

$$x_4 \longrightarrow \{ x_0, x_5, \dots \}$$

$$x_5 \longrightarrow \{ x_0, x_1, x_2, x_3, x_4, x_5, \dots \}$$

...

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$\dots$
-------	-------	-------	-------	-------	-------	---------

$$x_0 \longrightarrow \{ x_0, x_2, x_4, \dots \}$$

$$x_1 \longrightarrow \{ x_0, x_3, x_4, \dots \}$$

$$x_2 \longrightarrow \{ x_4, \dots \}$$

$$x_3 \longrightarrow \{ x_1, x_4, \dots \}$$

$$x_4 \longrightarrow \{ x_0, x_5, \dots \}$$

$$x_5 \longrightarrow \{ x_0, x_1, x_2, x_3, x_4, x_5, \dots \}$$

$\dots$

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$ $\longrightarrow$	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...

$x_1 \longrightarrow \{ x_0, x_3, x_4, \dots \}$

$x_2 \longrightarrow \{ x_4, \dots \}$

$x_3 \longrightarrow \{ x_1, x_4, \dots \}$

$x_4 \longrightarrow \{ x_0, x_5, \dots \}$

$x_5 \longrightarrow \{ x_0, x_1, x_2, x_3, x_4, x_5, \dots \}$

...

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

		$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$\dots$
$x_0$	$\longrightarrow$	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	$\dots$
$x_1$	$\longrightarrow$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	$\dots$

$$x_2 \longrightarrow \{ x_4, \dots \}$$
$$x_3 \longrightarrow \{ x_1, x_4, \dots \}$$
$$x_4 \longrightarrow \{ x_0, x_5, \dots \}$$
$$x_5 \longrightarrow \{ x_0, x_1, x_2, x_3, x_4, x_5, \dots \}$$

• • •

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0 \longrightarrow$	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_1 \longrightarrow$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	...
$x_2 \longrightarrow$	{ $x_4$ , ... }						
$x_3 \longrightarrow$	{ $x_1$ , $x_4$ , ... }						
$x_4 \longrightarrow$	{ $x_0$ , $x_5$ , ... }						
$x_5 \longrightarrow$	{ $x_0$ , $x_1$ , $x_2$ , $x_3$ , $x_4$ , $x_5$ , ... }						
...							

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

		$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$\dots$
$x_0$	→	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	$\dots$
$x_1$	→	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	$\dots$
$x_2$	→	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	$\dots$

$$x_3 \longrightarrow \{ x_1, x_4, \dots \}$$
$$x_4 \longrightarrow \{ x_0, x_5, \dots \}$$
$$x_5 \longrightarrow \{ x_0, x_1, x_2, x_3, x_4, x_5, \dots \}$$

• • •

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0 \longrightarrow$	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_1 \longrightarrow$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	...
$x_2 \longrightarrow$	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_3 \longrightarrow$	{ $x_1$ , $x_4$ , ... }						}
$x_4 \longrightarrow$	{ $x_0$ , $x_5$ , ... }						}
$x_5 \longrightarrow$	{ $x_0$ , $x_1$ , $x_2$ , $x_3$ , $x_4$ , $x_5$ , ... }						}
...							



*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$\dots$
$x_0 \rightarrow$	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	$\dots$
$x_1 \rightarrow$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	$\dots$
$x_2 \rightarrow$	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	$\dots$
$x_3 \rightarrow$	<b>N</b>	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	$\dots$

$$x_4 \longrightarrow \{ x_0, x_5, \dots \}$$
$$x_5 \longrightarrow \{ x_0, x_1, x_2, x_3, x_4, x_5, \dots \}$$

• • •

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0 \longrightarrow$	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_1 \longrightarrow$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	...
$x_2 \longrightarrow$	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_3 \longrightarrow$	<b>N</b>	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_4 \longrightarrow$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	...
$x_5 \longrightarrow$	$\{ x_0, x_1, x_2, x_3, x_4, x_5, \dots \}$						
...							

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$ →	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_1$ →	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	...
$x_2$ →	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_3$ →	<b>N</b>	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_4$ →	<b>Y</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	...
$x_5$ →	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	...
...							

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$ →	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_1$ →	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	...
$x_2$ →	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_3$ →	<b>N</b>	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_4$ →	<b>Y</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	...
$x_5$ →	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	...
...	...	...	...	...	...	...	...

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_1$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	...
$x_2$	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_3$	<b>N</b>	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_4$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	...
$x_5$	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	...
...	...	...	...	...	...	...	...

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_1$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	...
$x_2$	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_3$	<b>N</b>	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_4$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	...
$x_5$	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	...
...	...	...	...	...	...	...	...

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_1$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	...
$x_2$	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_3$	<b>N</b>	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_4$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	...
$x_5$	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	...
...	...	...	...	...	...	...	...

$\left\{ x_0', \quad \quad \quad x_5', \quad \dots \right\}$

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	N	N	...
$x_4$	N	N	N	N	N	N	...
$x_5$	N	N	N	N	N	N	...
...	...	...	...	...	...	...	...

For this choice of  $f$ , what set  
is shown in blue below?

Answer at **PollEv.com/cs103** or  
text **CS103** to **22333** once to join, then your answer.

*Flip all Y's to  
N's and vice-  
versa to get a  
new set*

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----



*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

*Flip all Y's to  
N's and vice-  
versa to get a  
new set*

{  $x_1', x_2', x_3', x_4', \dots$  }



*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

*Which row in the  
table is paired  
with this set?*

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_1$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	...
$x_2$	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_3$	<b>N</b>	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_4$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	...
$x_5$	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	...
...	...	...	...	...	...	...	...

*Which row in the  
table is paired  
with this set?*

<b>N</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	...
----------	----------	----------	----------	----------	----------	-----

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

*Which row in the  
table is paired  
with this set?*

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

*Which row in the  
table is paired  
with this set?*

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

*Which row in the  
table is paired  
with this set?*

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

*Which row in the  
table is paired  
with this set?*

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

*Which row in the  
table is paired  
with this set?*

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----



*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
Y	N	Y	N	Y	N	...
Y	N	N	Y	Y	N	...
N	N	N	N	Y	N	...
N	Y	N	N	Y	N	...
Y	N	N	N	N	Y	...
Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

Which row in the table is paired with this set?

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

*What set is this?*

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

*What set is this?*

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...
	N	Y	Y	Y	Y	N	...

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

$f(x_0)$

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_1$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	...
$x_2$	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_3$	<b>N</b>	<b>Y</b>	<b>N</b>	<b>N</b>	<b>Y</b>	<b>N</b>	...
$x_4$	<b>Y</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>	<b>Y</b>	...
$x_5$	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	...
...	...	...	...	...	...	...	...

<b>N</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>Y</b>	<b>N</b>	...
----------	----------	----------	----------	----------	----------	-----

$x_0 \in f(x_0)?$

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

$x_0 \notin f(x_0)$ ?

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

$f(x_1)$



*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

$x_1 \in f(x_1)?$

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

$x_1 \notin f(x_1)$ ?

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

$f(x_2)$

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

$x_2 \in f(x_2)?$

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

$x_2 \notin f(x_2)$ ?

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

$x_3 \notin f(x_3)$ ?

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

$x_4 \notin f(x_4)$ ?



*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

$x_5 \notin f(x_5)?$

N



*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
Y	N	Y	N	Y	N	...
Y	N	N	Y	Y	N	...
N	N	N	N	Y	N	...
N	Y	N	N	Y	N	...
Y	N	N	N	N	Y	...
Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...

...

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

*This is a drawing  
of our function  
 $f : S \rightarrow \wp(S)$ .*

	$x_0$	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	...
$x_0$	Y	N	Y	N	Y	N	...
$x_1$	Y	N	N	Y	Y	N	...
$x_2$	N	N	N	N	Y	N	...
$x_3$	N	Y	N	N	Y	N	...
$x_4$	Y	N	N	N	N	Y	...
$x_5$	Y	Y	Y	Y	Y	Y	...
...	...	...	...	...	...	...	...

$\{ x \in S \mid x \notin f(x) \}$

N	Y	Y	Y	Y	N	...
---	---	---	---	---	---	-----

# The Diagonal Set

- For any set  $S$  and function  $f : S \rightarrow \wp(S)$ , we can define a set  $D$  as follows:

$$D = \{ x \in S \mid x \notin f(x) \}$$

*(“The set of all elements  $x$  where  $x$  is not an element of the set  $f(x)$ .”)*

- This is a formalization of the set we found in the previous picture.
- Using this choice of  $D$ , we can formally prove that no function  $f : S \rightarrow \wp(S)$  is a bijection.

***Theorem:*** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

***Theorem:*** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

***Proof:*** Let  $S$  be an arbitrary set.

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ .

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ .

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.



**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

We will show that there is no  $y \in S$  such that  $f(y) = D$ .

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

We will show that there is no  $y \in S$  such that  $f(y) = D$ . To do so, we proceed by contradiction.

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

We will show that there is no  $y \in S$  such that  $f(y) = D$ . To do so, we proceed by contradiction. Suppose that there is some  $y \in S$  such that  $f(y) = D$ .

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

We will show that there is no  $y \in S$  such that  $f(y) = D$ . To do so, we proceed by contradiction. Suppose that there is some  $y \in S$  such that  $f(y) = D$ . By the definition of  $D$ , we know that

$$y \in D \text{ iff } y \notin f(y). \quad (2)$$

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

We will show that there is no  $y \in S$  such that  $f(y) = D$ . To do so, we proceed by contradiction. Suppose that there is some  $y \in S$  such that  $f(y) = D$ . By the definition of  $D$ , we know that

$$y \in D \text{ iff } y \notin f(y). \quad (2)$$

By assumption,  $f(y) = D$ .

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

We will show that there is no  $y \in S$  such that  $f(y) = D$ . To do so, we proceed by contradiction. Suppose that there is some  $y \in S$  such that  $f(y) = D$ . By the definition of  $D$ , we know that

$$y \in D \text{ iff } y \notin f(y). \quad (2)$$

By assumption,  $f(y) = D$ . Combined with (2), this tells us

$$y \in D \text{ iff } y \notin D. \quad (3)$$

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

We will show that there is no  $y \in S$  such that  $f(y) = D$ . To do so, we proceed by contradiction. Suppose that there is some  $y \in S$  such that  $f(y) = D$ . By the definition of  $D$ , we know that

$$y \in D \text{ iff } y \notin f(y). \quad (2)$$

By assumption,  $f(y) = D$ . Combined with (2), this tells us

$$y \in D \text{ iff } y \notin D. \quad (3)$$



**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

We will show that there is no  $y \in S$  such that  $f(y) = D$ . To do so, we proceed by contradiction. Suppose that there is some  $y \in S$  such that  $f(y) = D$ . By the definition of  $D$ , we know that

$$y \in D \text{ iff } y \notin f(y). \quad (2)$$

By assumption,  $f(y) = D$ . Combined with (2), this tells us

$$y \in D \text{ iff } y \notin D. \quad (3)$$

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

We will show that there is no  $y \in S$  such that  $f(y) = D$ . To do so, we proceed by contradiction. Suppose that there is some  $y \in S$  such that  $f(y) = D$ . By the definition of  $D$ , we know that

$$y \in D \text{ iff } y \notin f(y). \quad (2)$$

By assumption,  $f(y) = D$ . Combined with (2), this tells us

$$y \in D \text{ iff } y \notin D. \quad (3)$$

This is impossible.

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

We will show that there is no  $y \in S$  such that  $f(y) = D$ . To do so, we proceed by contradiction. Suppose that there is some  $y \in S$  such that  $f(y) = D$ . By the definition of  $D$ , we know that

$$y \in D \text{ iff } y \notin f(y). \quad (2)$$

By assumption,  $f(y) = D$ . Combined with (2), this tells us

$$y \in D \text{ iff } y \notin D. \quad (3)$$

This is impossible. We have reached a contradiction, so our assumption must have been wrong.

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

We will show that there is no  $y \in S$  such that  $f(y) = D$ . To do so, we proceed by contradiction. Suppose that there is some  $y \in S$  such that  $f(y) = D$ . By the definition of  $D$ , we know that

$$y \in D \text{ iff } y \notin f(y). \quad (2)$$

By assumption,  $f(y) = D$ . Combined with (2), this tells us

$$y \in D \text{ iff } y \notin D. \quad (3)$$

This is impossible. We have reached a contradiction, so our assumption must have been wrong. Therefore, there is no  $y \in S$  such that  $f(y) = D$ , so  $f$  is not surjective.

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

We will show that there is no  $y \in S$  such that  $f(y) = D$ . To do so, we proceed by contradiction. Suppose that there is some  $y \in S$  such that  $f(y) = D$ . By the definition of  $D$ , we know that

$$y \in D \text{ iff } y \notin f(y). \quad (2)$$

By assumption,  $f(y) = D$ . Combined with (2), this tells us

$$y \in D \text{ iff } y \notin D. \quad (3)$$

This is impossible. We have reached a contradiction, so our assumption must have been wrong. Therefore, there is no  $y \in S$  such that  $f(y) = D$ , so  $f$  is not surjective. This means that  $f$  is not a bijection, and since our choice of  $f$  was arbitrary, we conclude that there are no bijections between  $S$  and  $\wp(S)$ .

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

We will show that there is no  $y \in S$  such that  $f(y) = D$ . To do so, we proceed by contradiction. Suppose that there is some  $y \in S$  such that  $f(y) = D$ . By the definition of  $D$ , we know that

$$y \in D \text{ iff } y \notin f(y). \quad (2)$$

By assumption,  $f(y) = D$ . Combined with (2), this tells us

$$y \in D \text{ iff } y \notin D. \quad (3)$$

This is impossible. We have reached a contradiction, so our assumption must have been wrong. Therefore, there is no  $y \in S$  such that  $f(y) = D$ , so  $f$  is not surjective. This means that  $f$  is not a bijection, and since our choice of  $f$  was arbitrary, we conclude that there are no bijections between  $S$  and  $\wp(S)$ . Thus  $|S| \neq |\wp(S)|$ , as required.

**Theorem:** If  $S$  is a set, then  $|S| \neq |\wp(S)|$ .

**Proof:** Let  $S$  be an arbitrary set. We will prove that  $|S| \neq |\wp(S)|$  by showing that there are no bijections from  $S$  to  $\wp(S)$ . To do so, choose an arbitrary function  $f : S \rightarrow \wp(S)$ . We will prove that  $f$  is not surjective.

Starting with  $f$ , we define the set

$$D = \{ x \in S \mid x \notin f(x) \}. \quad (1)$$

We will show that there is no  $y \in S$  such that  $f(y) = D$ . To do so, we proceed by contradiction. Suppose that there is some  $y \in S$  such that  $f(y) = D$ . By the definition of  $D$ , we know that

$$y \in D \text{ iff } y \notin f(y). \quad (2)$$

By assumption,  $f(y) = D$ . Combined with (2), this tells us

$$y \in D \text{ iff } y \notin D. \quad (3)$$

This is impossible. We have reached a contradiction, so our assumption must have been wrong. Therefore, there is no  $y \in S$  such that  $f(y) = D$ , so  $f$  is not surjective. This means that  $f$  is not a bijection, and since our choice of  $f$  was arbitrary, we conclude that there are no bijections between  $S$  and  $\wp(S)$ . Thus  $|S| \neq |\wp(S)|$ , as required. ■

# The Big Recap

- We define equal cardinality in terms of bijections between sets.
- Lots of different sets of infinite size have the same cardinality.
- Cardinality acts like an equivalence relation – but only because we can prove specific properties of how it behaves by relying on properties of function.
- Cantor's theorem can be formalized in terms of surjectivity.



# Next Time

- ***Graphs***
  - A ubiquitous, expressive, and flexible abstraction!
- ***Properties of Graphs***
  - Building high-level structures out of lower-level ones!