

# Universal and Existential Statements

***Theorem:*** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Theorem:** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Proof:**

Which of the following should be the next sentence of this proof?

- A. "Pick any odd integer,  $n = 137$ ."
- B. "Pick any odd integer  $n$ ."
- C. "Pick any odd integer  $n$  and arbitrary integers  $r$  and  $s$  where  $r^2 - s^2 = n$ ."

Answer at **PollEv.com/cs103** or  
text **CS103** to **22333** once to join, then **A**, **B**, or **C**.

***Theorem:*** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

***Proof:*** Pick any odd integer  $n$ .

***Theorem:*** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

***Proof:*** Pick any odd integer  $n$ .

**Theorem:** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Proof:** Pick any odd integer  $n$ .

This is a very different sort of request than what we've seen in the past. How on earth do we go about proving something like this?

# Universal vs. Existential Statements

- A ***universal statement*** is a statement of the form

**For all  $x$ , [some-property] holds for  $x$ .**

- We've seen how to prove these statements.
- An ***existential statement*** is a statement of the form

**There is some  $x$  where [some-property] holds for  $x$ .**

- How do you prove an existential statement?

# Proving an Existential Statement

- Over the course of the quarter, we will see several different ways to prove an existential statement of the form

**There is an  $x$  where [some-property] holds for  $x$ .**

- ***Simplest approach:*** Search far and wide, find an  $x$  that has the right property, then show why your choice is correct.



***Theorem:*** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .


***Proof:*** Pick any odd integer  $n$ .

**Theorem:** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Proof:** Pick any odd integer  $n$ . Since  $n$  is odd, we know there is some integer  $k$  where  $n = 2k + 1$ .


**Theorem:** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Proof:** Pick any odd integer  $n$ . Since  $n$  is odd, we know there is some integer  $k$  where  $n = 2k + 1$ .




**Theorem:** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Proof:** Pick any odd integer  $n$ . Since  $n$  is odd, we know there is some integer  $k$  where  $n = 2k + 1$ .




**Theorem:** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Proof:** Pick any odd integer  $n$ . Since  $n$  is odd, we know there is some integer  $k$  where  $n = 2k + 1$ .



**Theorem:** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Proof:** Pick any odd integer  $n$ . Since  $n$  is odd, we know there is some integer  $k$  where  $n = 2k + 1$ .



**Theorem:** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Proof:** Pick any odd integer  $n$ . Since  $n$  is odd, we know there is some integer  $k$  where  $n = 2k + 1$ .

Now, let  $r = k+1$  and  $s = k$ .

**Theorem:** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Proof:** Pick any odd integer  $n$ . Since  $n$  is odd, we know there is some integer  $k$  where  $n = 2k + 1$ .

Now, let  $r = k+1$  and  $s = k$ . Then we see that

$$r^2 - s^2 = (k+1)^2 - k^2$$



**Theorem:** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Proof:** Pick any odd integer  $n$ . Since  $n$  is odd, we know there is some integer  $k$  where  $n = 2k + 1$ .

Now, let  $r = k+1$  and  $s = k$ . Then we see that

$$\begin{aligned} r^2 - s^2 &= (k+1)^2 - k^2 \\ &= k^2 + 2k + 1 - k^2 \end{aligned}$$

**Theorem:** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Proof:** Pick any odd integer  $n$ . Since  $n$  is odd, we know there is some integer  $k$  where  $n = 2k + 1$ .

Now, let  $r = k+1$  and  $s = k$ . Then we see that

$$\begin{aligned} r^2 - s^2 &= (k+1)^2 - k^2 \\ &= k^2 + 2k + 1 - k^2 \\ &= 2k + 1 \end{aligned}$$

**Theorem:** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Proof:** Pick any odd integer  $n$ . Since  $n$  is odd, we know there is some integer  $k$  where  $n = 2k + 1$ .

Now, let  $r = k+1$  and  $s = k$ . Then we see that

$$\begin{aligned} r^2 - s^2 &= (k+1)^2 - k^2 \\ &= k^2 + 2k + 1 - k^2 \\ &= 2k + 1 \\ &= n. \end{aligned}$$

**Theorem:** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Proof:** Pick any odd integer  $n$ . Since  $n$  is odd, we know there is some integer  $k$  where  $n = 2k + 1$ .

Now, let  $r = k+1$  and  $s = k$ . Then we see that

$$\begin{aligned} r^2 - s^2 &= (k+1)^2 - k^2 \\ &= k^2 + 2k + 1 - k^2 \\ &= 2k + 1 \\ &= n. \end{aligned}$$

This means that  $r^2 - s^2 = n$ , which is what we needed to show.

**Theorem:** For any odd integer  $n$ , there exist integers  $r$  and  $s$  where  $r^2 - s^2 = n$ .

**Proof:** Pick any odd integer  $n$ . Since  $n$  is odd, we know there is some integer  $k$  where  $n = 2k + 1$ .

Now, let  $r = k+1$  and  $s = k$ . Then we see that

$$\begin{aligned} r^2 - s^2 &= (k+1)^2 - k^2 \\ &= k^2 + 2k + 1 - k^2 \\ &= 2k + 1 \\ &= n. \end{aligned}$$

This means that  $r^2 - s^2 = n$ , which is what we needed to show. ■

***Follow-Up Question:*** There are some integers that can't be written as  $r^2 - s^2$  for any integers  $r$  and  $s$ .

Can you prove that every integer can be formed by adding and subtracting some combination of at most *three* perfect squares?