




Bilayer start 128 lipids, equilibrate at $T = 0.8$, skip first 1000τ




$A_{pl} = 0.5 \text{ nm}^2$




$A_{pl} = 0.85 \text{ nm}^2$



$A_{pl} = 12 \text{ nm}^2$








→ 128 lipids: $\text{apl} = 0.85 \text{ nm}^2$, $\text{wpl} = 22$, $\rho = 3$

→ 256 lipids: $\text{apl} = 0.88 \text{ nm}^2$, $\text{wpl} = 24$, $\rho = 3$




T = 1, 128 lipids





- 1) Lipid soup
- 2) Equilibration at T = 2
- 3) Cool down to T = 1
- 4) Production at T = 1




$T = 1$, 128 lipids





$T = 0.8$, 128 lipids





- 1) Lipid soup
- 2) Equilibration at $T = 2$
- 3) Cool down to $T = 0.8$
- 4) Production at $T = 0.8$




$T = 0.8$, 128 lipids






$T = 1$, 256 lipids





- 1) Lipids soup
- 2) Equilibrate at $T = 2$
- 3) Cool down to $T = 1$
- 4) Deform Box to fit into quadratic form
- 5) Production at $T = 1$




$T = 1$, 256 lipids



$T = 0.8$, 256 lipids



- 1) Lipids soup
- 2) Equilibrate at $T = 2$
- 3) Cool down to $T = 1$
- 4) Deform Box to fit into quadratic form
- 5) Production at $T = 1$



$T = 0.8$, 256 lipids

