

Dependent T-Test in SPSS Statistics - The procedure for running the test, generating the output and understanding the output using a relevant example | Laerd Statistics

Laerd Statistics [Login](#)[Cookies & Privacy](#)

[Take the Tour](#) [Plans & Pricing](#) [SIGN UP](#)

Dependent T-Test using SPSS Statistics

Introduction

The dependent t-test (called the paired-samples t-test in SPSS Statistics) compares the means between two related groups on the same continuous, dependent variable. For example, you could use a dependent t-test to understand whether there was a difference in smokers' daily cigarette consumption before and after a 6 week hypnotherapy programme (i.e., your dependent variable would be "daily cigarette consumption", and your two related groups would be the cigarette consumption values "before" and "after" the hypnotherapy programme). If your dependent variable is dichotomous, you should instead use McNemar's test.

This "quick start" guide shows you how to carry out a dependent t-test using SPSS Statistics, as well as interpret and report the results from this test. However, before we introduce you to this procedure, you need to understand the different assumptions that your data must meet in order for a dependent t-test to give you a valid result. We discuss these assumptions next.

SPSS Statistics

Assumptions

When you choose to analyse your data using a dependent t-test, part of the process involves checking to make sure that the data you want to analyse can actually be analysed using a dependent t-test. You need to do this because it is only appropriate to use a dependent t-test if your data "passes" four assumptions that are required for a dependent t-test to give you a valid result. In practice, checking for these four assumptions just adds a little bit more time to your analysis, requiring you to click a few more buttons in SPSS Statistics when performing your analysis, as well as think a little bit more about your data, but it is not a difficult task.

Before we introduce you to these four assumptions, do not be surprised if, when analysing your own data using SPSS Statistics, one or more of these assumptions is violated (i.e., is not met). This is not uncommon when working with real-world data rather than textbook examples, which often only show you how to carry out a dependent t-test when everything goes well! However, don't worry. Even when your data fails certain assumptions, there is often a solution to overcome this. First, let's take a look at these four assumptions:

- **Assumption #1:** Your **dependent variable** should be measured on a **continuous** scale (i.e., it is measured at the **interval** or **ratio** level). Examples of variables that meet this criterion include revision time (measured in hours), intelligence (measured using IQ score), exam performance (measured from 0 to 100), weight (measured in kg), and so forth. You can learn more about continuous variables in our article: [Types of Variable](#).
- **Assumption #2:** Your **independent variable** should consist of **two categorical**, "**related groups**" or "**matched pairs**". "Related groups" indicates that the same subjects are present in both groups. The reason that it is possible to have the same subjects in each group is because each subject has been measured on two occasions on the same dependent variable. For example, you might have measured 10 individuals' performance in a spelling test (the dependent variable) before and after they underwent a new form of computerised

teaching method to improve spelling. You would like to know if the computer training improved their spelling performance. The first related group consists of the subjects at the beginning of (prior to) the computerised spelling training and the second related group consists of the same subjects, but now at the end of the computerised training. The dependent t-test can also be used to compare different subjects, but this does not happen very often. Nonetheless, to learn more about the different study designs that can be analysed using a dependent t-test, see our enhanced dependent t-test guide.

- **Assumption #3:** There should be **no significant outliers** in the **differences** between the two related groups. Outliers are simply single data points within your data that do not follow the usual pattern (e.g., in a study of 100 students' IQ scores, where the mean score was 108 with only a small variation between students, one student had a score of 156, which is very unusual, and may even put her in the top 1% of IQ scores globally). The problem with outliers is that they can have a negative effect on the dependent t-test, reducing the validity of your results. In addition, they can affect the statistical significance of the test. Fortunately, when using SPSS Statistics to run a dependent t-test on your data, you can easily detect possible outliers. In our enhanced dependent t-test guide, we (a) show you how to use SPSS Statistics to compute the difference scores, (b) show you how to detect outliers using SPSS Statistics, and (c) discuss some of the options you have in order to deal with outliers.
- **Assumption #4:** The **distribution of the differences** in the **dependent variable** between the two related groups should be **approximately normally distributed**. We talk about the dependent t-test only requiring **approximately** normal data because it is quite "robust" to violations of normality, meaning that the assumption can be a little violated and still provide valid results. You can test for normality using the Shapiro-Wilk test of normality, which is easily tested for using SPSS Statistics. In addition to showing you how to do this in our enhanced dependent t-test guide, we also explain what you can do if your data fails this assumption (i.e., if it fails it more than a little bit).

You can check assumptions #3 and #4 using SPSS Statistics. Before doing this, you should make sure that your data meets assumptions #1 and #2, although you don't need SPSS Statistics to do this. When moving on to assumptions #3 and #4, we suggest testing them in this order because it represents an order where, if a violation to the assumption is not correctable, you will no longer be able to use a dependent t-test (although you may be able to run another statistical test on your data instead). Just remember that if you do not run the statistical tests on these assumptions correctly, the results you get when running a dependent t-test might not be valid. This is why we dedicate a number of sections of our enhanced dependent t-test guide to help you get this right. You can find out about our enhanced content as a whole on our Features: **Overview** page, or more specifically, learn how we help with testing assumptions on our Features: **Assumptions** page.

In the section, Test Procedure in SPSS Statistics, we illustrate the SPSS Statistics procedure required to perform a dependent t-test assuming that no assumptions have been violated. First, we set out the example we use to explain the dependent t-test procedure in SPSS Statistics.

Get access to all the enhanced SPSS guides in Laerd Statistics



“ Thanks again for making such a helpful website and making a subscription completely affordable

”

Darlynda, USA

“ I just signed up for Laerd Statistics and must say that your website is just what I needed! Your guides are very detailed and easy to follow, which is just perfect!

”

Anne, Denmark

TAKE THE TOUR

PLANS & PRICING

SPSS Statistics

Example


A group of Sports Science students ($n = 20$) are selected from the population to investigate whether a 12-week plyometric-training programme improves their standing long jump performance. In order to test whether this training improves performance, the students are tested for their long jump performance before they undertake a plyometric-training programme and then again at the end of the programme (i.e., the dependent variable is "standing long jump performance", and the two related groups are the standing long jump values "before" and "after" the 12-week plyometric-training programme).

SPSS Statistics

Test Procedure in SPSS Statistics


The six steps below show you how to analyse your data using a dependent t-test in SPSS Statistics when the four assumptions in the previous section, Assumptions, have not been violated. At the end of these six steps, we show you how to interpret the results from this test. If you are looking for help to make sure your data meets assumptions #3 and #4, which are required when using a dependent t-test, and can be tested using SPSS Statistics, you can learn how we help with testing assumptions on our Features: **Assumptions** page. We also show you how to correctly enter your data into SPSS Statistics in order to run a dependent t-test, as well as explaining how to deal with missing values (e.g., if a participant completed a pre-test but failed to turn up to the post-test). However, in this "quick start" guide, we focus on the six steps required to run the dependent t-test procedure using SPSS Statistics.

- Click **Analyze > Compare Means > Paired-Samples T Test...** on the top menu, as shown below:





Published with written permission from SPSS Statistics, IBM Corporation.

- You will be presented with the **Paired-Samples T Test** dialogue box, as shown below:



Published with written permission from SPSS Statistics, IBM Corporation.

- Transfer the variables JUMP1 and JUMP2 into the Paired Variables: box. There are two ways to do this: (a) click on both variables whilst holding down the shift key (which highlights them) and then pressing the  button; or (b) drag-and-drop each variable separately into the boxes. If you are using older versions of SPSS Statistics, you will need to transfer the variables using the former method. You will end up with a screen similar to the one shown below:



Published with written permission from SPSS Statistics, IBM Corporation.

Note:



button shifts the pair of variables you have highlighted down one level.




button shifts the pair of variables you have highlighted up one level.



button shifts the order of the variables within a variable pair.

- If you need to change the confidence level limits or exclude cases, click on the **Options...** button. You will be presented with the **Paired-Samples T Test: Options** dialogue box, as shown below:



Published with written permission from SPSS Statistics, IBM Corporation.

- Click on the **Continue** button. You will be returned to the **Paired-Samples T Test** dialogue box.
- Click on the **OK** button.

Join the 10,000s of students, academics and professionals who rely on Laerd Statistics. **TAKE THE TOUR PLANS & PRICING**

SPSS Statistics

Output of the Dependent T-Test in SPSS Statistics

SPSS Statistics generates three tables in the **Output Viewer** under the title "T-Test", but you only need to look at two tables: the **Paired Samples Statistics** table and the **Paired Samples Test** table. In addition, you will need to interpret the boxplots that you created to check for outliers and the output from the Shapiro-Wilk test for normality, which you used to determine whether the distribution of the differences in the dependent variable between the two related groups were approximately normally distributed. This is explained in our enhanced guide. However, in this "quick start" guide, we focus on the two main tables you need to understand if your data has met all the necessary assumptions:

Paired Sample Statistics Table

The first table, titled **Paired Samples Statistics**, is where SPSS Statistics has generated descriptive statistics for your variables. You could use the results here to describe the characteristics of the first and second jumps in your write-up.

Paired Samples Statistics

		Mean	N	Std. Deviation	Std. Error Mean
Pair 1	JUMP1	2.4815	20	.16135	.03608
	JUMP2	2.5155	20	.15982	.03574

Published with written permission from SPSS Statistics, IBM Corporation.

Paired Samples Test Table

The **Paired Samples Test** table is where the results of the dependent t-test are presented. A lot of information is presented here and it is important to remember that this information refers to the **differences** between the two jumps (the subtitle reads "Paired Differences"). As such, the columns of the table labelled "**Mean**", "**Std. Deviation**",

"Std. Error Mean" and "95% Confidence Interval of the Difference" refer to the mean difference between the two jumps and the standard deviation, standard error and 95% confidence interval of this mean difference, respectively. The last three columns express the results of the dependent t-test, namely the *t*-value ("**t**"), the degrees of freedom ("**df**") and the significance level ("**Sig. (2-tailed)**").

Paired Samples Test

	Paired Differences					t	df
	Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference			
				Lower	Upper		
Pair 1 JUMP1 - JUMP2	-.03400	.03185	.00712	-.04891	-.01909	-4.773	19

Published with written permission from SPSS Statistics, IBM Corporation.

You are essentially conducting a one-sample t-test on the differences between the groups.

SPSS Statistics

Reporting the Output of the Dependent T-Test

You might report the statistics in the following format: $t(\text{degrees of freedom}) = t\text{-value}$, $p = \text{significance level}$. In our case this would be: $t(19) = -4.773$, $p < 0.0005$. Due to the means of the two jumps and the direction of the *t*-value, we can conclude that there was a statistically significant improvement in jump distance following the plyometric-training programme from 2.48 ± 0.16 m to 2.52 ± 0.16 m ($p < 0.0005$); an improvement of 0.03 ± 0.03 m.

Note: SPSS Statistics can output the results to many decimal places, but you should understand your measurement scale in order to know whether it is appropriate to report your results with such precision.

In our enhanced dependent t-test guide, we show you how to write up the results from your assumptions tests and dependent t-test procedure if you need to report this in a dissertation/thesis, assignment or research report. We do this using the Harvard and APA styles. It is also worth noting that in addition to reporting the results from your assumptions and dependent t-test, you are increasingly expected to report effect sizes. Whilst there are many different ways you can do this, we show you how to calculate effect sizes from your SPSS Statistics results in our enhanced dependent t-test guide. Effect sizes are important because whilst the dependent t-test tells you whether differences between group means are "real" (i.e., different in the population), it does not tell you the "size" of the difference. Providing the effect size in your results helps to overcome this limitation. You can learn more about our enhanced content on our Features: **Overview** page.

Join the 10,000s of students, academics and professionals who rely on Laerd Statistics. **TAKE THE TOUR PLANS & PRICING**

1

Home About Us Contact Us Terms & Conditions Privacy & Cookies © 2018 Lund Research Ltd