

Subtype-dependent LOX-family transcript changes in aging murine thymic stroma

Author: Aliaksandr Karatseyeu

Affiliation: Independent researcher

Correspondence: <https://github.com/G1F12/ThymusLOXScan>

Version: v5.6 GSE231906 human aging context pilot

Abstract

The lysyl oxidase (LOX) family has been implicated in extracellular matrix remodeling and aging in multiple tissues, but its role in thymic stromal aging remains unclear. Here, we performed a computational reanalysis of the public GSE240016 single-cell RNA-seq dataset, comprising 22,932 CD45-negative thymic stromal cells from young (2-month) and aged (18-month) female C57BL/6 mice. Pseudobulk and sample-level summaries showed subtype-dependent LOX-family transcript differences rather than a uniform fibroblast-wide pattern. Capsular fibroblasts showed lower *Lox* in aged samples ($\log_2FC = -1.46$, $p_{adj} = 7.5e-5$), while medullary fibroblasts showed divergent changes, with *Lox11* increased and *Lox12* decreased in aged samples. In the mTEC1 epithelial compartment, *Lox12* showed an aged-lower pseudobulk estimate and very low aged-cell detection, but this comparison included only two young and two aged biological samples. In a very small fibroblast composition-adjusted sensitivity model, the age coefficients for *Lox*, *Lox11*, and *Lox12* remained aged-lower, but the model had only six samples and one residual degree of freedom. Detection-rate decomposition suggested that several findings, especially medullary fibroblast and mTEC1 *Lox12*, largely reflected fewer expressing cells. Focused dropout/depth sensitivity analysis supported the aged-lower mTEC1 *Lox12* direction after depth-matched checks but classified dropout/depth artifact risk as partial/unresolved; public protein/spatial resources provided weak indirect context only, leaving direct LOXL2 protein validation to future IHC or related assays. Annotation sanity checks found expected marker expression to varying degrees in capsFB, intFB, medFB, and mTEC1 labels, and the four focal transcript directions had the same sign in stricter marker-positive subsets. External comparisons provided broad directional context only. GSE223049 sorted bulk RNA-seq was directionally consistent with broad aged-lower thymic fibroblast *Lox* and *Lox12* and broad thymic epithelial *Lox12*. E-MTAB-8560 TEC Smart-seq2 data showed an aged-lower *Lox12* tendency in broad TEC and mTEC-like groups across the first year of mouse life, but not exact mTEC1 recapitulation. A controlled GSE231906 pilot made aged-human thymus LOX-family transcript context technically evaluable and recovered LOX-family genes plus TEC/mTEC-like compartments after metadata-to-expression joining, while the LOXL2 mTEC-like aging context remained mixed. Together, these analyses nominate subtype-dependent LOX-family transcript differences as a hypothesis-generating feature of murine thymic stromal aging, with *Lox12* emerging as a recurrent candidate transcript marker requiring orthogonal experimental follow-up.

Introduction

Age-associated thymic involution is accompanied by reduced thymic output and changes in the thymic stromal microenvironment, including epithelial and mesenchymal compartments that support T cell development [1-4]. Recent single-cell and spatial transcriptomic analyses of murine thymic aging identified age-associated epithelial states and stromal changes in public datasets [5,6]. These public data provide an opportunity to examine additional stromal gene programs in a transparent reanalysis.

The lysyl oxidase family, including LOX and LOX-like enzymes LOXL1-4, encodes secreted copper-dependent amine oxidases involved in extracellular matrix biology, particularly collagen and elastin crosslinking [7,8]. LOX-family genes have also been linked to tissue remodeling, fibrosis, cancer-associated matrix organization, and age-related extracellular matrix changes in multiple contexts [9,10]. Individual family members can have distinct biological associations: LOXL1 is required for elastic fiber homeostasis [11], while LOXL2 has been implicated in collagen IV crosslinking, basement membrane organization, and epithelial or endothelial tissue remodeling [12,13].

Here we reanalyzed the public GSE240016 murine thymic stromal single-cell RNA-seq dataset to assess LOX-family expression in young and aged CD45-negative thymic stromal cells [5,6]. We used pseudobulk differential expression with biological samples as replicates as the primary inferential framework, with descriptive single-cell summaries, per-sample checks, and correlation analyses as supporting context. The goal was not to prove a mechanism, but to identify candidate stromal expression patterns that may motivate future experimental testing and independent subtype-resolved reanalysis.

Results

Dataset and analysis overview

We analyzed the public GSE240016 annotated single-cell RNA-seq dataset of CD45-negative murine thymic stromal cells from young (02mo) and aged (18mo) mice at steady state [5,6]. The repository input file was `data/raw/GSE240016_CD45neg_thymic_stroma_d0+annotation.h5ad`, corresponding to the GEO supplementary file named `GSE240016_CD45neg_thymic_stroma_d0+annotation.h5ad`. The analyzed local file contained 22,932 cells and metadata fields for age, biological sample, broad cell type, and fine subtype. Downstream analyses focused on the LOX-family genes *Lox*, *Lox11*, *Lox12*, *Lox13*, and *Lox14*.

The main inferential analysis used pseudobulk DESeq2, in which raw counts were summed within each biological sample and cell type or subtype before age comparison. Major pseudobulk results are reported in `results/tables/lox_pseudobulk_complete_results.tsv` and Supplementary Table 2. Cell counts by sample and annotation are reported in Supplementary Table 1. Descriptive single-cell tests are reported in `results/sc_mannwhitney_FB_combined.csv`, `results/sc_mannwhitney_mTEC1.csv`, and Supplementary Table 3. Correlation results are reported in `results/sc_spearman_correlations.csv` and Supplementary Table 4.

Broad fibroblast summaries show reduced LOX-family expression but are composition-sensitive

At the broad fibroblast level, pseudobulk DESeq2 estimated lower expression of several LOX-family genes in aged samples. *Lox* was lower in aged fibroblasts ($\log_2FC = -1.29$, $padj = 4.36e-5$), as were *Lox11* ($\log_2FC = -0.62$, $padj = 0.014$) and *Lox12* ($\log_2FC = -0.52$, $padj = 0.021$). These broad fibroblast-level pseudobulk results used three young and three aged biological samples and are shown in Figure 1 and `results/tables/lox_pseudobulk_complete_results.tsv`. However, broad fibroblast summaries pool capsular, interstitial, medullary, and fat-associated fibroblast states, so they are not treated here as the central biological conclusion.

Descriptive single-cell tests showed the same general direction for all five LOX-family genes in pooled fibroblasts, with lower expression in aged cells for *Lox*, *Lox11*, *Lox12*, *Lox13*, and *Lox14*. The largest rank-biserial effect sizes were observed for *Lox11* (-0.226), *Lox* (-0.154), and *Lox12* (-0.131), whereas *Lox14* had a negligible effect size (-0.004) despite nominal statistical significance. These single-cell tests are descriptive because cells are not independent biological replicates. Follow-up sample-level sensitivity analysis found that broad fibroblast age coefficients had the same aged-lower sign after adjustment for capsFB, intFB, medFB, and Fat fractions, but with only six samples and one residual degree of freedom in the adjusted model. Thus, reduced broad fibroblast expression is best presented as partly influenced by subtype composition and as compartment-level context rather than primary evidence for the subtype-resolved findings. Sources: `results/sc_mannwhitney_FB_combined.csv`, Supplementary Table 3, `results/tables/fb_composition_adjusted_lox.tsv`, and `reports/fb_composition_adjustment.md`.

Fibroblast subtypes show divergent LOX-family age-associated patterns

Subtype-resolved pseudobulk analysis provided the main fibroblast transcript summaries and suggested that LOX-family age-associated patterns were not uniform across annotated fibroblast populations. Capsular fibroblasts showed reduced *Lox* in aged samples ($\log_2FC = -1.46$, $padj = 7.5e-5$; three young and three aged biological samples). This result is shown in Figure 1 and Figure 2A-B and reported in `results/tables/lox_pseudobulk_complete_results.tsv`.

Medullary fibroblasts showed divergent isoform behavior. *Lox11* was higher in aged medullary fibroblasts ($\log_2FC = 0.75$, $padj = 0.023$), whereas *Lox12* was lower ($\log_2FC = -1.00$, $padj = 0.0067$). Both medullary fibroblast comparisons used three young and three aged biological samples. Detection-rate summaries suggested different expression architectures for these findings: medFB *Lox11* had higher aged detection but lower mean expression among detecting cells, whereas medFB *Lox12* was more consistent with fewer detecting cells in aged samples. These results show a subtype-dependent LOX-family transcript pattern rather than a single pooled-fibroblast decrease, but they do not establish functional extracellular matrix changes. Sources: Figure 1, Figure 2A-B, `results/tables/lox_pseudobulk_complete_results.tsv`, `results/tables/lox_detection_rates_by_sample.tsv`, `reports/lox_detection_rate_analysis.md`, and Supplementary Table 2.




Figure 1. LOX-family pseudobulk differential expression

mTEC1 Loxl2 shows aged-lower expression and low aged detection in a n=2 vs n=2 comparison

An epithelial LOX-family candidate in this analysis was aged-lower Loxl2 in mTEC1. Pseudobulk DESeq2 estimated lower Loxl2 expression in aged mTEC1 samples ($\log_2\text{FC} = -3.29$, $\text{padj} = 9.6\text{e-}4$). This comparison included two young and two aged biological samples, with 579 young and 1,053 aged mTEC1 cells contributing to pseudobulk profiles in the complete results table. Because $n=2$ per age group limits inference, this result should be treated as a candidate mTEC1-associated transcript pattern requiring further independent testing rather than as functional evidence. Source: Figure 1, Figure 2A-B, [results/tables/lox_pseudobulk_complete_results.tsv](#), [results/tables/mtec1_loxl2_per_sample_expression.tsv](#), and [reports/mtec1_loxl2_robustness.md](#).

Per-sample inspection showed that both young mTEC1 pseudobulk samples had higher normalized Loxl2 expression than both aged mTEC1 samples. Detection-rate analysis indicated that the mTEC1 Loxl2 decrease primarily reflected fewer detecting cells in aged samples, with very low aged detection. A formal leave-one-sample-out DESeq2 refit was not appropriate because omitting one sample would leave only one biological replicate in one age group. The per-sample plot is available at [results/figures/per_sample/mtec1_loxl2_per_sample.pdf](#).

A focused dropout/depth audit found that young mTEC1 samples had higher Loxl2 detection than aged mTEC1 samples, and the young-higher/aged-lower direction persisted in pairwise sample comparisons and depth-matched downsampling. However, aged mTEC1 samples had lower sequencing depth and Loxl2 detection was sparse, so dropout/depth artifact was classified as partial/unresolved rather than excluded. Sources: [reports/mtec1_loxl2_dropout_depth_audit.md](#), [results/tables/mtec1_loxl2_dropout_depth_audit.tsv](#), [results/tables/mtec1_loxl2_pairwise_sample_direction.tsv](#), and [results/tables/mtec1_loxl2_downsampling_summary.tsv](#).

Sensitivity analyses and unresolved technical concerns

Several sensitivity analyses were added to clarify how much confidence should be placed in the highlighted LOX-family directions. In a fibroblast subtype-composition sensitivity model, broad fibroblast age coefficients for Lox, Lox11, and Lox12 remained aged-lower after adjustment for capsFB, intFB, medFB, and Fat fractions. However, this model used only six broad fibroblast samples and included age plus four subtype-fraction covariates, leaving one residual degree of freedom. The adjusted coefficients are therefore best interpreted as directional sensitivity checks rather than definitive evidence for broad fibroblast-intrinsic age effects.

Detection-rate decomposition suggested that several transcript differences coincided partly or largely with fewer cells having detectable counts for the relevant gene. This was especially clear for medFB Lox12, where detection was lower in aged samples, and for mTEC1 Lox12, where aged samples showed very low detection. capsFB Lox and pooled fibroblast Lox reflected both fewer detecting cells and lower expression among detecting cells, while medFB Lox11 behaved differently, with higher aged detection but lower nonzero expression among detecting cells. These patterns favor presenting detection and expression as linked but non-identical summaries.

The highlighted directions had the same sign in quality-control threshold checks, raw-count CPM summaries, normalized-expression summaries, exclusion of sorted/enriched sample preparations, and leave-one-sample-out direction checks where feasible. These checks show that the reported directions are not unique to one normalization representation, one QC filter, or one obvious outlier sample. They do not eliminate batch or sample-preparation confounding, sequencing-depth differences, dropout, low-count instability, or annotation uncertainty. Sources: `reports/hypothesis_falsification.md`, `reports/fb_composition_adjustment.md`, `results/tables/fb_composition_adjusted_lox.tsv`, `reports/lox_detection_rate_analysis.md`, and `results/tables/lox_detection_rates_by_sample.tsv`.

Annotation uncertainty and stricter marker-positive subsets

Because the central claims depend on published fine annotations, we performed a marker-based annotation sanity check for capsFB, intFB, medFB, and mTEC1. The marker sets included capsFB markers `Dpp4`, `Smpd3`, `Pi16`, and `Pdgfra`; intFB markers `Inmt`, `Gpx3`, and `Pdgfra`; medFB markers `Ptn`, `Postn`, and `Pdgfra`; and mTEC1 markers `Cc121a`, `Itgb4`, `Ly6a`, `Epcam`, `H2-Aa`, `Krt8`, and `Krt5`. Marker expression was more consistent with the expected medFB and mTEC1 labels than with capsFB and intFB labels, where marker support was partial, especially in aged samples.

All four focal LOX-family directions had the same sign in stricter marker-positive subsets: capsFB Lox remained lower in aged samples, medFB Lox11 remained higher, medFB Lox12 remained lower, and mTEC1 Lox12 remained lower. In strict marker-positive subsets, the aged-minus-young $\Delta \log_2(\text{CPM}+1)$ values were -1.974 for capsFB Lox, +1.036 for medFB Lox11, -0.958 for medFB Lox12, and -2.657 for mTEC1 Lox12. These findings make a simple marker-threshold artifact less obvious under this filtering strategy, but they do not prove that the published annotations are correct or that annotation uncertainty is fully resolved. Sources: `reports/annotation_uncertainty_check.md`, `results/figures/annotation_sanity/strict_marker_positive_lox_age_summary.tsv`, and `results/figures/annotation_sanity/strict_marker_positive_lox_summary.png`.

Independent external comparison in GSE223049

We next used GSE223049, an independent mouse sorted bulk RNA-seq dataset comparing young 2-month and aged 22-24-month samples across sorted immune and stromal cell types. The relevant external populations were broad Thymic_fibroblasts and broad Thymic_epithelial samples. This sorted bulk RNA-seq design cannot test capsFB, medFB, or mTEC1 specificity, and the aged group is older than the 18-month group in GSE240016. It can nevertheless provide a descriptive check of whether broad thymic fibroblast and broad thymic epithelial LOX-family directions are similar outside the original single-cell dataset.

In GSE223049 broad thymic fibroblasts, mean Lox was lower in aged samples, with $\Delta \log_2(\text{CPM}+1) = -0.354$, and mean Lox12 was lower, with $\Delta \log_2(\text{CPM}+1) = -1.206$. Broad thymic fibroblast Lox11 showed only a weak decrease, $\Delta \log_2(\text{CPM}+1) = -0.112$, and therefore did not test or replicate the medFB-specific Lox11 increase observed in GSE240016. In GSE223049 broad thymic epithelial samples, mean Lox12 was lower in aged samples, with $\Delta \log_2(\text{CPM}+1) = -0.588$. Thus, GSE223049 sorted bulk RNA-seq is directionally similar to aged-lower broad fibroblast Lox/Lox12 and broad epithelial Lox12, especially Lox12, but it does not test exact fibroblast-subtype or mTEC1-specific claims and is not subtype-resolved evidence. Sources: `results/tables/external_gse223049_lox_validation.tsv`, `results/tables/external_gse223049_lox_validation_summary.tsv`, `reports/external_dataset_search.md`, `reports/external_validation_plan.md`, and `reports/highest_impact_next_analysis.md`.

Cross-dataset external comparisons

We next assembled a cross-dataset comparison matrix across the internal GSE240016 pseudobulk results, GSE223049 sorted bulk RNA-seq, E-MTAB-8560 TEC Smart-seq2 age-series summaries, and GSE231906 human thymus metadata. Each row was assigned to a specific claim mapping, including broad fibroblast Lox aged-lower, broad fibroblast Lox12 aged-lower, capsFB Lox aged-lower, medFB Lox11 aged-higher, medFB Lox12 aged-lower, epithelial Lox12 aged-lower, and mTEC1 Lox12 aged-lower. A simple consistency score was used only as a directional summary: +1 for an appropriate external row with the same direction, 0 for not testable or inconclusive rows, and -1 for an appropriate external row in the opposite direction. Broad sorted bulk rows were not counted as subtype-specific evidence.

The cross-dataset matrix provided descriptive directional context for three broad or appropriately resolved transcript-level comparisons. First, broad fibroblast Lox aged-lower and broad fibroblast Lox12 aged-lower were directionally similar between internal GSE240016 summaries and GSE223049 broad thymic fibroblast bulk RNA-seq. Second, epithelial Lox12 aged-lower was directionally

similar across GSE240016, GSE223049 broad thymic epithelial samples, and E-MTAB-8560 broad TEC age-series summaries. Third, the mTEC1 Lox12 direction had approximate mTEC-like directional context from E-MTAB-8560 mTEC-like, mTEClo, and mTEChi summaries, but this was not exact mTEC1 recapitulation. In contrast, capsFB Lox aged-lower, medFB Lox11 aged-higher, and medFB Lox12 aged-lower remained internally observed or externally not testable at exact subtype resolution. The later controlled GSE231906 pilot made aged-human LOX-family transcript context technically evaluable, but the LOXL2 mTEC-like context was mixed and is not used as a validation or conservation claim. No claim was classified as contradicted by an appropriate external dataset row. Sources: results/tables/cross_dataset_lox_validation_matrix.tsv, results/tables/cross_dataset_lox_consistency_summary.tsv, reports/cross_dataset_lox_analysis.md, and reports/cross_dataset_figure_notes.md.

A later E-MTAB-8560 R/Bioconductor export and SDRF metadata join provided per-mouse summaries for external transcript-level context. E-MTAB-8560 provides independent mTEC-focused transcript-level context, with the clearest aged-lower Lox12 pattern in mTEClo and cTEC, weaker support in mTEChi, and overall mixed/inconclusive classification. This does not constitute exact GSE240016 mTEC1 validation.

Controlled GSE231906 human thymus aging pilot

GSE231906 was analyzed because prior human TEC/epithelial datasets provided developmental or metadata-limited context, whereas GSE231906 offered a possible aged-human thymus dataset with donor/sample labels and stromal/TEC annotations. The goal was to test whether aged-human LOX-family transcript summaries could be made technically evaluable under controlled data-handling constraints, not to test conservation or validate the mouse GSE240016 mTEC1 Lox12 result.

The controlled pilot was technically usable. The metadata workbook was parsed with 516,352 metadata rows. The locally downloaded processed archive was audited without committing large data; the manifest contained 217 archive members and 59 candidate expression units. Pilot extraction completed for GSM8945751_donor3, the metadata join was successful, and the barcode-to-metadata join match rate was 0.909477.

LOX-family genes LOX, LOXL1, LOXL2, LOXL3, and LOXL4 were present. Recoverable target compartments included epithelial/Epi, mTEC-like, cTEC-like, post_AIRE_mTEC, immature TEC, mesenchymal/fibroblast-like, and endothelial contexts. Donor/sample-aware aging context was feasible, detection summaries were generated, and descriptive mean-value summaries were generated only as descriptive summaries because expression matrix semantics remain partial.

The LOXL2 mTEC-like aging context was mixed, with evidence grade mixed_transcript_level_context. The controlled GSE231906 pilot makes aged-human thymus LOX-family transcript context technically evaluable. LOX-family genes and TEC/mTEC-like compartments were recoverable after metadata-to-expression joining, but the LOXL2 mTEC-like aging context was mixed. We therefore interpret GSE231906 as mixed aged-human transcript-level context, not human validation or conservation of the mouse GSE240016 mTEC1 Lox12 candidate signal.

This result strengthens the rationale for orthogonal mouse LOXL2 IHC/IF or Lox12 RNA-ISH/RNAscope by showing that aged-human context is evaluable but not definitive. It does not establish human conservation, exact replication, mechanism, protein-level support, LOX activity, extracellular matrix crosslinking, thymic function, therapeutic relevance, or causality.

dataset	age/development relevance	metadata strength	LOX gene coverage	TEC/mTEC-like coverage	matrix semantics	evidence grade	interpretation
Park TEC	Developmental; not aging-focused	Donor/sample-aware	LOX-family detection context	TEC subtypes	X detection; no means	exploratory_transcript_level_context	Developmental TEC context only
Yayon TEC	Developmental/early-life; not aging-focused	Donor/sample-aware	LOX-family detection context	TEC subtypes	X detection; no means	exploratory_transcript_level_context	Early-life TEC context only
GSE147520 epithelial	Fetal/postnatal/adult-labeled; limited adult coverage	Sample-aware; donor/sex unavailable	LOX-family detection from raw.X	Epithelial context	raw.X detection; no means	metadata_limited_transcript_level_context	Sample-aware epithelial context only
GSE231906	Aged-human thymus context	Donor/sample-aware joined metadata	LOX, LOXL1, LOXL2, LOXL3, LOXL4 present	Epi, mTEC-like, cTEC-like, post_AIRE_mTEC, immature TEC, mesenchymal/fibroblast-like, endothelial	Partial; descriptive means only	mixed_transcript_level_context	Mixed LOXL2 mTEC-like context; not human validation or conservation

Fibroblast Loxl2 correlates with Col1a1 and Vim but not Snai1

In pooled fibroblasts, per-cell Spearman correlations showed weak positive associations between Loxl2 and structural or mesenchymal markers Col1a1 ($\rho = 0.193$, $p = 3.2\text{e-}111$) and Vim ($\rho = 0.145$, $p = 8.0\text{e-}63$). Loxl2 was not meaningfully correlated with Snai1 ($\rho = 0.005$, $p = 0.578$). These descriptive correlations are consistent with Loxl2 co-varying weakly with selected structural or mesenchymal transcript markers more than with Snai1, but they do not establish pathway direction or protein-level co-regulation. Source: Figure 2C, results/sc_spearman_correlations.csv, and Supplementary Table 4.

LOXL2 protein and spatial public-resource feasibility

An in-silico LOXL2 protein/spatial feasibility audit found weak indirect public support. Human thymus spatial RNA and epithelial RNA resources provided context for LOXL2 detectability in thymic epithelial/medullary settings, but public protein resources did not provide direct mouse mTEC1 LOXL2 protein localization or age-aware thymic epithelial protein evidence. These results do not substitute for IHC. Sources: reports/loxl2_protein_spatial_validation_feasibility.md, results/tables/loxl2_protein_spatial_evidence_audit.tsv, and results/tables/loxl2_ihc_antibody_feasibility.tsv.




Figure 2. LOX-family expression changes and descriptive correlations

Methods

Dataset and preprocessing

Single-cell RNA-seq data were analyzed from the public murine thymic stromal dataset GSE240016 [5,6]. The primary input file used by the repository was data/raw/GSE240016_CD45neg_thymic_stroma_d0+annotation.h5ad. GEO lists this processed supplementary file at:

https://www.ncbi.nlm.nih.gov/geo/download/?acc=GSE240016&format=file&file=GSE240016_CD45neg_thymic_stroma_d0%2Bannotation.h5ad

The local SHA256 checksum of the analyzed file was:

A5CF4BAEB408412947DE2AE1867B4AD42271088C34C9DD212479C8B19E119FB1

Metadata fields used by the analysis included `stage`, `sample`, `cell_type`, `cell_type_subset`, `n_genes_by_counts`, `total_counts`, and `mito_frac`. The age comparison was encoded as 02mo versus 18mo.

Preprocessing was performed in `notebooks/03_preprocessing_thymus.ipynb`. Gene names were made unique, `stage` was copied to `age_group`, and quality-control distributions were inspected. Cells were retained with `n_genes_by_counts >= 200`, `n_genes_by_counts <= 6000`, and `mito_frac <= 0.20`; genes were retained with `sc.pp.filter_genes(..., min_cells=3)`. The filtered object was written to `data/processed/thymus_preprocessed.h5ad`. Existing UMAP coordinates and published annotation fields were preserved for visualization.

Cell type annotations

Cell type labels were not newly inferred in this repository. The workflow retained the published annotations present in the GSE240016 processed AnnData object and inspected them in `notebooks/04_cell_type_annotation.ipynb` [5,6]. Broad labels included fibroblasts (FB), endothelial cells (EC), thymic epithelial cells (TEC), vascular smooth muscle/pericyte-like cells, mesenchymal epithelial cells, and non-myelinating Schwann cells. Fine subtype labels included 3:capsFB, 4:intFB, 5:medFB, and 13:mTEC1. The annotated object was written to `data/processed/thymus_annotated.h5ad`.

Pseudobulk differential expression

The primary inferential analysis was pseudobulk DESeq2 implemented in `scripts/make_pseudobulk_results_table.py` and related code in `scripts/pseudobulk_deseq2_lox.py`. Raw counts from `adata.raw.X` were summed within each biological sample and cell type or subtype. Sample-by-annotation groups with fewer than 10 cells were excluded. Biological samples, not individual cells, were used as replicates, following recommendations from pseudobulk and multi-sample single-cell differential expression literature [14-16].

DESeq2 modeling was performed with PyDESeq2 using design `~ stage`, reference level 02mo, and contrast 18mo versus 02mo [17,18]. Positive log2 fold changes therefore indicate higher expression in aged samples. Results were extracted for `Lox`, `Lox11`, `Lox12`, `Lox13`, and `Lox14`, including `baseMean`, `log2FoldChange`, `lfcSE`, Wald statistic, nominal p-value, and Benjamini-Hochberg adjusted p-value. The complete current result tables were written to `results/tables/lox_pseudobulk_complete_results.tsv` and `results/tables/lox_pseudobulk_complete_results.csv`.

Descriptive single-cell tests

Single-cell-level tests were used descriptively rather than as the primary inferential framework. Normalized per-cell expression values from `adata.X` were compared between age groups with two-sided Mann-Whitney U tests, and rank-biserial effect sizes were calculated. Current manuscript single-cell summaries are stored in `results/sc_mannwhitney_FB_combined.csv` and `results/sc_mannwhitney_mTEC1.csv`, and are exported in Supplementary Table 3. Because these files are derived outputs without a standalone generator script identified in the current repository, they are used only as descriptive supporting summaries.

Fibroblast subtype-composition sensitivity analysis

To perform a limited directional sensitivity check for fibroblast subtype mixture, sample-level broad fibroblast pseudobulk expression was modeled with age while adjusting for capsFB, intFB, medFB, and Fat fractions within each biological sample. This analysis was implemented in `scripts/fb_composition_adjusted_lox.py` and written to `results/tables/fb_composition_adjusted_lox.tsv` with a narrative summary in `reports/fb_composition_adjustment.md`. Because the adjusted model used only six broad fibroblast samples and included age plus four subtype-fraction covariates, it was treated as a sensitivity analysis rather than definitive inference.

Detection-rate analysis

Cell-level detection rates were computed from raw counts in `adata.raw.X` with detection defined as raw count greater than zero. For each biological sample, selected cell type or subtype, and LOX-family gene, the analysis recorded cell count, summed raw counts, number of detecting cells, detection rate, and mean raw expression among detecting cells. This analysis was implemented in `scripts/lox_detection_rates_by_sample.py`, with outputs in `results/tables/lox_detection_rates_by_sample.tsv` and `reports/lox_detection_rate_analysis.md`.

mTEC1 Lox12 dropout/depth audit

The mTEC1 Lox12 dropout/depth audit was implemented in `scripts/mtec1_lox12_dropout_depth_audit.py`. The audit used raw counts from `adata.raw.X` for annotated 13:mTEC1 cells and summarized per-sample Lox12 detection rate, pseudobulk CPM,

total_counts, n_genes_by_counts, and mito_frac. It also performed pairwise young-versus-aged sample comparisons and depth-matched downsampling with a fixed random seed. A descriptive QC-adjusted logistic model was fit with depth/QC covariates, but cell-level models were treated as descriptive because cells are not independent biological replicates. The final dropout/depth artifact-risk classification was Partial / unresolved, reflecting directional persistence after these checks together with sparse detection, lower aged mTEC1 depth, and only two biological samples per age group. Outputs are stored in reports/mtec1_loxl2_dropout_depth_audit.md, results/tables/mtec1_loxl2_dropout_depth_audit.tsv, results/tables/mtec1_loxl2_pairwise_sample_direction.tsv, results/tables/mtec1_loxl2_downsampling_summary.tsv, and results/tables/mtec1_loxl2_depth_adjusted_logistic.tsv.

Marker-positive annotation sanity check

Annotation uncertainty was evaluated by checking marker expression within the published capsFB, intFB, medFB, and mTEC1 labels and by repeating key sample-level LOX-family summaries in stricter marker-positive subsets. Marker sets were taken from the original study's described annotation markers and included fibroblast subtype markers plus broad lineage markers. Strict marker-positive cells were summarized by biological sample, age, subtype, and focal LOX-family gene using $\log_2(\text{CPM}+1)$ and detection-rate summaries. This analysis was implemented in scripts/annotation_uncertainty_check.py, with narrative output in reports/annotation_uncertainty_check.md, tabular outputs in results/figures/annotation_sanity/marker_summary_by_sample.tsv, results/figures/annotation_sanity/marker_expression_long.tsv, results/figures/annotation_sanity/strict_marker_positive_lox_by_sample.tsv, and results/figures/annotation_sanity/strict_marker_positive_lox_age_summary.tsv, and supplementary marker-sanity figures in results/figures/annotation_sanity/.

GSE223049 external comparison

Independent external comparison used GSE223049, a mouse sorted bulk RNA-seq dataset with 2-month and 22-24-month samples across sorted cell types. The analysis focused on the sorted broad Thymic_fibroblasts and broad Thymic_epithelial populations. Counts were downloaded from GEO by scripts/external_validation_gse223049_lox.py, library-size normalized to CPM, transformed as $\log_2(\text{CPM}+1)$, and summarized by age group for Lox, Lox11, Lox12, Lox13, and Lox14. This was treated as a descriptive external comparison of broad cell-type directionality, not as subtype-resolved evidence. Outputs are stored in results/tables/external_gse223049_lox_validation.tsv, results/tables/external_gse223049_lox_validation_summary.tsv, and the local input directory data/external/GSE223049/.

E-MTAB-8560, GSE231906, and cross-dataset summaries

E-MTAB-8560 was analyzed as an external mouse TEC Smart-seq2 age-series using scripts/external_validation_emtab8560_tec_lox.py. The script parsed public ArrayExpress/BioStudies metadata and Bioconductor MouseThymusAgeing processed SMART-seq2 resources, summarized LOX-family expression by age in broad TEC, cTEC, mTEClo, mTEChi, and mTEC-like groups, and generated descriptive age-trend tables and figures. A later R/Bioconductor export joined official SDRF metadata to the MouseThymusAgeing colData, allowing per-mouse summaries for cTEC, mTEClo, mTEChi, and combined mTEClo/mTEChi context while retaining explicit batch limitations. Outputs are stored in results/tables/external_emtab8560_tec_lox_by_age.tsv, results/tables/external_emtab8560_tec_lox_summary.tsv, results/tables/external_emtab8560_mtec_lox_models.tsv, results/tables/external_emtab8560_mtec_lox_permutation.tsv, results/tables/external_emtab8560_mtec_lox_global_fdr.tsv, results/figures/external_validation/emtab8560/, reports/emtab8560_tec_external_validation.md, reports/external_emtab8560_biological_unit_audit.md, reports/external_emtab8560_mtec_lox_reanalysis.md, and reports/external_emtab8560_vs_gse240016_interpretation.md.

GSE231906 was handled with a guarded human metadata and expression-pilot workflow. An initial metadata-only pipeline in scripts/external_validation_gse231906_human_lox.py parsed the cell-level metadata workbook and identified candidate fibroblast-like/mesenchymal, epithelial, mTEC-like, cTEC-like, and endothelial groups. The v5.6 controlled pilot then used scripts/human_thymus_stage4_gse231906_metadata_audit.py, scripts/human_thymus_stage4_gse231906_archive_manifest.py, scripts/human_thymus_stage4_gse231906_pilot_extract.py, scripts/human_thymus_stage4_gse231906_join_pilot.py, scripts/human_thymus_stage4_gse231906_target_gene_extract.py, and scripts/human_thymus_stage4_gse231906_aging_context_summary.py to audit the large processed archive locally, extract one pilot expression unit, join barcodes to metadata, and generate donor/sample-aware LOX-family summaries. Expression matrix semantics remain partial, so mean-value summaries were treated as descriptive only and no cell-level statistical tests were run. Outputs are stored in results/tables/human_thymus_stage4_gse231906_*, reports/human_thymus_stage4_gse231906_balanced_report.md, reports/human_thymus_stage4_gse231906_decision_memo.md, and reports/human_thymus_stage4_gse231906_one_page_summary.md.

Cross-dataset summaries were built with `scripts/build_cross_dataset_lox_matrix.py`, which combines internal GSE240016, GSE223049, E-MTAB-8560, and GSE231906 outputs into a claim-level matrix. Direction consistency was scored as +1, 0, or -1 only for summary purposes. Figure panels were generated with `scripts/figures/plot_cross_dataset_lox_validation.py`. Outputs are stored in `results/tables/cross_dataset_lox_validation_matrix.tsv`, `results/tables/cross_dataset_lox_consistency_summary.tsv`, `results/figures/external_validation/cross_dataset/`, `reports/cross_dataset_lox_analysis.md`, and `reports/cross_dataset_figure_notes.md`.

LOXL2 protein/spatial feasibility audit

The LOXL2 protein/spatial feasibility audit checked public resources for direct or indirect support relevant to the epithelial/mTEC1 transcript finding. Resources included the Human Protein Atlas (HPA), ProteomicsDB, PRIDE/ProteomeXchange, CELLxGENE, UniProt, and antibody resources. The audit produced `results/tables/lox12_protein_spatial_evidence_audit.tsv` and `results/tables/lox12_ihc_antibody_feasibility.tsv`, with narrative summary in `reports/lox12_protein_spatial_validation_feasibility.md`. The final classification was weak indirect support. No direct mouse mTEC1 LOXL2 protein localization was found.

Correlation analysis

Fibroblast co-expression analyses used normalized per-cell expression values from `data/processed/thymus_annotated.h5ad`. Spearman correlations were calculated between `Lox12` and `Col1a1`, `Vim`, and `Snai1`, with results stored in `results/sc_spearman_correlations.csv` and Supplementary Table 4. These correlations were treated as descriptive because they were calculated at the single-cell level and do not model biological samples as independent replicates.

MAGIC imputation and exploratory analyses

MAGIC imputation was implemented in `scripts/magic_imputation_lox.py` as a sensitivity analysis for fibroblast LOX-family expression. The script used normalized fibroblast expression values, ran MAGIC with `knn=10` and `t=3`, and wrote results to `results/magic_imputation_LOX.csv`. The code also calculates Mann-Whitney statistics on imputed values; these p-values are a methodological concern because imputation can alter variance and dependence structure. MAGIC results should therefore be treated as visualization or qualitative sensitivity outputs only, not as primary inferential evidence.

Exploratory GSEA, signature scoring, and PAGA analyses are present in `notebooks/06_GSEA_TGFb_analysis.ipynb`, but the expected output files were not present in the current repository state. These analyses are not used as completed manuscript results in this draft.

Figure and table generation

Final manuscript figures were generated with `scripts/figures/plot_final_volcano.py` and `scripts/figures/plot_final_summary.py`. Outputs are stored in `results/figures/final/` as both PNG and PDF files. Per-sample robustness plots were generated with `scripts/plot_per_sample_lox_expression.py`. Supplementary tables were generated with `supplementary_tables/make_supplementary_tables.py`.

Software

The reproducible Python environment is specified in `environment.yml` and `requirements.txt`. The current workflow uses Python 3.11, Scanpy, AnnData, NumPy, pandas, SciPy, Matplotlib, seaborn, statsmodels, PyDESeq2, MAGIC, GSEAPy, python-igraph, and leidenalg. The primary GSE240016 pseudobulk analyses in this repository use Python/PyDESeq2 rather than an R/Bioconductor DESeq2 workflow. The E-MTAB-8560 external-context analysis uses an R/Bioconductor export path for `MouseThymusAgeing/SingleCellExperiment` resources, followed by derived per-mouse summaries and Python-based audit/reanalysis scripts.

Discussion

This computational reanalysis suggests a narrow central conclusion: age-associated LOX-family transcript differences in thymic stroma are subtype-dependent, with broad external directional context for selected fibroblast and epithelial directions. The largest internal fibroblast observations were reduced `Lox` in capsular fibroblasts, increased `Lox11` in medullary fibroblasts, and reduced `Lox12` in medullary fibroblasts. Broad fibroblast summaries also showed aged-lower `Lox`, `Lox11`, and `Lox12`, and these directions had the same sign in a composition-adjusted sensitivity model, but the small sample size and subtype-mixture changes mean these broad effects should not be treated as clean fibroblast-wide intrinsic changes.

At the transcript level, the observed pattern does not resemble a simple model in which all LOX-family genes increase uniformly across stromal compartments with age. Instead, the aged thymic stromal compartment showed a mixture of decreases and increases depending on annotated cell state and gene family member. One fibroblast example was the medullary fibroblast split between increased *Lox11* and decreased *Lox12*. This suggests that the annotated aged thymic stroma may contain altered proportions or expression levels of LOX-positive stromal states rather than a single monotonic LOX-family response across all fibroblasts.

Lox12 appeared repeatedly across the transcript-level summaries in this analysis. It appears in multiple contexts: broad fibroblast summaries, medullary fibroblast pseudobulk results, mTEC1 low-detection results, detection-rate decomposition, marker-positive annotation sanity checks, GSE223049 broad sorted bulk comparisons, and E-MTAB-8560 TEC age-series summaries. These observations make *Lox12* a recurrent candidate transcript marker of age-associated thymic stromal differences. They do not show that *Lox12* drives the phenotype, that LOXL2 protein is reduced, that LOX enzymatic activity changes, or that extracellular matrix crosslinking is altered.

The mTEC1 *Lox12* result is potentially interesting but remains fragile. The direction was aged-lower in per-sample summaries and had the same sign after stricter marker-positive filtering, and it persisted in a focused dropout/depth audit that included pairwise sample comparisons and depth-matched downsampling. An exact permutation check showed that the sample-level direction was ordered, but p-value resolution was limited by the n=2 versus n=2 design. However, the original mTEC1 comparison had only two biological samples per age group, aged mTEC1 cells had lower sequencing depth, and *Lox12* detection was sparse, so dropout/depth artifact remains a partial unresolved contributor. GSE223049 sorted bulk RNA-seq showed a broad epithelial aged-lower *Lox12* direction and E-MTAB-8560 showed aged-lower *Lox12* tendencies in broad TEC and mTEC/cTEC contexts, but the external datasets do not share the exact mTEC1 annotation. The safest interpretation is that *Lox12* shows a candidate age-associated transcript difference in annotated mTEC1 cells that has broad or approximate external directional context but still requires exact subtype-resolved testing.

Public protein and spatial resources provided weak indirect context only. Human thymus spatial RNA and epithelial RNA resources support LOXL2 detectability in thymic epithelial or medullary settings, and public antibody/protein resources can help prioritize wet-lab follow-up. However, these public resources did not provide direct mouse mTEC1 LOXL2 protein localization or age-aware thymic epithelial protein evidence, so they do not substitute for IHC or another direct protein assay.

The external comparisons separate broad directionality from subtype specificity. GSE223049 sorted bulk RNA-seq shows aged-lower mean broad thymic fibroblast *Lox* and *Lox12* and aged-lower mean broad thymic epithelial *Lox12*, but it cannot test capsFB, medFB, or mTEC1 specificity. It also does not test or replicate the medFB *Lox11* increase, because broad thymic fibroblast *Lox11* was weakly lower in aged GSE223049 samples. E-MTAB-8560 adds mTEC-focused transcript-level context for *Lox12*, with clearer aged-lower patterns in mTEC/cTEC than in mTEChi, but it spans the first year of mouse life and does not map exactly to the mTEC1 label used in GSE240016. GSE231906 adds supportive technical feasibility for aged-human thymus LOX-family transcript summaries and mixed transcript-level biological context for LOXL2 in mTEC-like compartments. Thus, the external evidence is most useful for broad fibroblast, epithelial, mTEC-focused, and human-context framing and does not establish exact subtype assignments, human conservation, or validation of the mouse mTEC1 candidate signal.

Future work should test these observations in balanced, independent thymic aging cohorts and with subtype-resolved methods. Spatial transcriptomics, RNA in situ hybridization, immunostaining, or sorted-cell qPCR could assess whether the transcript patterns reproduce at specific stromal and epithelial interfaces. Protein abundance, secretion, enzymatic activity, collagen and elastin organization, extracellular-matrix crosslinking, and thymic functional assays would be required before drawing mechanistic conclusions. Perturbation experiments would be needed before asking whether *Lox12* or other LOX-family members have any functional role in thymic aging rather than merely marking transcript differences among stromal states.

Limitations

This study has several important limitations. First, it remains a computational reanalysis centered on one public single-cell RNA-seq dataset, GSE240016. Second, the independent comparison from GSE223049 comes from sorted bulk RNA-seq rather than subtype-resolved single-cell data, so it is consistent only with broad fibroblast and broad epithelial directionality. Third, E-MTAB-8560 provides mTEC-focused transcript-level context but no fibroblast compartment, no exact mTEC1 annotation, and an age range ending at 52 weeks. E-MTAB-8560 is not exact GSE240016 mTEC1 replication; mTEC/cTEC support is clearer than mTEChi, batch-aware modeling remained limited, and the overall classification is mixed/inconclusive. Fourth, GSE231906 is technically usable for aged-human thymus LOX-family transcript context, and donor/sample-aware aging summaries were feasible, but expression matrix semantics remain partial. GSE231906 mean-value summaries are descriptive only, no cell-level tests were run, and the LOXL2 mTEC-like context is mixed rather than definitive. It does not validate the mouse result and does not establish human conservation. Fifth, the mTEC1 *Lox12* comparison in the original single-cell dataset includes only two biological samples per age group, and medFB *Lox11* increase lacks subtype-resolved external reanalysis. For mTEC1 *Lox12*, dropout/depth artifact remains a partial unresolved contributor despite directional persistence after depth-matched and pairwise sample checks. Exact permutation showed ordered sample-level direction, but p-value resolution is limited by n=2 versus n=2. Official GSE240016 GEO/SRA metadata mapping was improved, but batch

confounding was not ruled out. In addition, public protein/spatial resources provide only weak indirect support and no direct mouse mTEC1 LOXL2 protein localization or age-aware thymic epithelial protein evidence. Sixth, transcript abundance does not establish protein abundance, protein localization, secretion, LOX enzymatic activity, extracellular matrix composition, collagen or elastin crosslinking, tissue mechanics, thymic rejuvenation, or functional consequences.

Additional technical limitations remain. Age is partly entangled with sample identity, sequencing depth, and sample-preparation context, and batch or preparation effects cannot be fully excluded. Detection/dropout and low-count issues are substantial for several LOX-family findings, especially mTEC1 Loxl2. Single-cell-level tests and correlations are descriptive because individual cells are not independent biological replicates. The analysis relies on published annotations; marker-positive subset checks reduce but do not eliminate annotation uncertainty, with partial marker support for capsFB and intFB, especially in aged samples. External datasets also use different age ranges, species, technologies, sorting strategies, annotation systems, and compartment definitions, so cross-dataset agreement should be read as directional context rather than evidence for identical cell states. Finally, the study compares limited age groups and does not resolve trajectory, sex specificity, strain specificity, reversibility, or causality.

Data and Code Availability

Raw and processed single-cell RNA-seq data analyzed in this study are publicly available at the NCBI Gene Expression Omnibus under accession number GSE240016 [5]. The specific processed input file used here is available from GEO as:

https://www.ncbi.nlm.nih.gov/geo/download/?acc=GSE240016&format=file&file=GSE240016_CD45neg_thymic_stroma_d0%2Bannotation.h5ad

The local SHA256 checksum of the analyzed input file is:

A5CF4BAEB408412947DE2AE1867B4AD42271088C34C9DD212479C8B19E119FB1

Reviewers can recompute this checksum with `sha256sum data/raw/GSE240016_CD45neg_thymic_stroma_d0+annotation.h5ad` on Linux/macOS or `Get-FileHash data\raw\GSE240016_CD45neg_thymic_stroma_d0+annotation.h5ad -Algorithm SHA256` on Windows PowerShell. Analysis code, generated tables, reports, and manuscript-related files are available at <https://github.com/G1F12/ThymusLOXScan>.

New sensitivity, annotation-sanity, and external-comparison analyses for this version include `scripts/fb_composition_adjusted_lox.py`, `scripts/lox_detection_rates_by_sample.py`, `scripts/annotation_uncertainty_check.py`, `scripts/external_validation_gse223049_lox.py`, `scripts/external_validation_emtab8560_tec_lox.py`, `scripts/external_validation_gse231906_human_lox.py`, `scripts/build_cross_dataset_lox_matrix.py`, `scripts/figures/plot_cross_dataset_lox_validation.py`, `scripts/mtec1_loxl2_dropout_depth_audit.py`, and the `scripts/human_thymus_stage4_gse231906_*.py` controlled GSE231906 pilot scripts; reports in `reports/fb_composition_adjustment.md`, `reports/lox_detection_rate_analysis.md`, `reports/annotation_uncertainty_check.md`, `reports/external_dataset_search.md`, `reports/external_dataset_search_v2.md`, `reports/external_validation_plan.md`, `reports/gse223049_external_validation_v2.md`, `reports/emtab8560_tec_external_validation.md`, `reports/gse231906_human_external_validation.md`, `reports/cross_dataset_lox_analysis.md`, `reports/cross_dataset_figure_notes.md`, `reports/highest_impact_next_analysis.md`, `reports/hypothesis_falsification.md`, `reports/mtec1_loxl2_dropout_depth_audit.md`, `reports/loxl2_protein_spatial_validation_feasibility.md`, `reports/human_thymus_stage4_gse231906_balanced_report.md`, `reports/human_thymus_stage4_gse231906_decision_memo.md`, and `reports/human_thymus_stage4_gse231906_one_page_summary.md`; and tables in `results/tables/fb_composition_adjusted_lox.tsv`, `results/tables/lox_detection_rates_by_sample.tsv`, `results/tables/external_gse223049_lox_validation.tsv`, `results/tables/external_gse223049_lox_validation_summary_v2.tsv`, `results/tables/external_emtab8560_tec_lox_summary.tsv`, `results/tables/external_gse231906_human_lox_summary.tsv`, `results/tables/cross_dataset_lox_validation_matrix.tsv`, `results/tables/cross_dataset_lox_consistency_summary.tsv`, `results/figures/annotation_sanity/strict_marker_positive_lox_by_sample.tsv`, `results/figures/annotation_sanity/strict_marker_positive_lox_age_summary.tsv`, `results/tables/mtec1_loxl2_dropout_depth_audit.tsv`, `results/tables/mtec1_loxl2_pairwise_sample_direction.tsv`, `results/tables/mtec1_loxl2_downsampling_summary.tsv`, `results/tables/mtec1_loxl2_depth_adjusted_logistic.tsv`, `results/tables/loxl2_protein_spatial_evidence_audit.tsv`, `results/tables/loxl2_ihc_antibody_feasibility.tsv`, and `results/tables/human_thymus_stage4_gse231906_*`.

References

1. Palmer DB. The effect of age on thymic function. *Front Immunol.* 2013;4:316. doi:10.3389/fimmu.2013.00316.

2. Thomas R, Wang W, Su DM. Contributions of age-related thymic involution to immunosenescence and inflammaging. *Immun Ageing*. 2020;17:2. doi:10.1186/s12979-020-0173-8.
3. Ki S, Park D, Selden HJ, Seita J, Chung H, Kim J, Iyer VR, Ehrlich LIR. Global transcriptional profiling reveals distinct functions of thymic stromal subsets and age-related changes during thymic involution. *Cell Rep*. 2014;9(1):402-415. doi:10.1016/j.celrep.2014.08.070.
4. Hamazaki Y. Adult thymic epithelial cell (TEC) progenitors and TEC stem cells: models and mechanisms for TEC development and maintenance. *Eur J Immunol*. 2015;45(11):2985-2993. doi:10.1002/eji.201545844.
5. Kousa AI, Jahn L, Zhao K, Flores AE, Acenas D, Lederer E, et al. Age-related epithelial defects limit thymic function and regeneration. *Nat Immunol*. 2024;25(9):1593-1606. doi:10.1038/s41590-024-01915-9. PMID:39112630.
6. Gene Expression Omnibus. GSE240016: Age-related epithelial defects limit thymic function and regeneration [single cell]. NCBI GEO. Public on 2024-06-27. <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE240016>.
7. Lucero HA, Kagan HM. Lysyl oxidase: an oxidative enzyme and effector of cell function. *Cell Mol Life Sci*. 2006;63(19-20):2304-2316. doi:10.1007/s00018-006-6149-9.
8. Vallet SD, Guérault M, Belloy N, Dauchez M, Ricard-Blum S. Lysyl oxidases: from enzyme activity to extracellular matrix cross-links. *Essays Biochem*. 2019;63(3):349-364. doi:10.1042/EBC20180050.
9. Cox TR, Erler JT. Remodeling and homeostasis of the extracellular matrix: implications for fibrotic diseases and cancer. *Dis Model Mech*. 2011;4(2):165-178. doi:10.1242/dmm.004077.
10. Tschumperlin DJ, Ligresti G, Hilscher MB, Shah VH. Mechanosensing and fibrosis. *J Clin Invest*. 2018;128(1):74-84. doi:10.1172/JCI93561.
11. Liu X, Zhao Y, Gao J, Pawlyk B, Starcher B, Spencer JA, Yanagisawa H, Zuo J, Li T. Elastic fiber homeostasis requires lysyl oxidase-like 1 protein. *Nat Genet*. 2004;36(2):178-182. doi:10.1038/ng1297. PMID:14745449.
12. Añazco C, López-Jiménez AJ, Rafi M, Vega-Montoto L, Zhang MZ, Hudson BG, Vanacore RM. Lysyl oxidase-like-2 cross-links collagen IV of glomerular basement membrane. *J Biol Chem*. 2016;291(50):25999-26012. doi:10.1074/jbc.M116.738856. PMID:27770022.
13. Bignon M, Pichol-Thievend C, Hardouin J, Malbouyres M, Brechot N, Nasciutti L, et al. Lysyl oxidase-like protein-2 regulates sprouting angiogenesis and type IV collagen assembly in the endothelial basement membrane. *Blood*. 2011;118(14):3979-3989. doi:10.1182/blood-2010-10-313296. PMID:21835952.
14. Crowell HL, Soneson C, Germain PL, Calini D, Collin L, Raposo C, Malhotra D, Robinson MD. muscat detects subpopulation-specific state transitions from multi-sample multi-condition single-cell transcriptomics data. *Nat Commun*. 2020;11:6077. doi:10.1038/s41467-020-19894-4. PMID:33257685.
15. Squair JW, Gautier M, Kathe C, Anderson MA, James ND, Hutson TH, et al. Confronting false discoveries in single-cell differential expression. *Nat Commun*. 2021;12:5692. doi:10.1038/s41467-021-25960-2.
16. Murphy AE, Skene NG. A balanced measure shows superior performance of pseudobulk methods in single-cell RNA-sequencing analysis. *Nat Commun*. 2022;13:7851. doi:10.1038/s41467-022-35519-4.
17. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*. 2014;15:550. doi:10.1186/s13059-014-0550-8.
18. Muzellec B, Telenczuk M, Cabeli V, Andreux M. PyDESeq2: a python package for bulk RNA-seq differential expression analysis. *Bioinformatics*. 2023;39(9):btad547. doi:10.1093/bioinformatics/btad547.

Figure Legends

Figure 1. LOX-family pseudobulk differential expression in aging thymic stroma. Volcano plot of LOX-family pseudobulk DESeq2 results comparing aged 18-month samples with young 2-month samples. Each point represents one LOX-family gene tested in a broad cell type or fine subtype. The vertical dashed line marks $\log_2FC = 0$, and the horizontal dotted line marks $\text{padj} = 0.05$. Labeled points indicate LOX-family results meeting the table's FDR threshold in this small-sample pseudobulk analysis. Source figures: [results/figures/final/Fig1_pseudobulk_volcano_final.pdf](#) and [results/figures/final/Fig1_pseudobulk_volcano_final.png](#). Source table: [results/tables/lox_pseudobulk_complete_results.tsv](#).

Figure 2. Summary of LOX-family expression changes and descriptive correlations. (A) Pseudobulk $\log_2$ fold changes for LOX-family genes in selected annotated fibroblast and mTEC1 subtypes. Asterisks indicate $\text{padj} < 0.05$. (B) Pseudobulk LOX-family results meeting the table's FDR threshold, with color indicating $\log_2$ fold change aged versus young and bubble size reflecting adjusted significance. (C) Descriptive single-cell Spearman correlations between fibroblast *Loxl2* and *Col1a1*, *Vim*, or *Snai1*. (D) Descriptive single-cell rank-biserial effect sizes for pooled fibroblasts and mTEC1. Source figures: [results/figures/final/Fig2_summary_4panel_final.pdf](#) and [results/figures/final/Fig2_summary_4panel_final.png](#). Source tables: [results/tables/lox_pseudobulk_complete_results.tsv](#), [results/sc_spearman_correlations.csv](#), [results/sc_mannwhitney_FB_combined.csv](#), and [results/sc_mannwhitney_mTEC1.csv](#).

Figure 3. Cross-dataset LOX-family direction consistency. Cross-dataset summary of internal and external LOX-family direction checks. The heatmap summarizes whether each dataset/cell-group row is internally observed, directionally consistent, not testable, or

opposite in direction for each claim mapping. The external effect-size point plot shows descriptive high-age-minus-low-age deltas only where the available metrics are reasonably comparable across external summaries. The claim-level plot separates directionally consistent broad or approximate findings from internal-only findings. These panels do not label any result as subtype-resolved external evidence; GSE223049 is broad sorted bulk RNA-seq, E-MTAB-8560 provides TEC/mTEC-like rather than exact mTEC1 context, and GSE231906 is treated in v5.6 as mixed aged-human transcript-level context. Source figures: [results/figures/external_validation/cross_dataset/cross_dataset_lox_direction_heatmap.pdf](#), [results/figures/external_validation/cross_dataset/cross_dataset_lox_external_effect_points.pdf](#), and [results/figures/external_validation/cross_dataset/cross_dataset_lox_claim_consistency.pdf](#). Source tables: [results/tables/cross_dataset_lox_validation_matrix.tsv](#) and [results/tables/cross_dataset_lox_consistency_summary.tsv](#).

Supplementary Figure. Per-sample LOX-family pseudobulk expression checks. Per-biological-sample normalized pseudobulk expression values for key LOX-family findings, grouped by age. These plots are per-sample descriptive visualizations and do not replace pseudobulk DESeq2 inference. Source figures: [results/figures/per_sample/lox_per_sample_pseudobulk_expression.pdf](#) and [results/figures/per_sample/mtec1_lox12_per_sample.pdf](#).

Supplementary Figure. Annotation marker sanity checks. Marker score, marker detection, marker expression, and strict marker-positive LOX-family summary plots for capsFB, intFB, medFB, and mTEC1 annotations. These plots evaluate consistency with expected marker expression and do not prove annotation correctness. Source figures and tables are stored in [results/figures/annotation_sanity/](#), including [results/figures/annotation_sanity/strict_marker_positive_lox_summary.png](#), [results/figures/annotation_sanity/marker_summary_by_sample.tsv](#), and [results/figures/annotation_sanity/strict_marker_positive_lox_age_summary.tsv](#).