

1. Consider two sets of samples drawn from the same population that are randomly selected. Set X has a sample size = 10, and set Y has a sample size = 100. Which of the following statements is accurate about the confidence interval for the mean of the samples?

1 / 1 point

- ☒ The confidence interval for set X is **larger** than the confidence interval for set Y.
- ☐ The confidence interval for set X is **smaller** than the confidence interval for set Y.
- ☐ The confidence interval for set X **equals** the confidence interval for set Y.
- ☐ There isn't enough information to answer the question.

✓ **Correct**

Confidence interval estimates the range within a population parameter, in this case, the population mean. Since set X has a smaller sample size, the sample distribution gets further away from the normal distribution, and the standard deviation is larger. With a larger standard deviation, this results in a larger confidence interval.

2. Suppose you have a sample of 100 heights of individuals from a specific population. Assume the **population standard deviation** is 1 cm, and the **sample mean** is 175cm from a **random sample** of 100 individuals. What expression describes the margin of error for a confidence level of 99%?

1 / 1 point

- ☐ $z_{0.01} \cdot \frac{1}{10}$
- ☐ $z_{0.005} \cdot \frac{1}{100}$
- ☒ $z_{0.005} \cdot \frac{1}{10}$
- ☐ $z_{0.1} \cdot \frac{1}{100}$

✓ **Correct**

Since the question asks for a confidence level of 99% $\alpha/2 = \frac{0.01}{2} = 0.005$. The formula for the margin of error is $z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}$.

3. To calculate a confidence interval for the **mean** of a population, what assumptions must be made? **Select all that apply.**

1 / 1 point

- ☒ The sample is a random sample.

✓ **Correct**

A random sample is crucial for a reliable confidence interval for the population mean as it ensures representativeness. Random sampling provides an equal chance for each population member to be included, reducing bias and enhancing

- ☐ The population must follow a normal distribution.

- ☒ The sample size must be big enough (usually over 30).

✓ **Correct**

The Central Limit Theorem says the average sample mean **converges to** a normal distribution. It means that the bigger the sample, the closer it is to a normal, so we must ensure that there are enough points for this approximation to be good enough.

- ☐ The sample must have a mean = 0 and a standard deviation = 1.

4. You have a sample size of 20 from a population with unknown mean and standard deviation. You measured that the **sample mean** $\bar{X} = 50$ and the **sample standard deviation is** $s = 10$. A confidence interval of 95% of confidence level is given by:

1 / 1 point

Hint: $t_{0.975} = 2.093$

- ☐ (48.95, 51.05)
- ☒ (45.32, 54.68)
- ☐ (45.2, 54.8)
- ☐ (48.9, 51.1)

✔ Correct

5. A manufacturing company takes a sample of 100 items in its product warehouse and determines that 22% of the sample contains a defect. Calculate the **proportion margin of error** with a 95% confidence interval.

1 / 1 point

Hint: $z_{\alpha/2} = 1.96$

- ☐ 0.0336
- ☒ 0.0812
- ☐ 0.0919
- ☐ 0.3363

✔ Correct

When calculating the confidence interval for proportions, the formula is

$$\text{margin of error} = z_{\alpha/2} \cdot \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$$