

String amplitudes and automorphic representations (Lecture 1)

Daniel Persson

*Department of Mathematical Sciences
Chalmers University of Technology*



Bhaskara Math Seminars — Harish-Chandra Lecture Series
August 26, 2024


Outline

1. Motivation

2. Automorphic forms and representation theory

3. Small representations

4. Outlook



Suggested literature

Survey: “String scattering amplitudes and small automorphic representations”
w/ Guillaume Bossard and Axel Kleinschmidt, 2024
(Waseda number theory workshop - proceedings)

Book: “Eisenstein series and automorphic representations
- with applications in string theory”
w/ Philipp Fleig, Axel Kleinschmidt and Henrik Gustafsson
Cambridge University Press, 2018



Suggested literature


Survey: “String scattering amplitudes and small automorphic representations”
w/ Guillaume Bossard and Axel Kleinschmidt, 2024

Book: “Eisenstein series and automorphic representations - with applications in string theory”
w/ Philipp Fleig, Axel Kleinschmidt and Henrik Gustafsson
Cambridge University Press, 2018

Thanks to all collaborators and friends for discussions over the years:
Ahlén, Marcus Berg, Guillaume Bossard, Philipp Fleig, Solomon Friedberg, David Ginzburg, Dmitry Gourevitch, Michael Green, Henrik Gustafsson, Axel Kleinschmidt, Liu, Steve Miller, Manish Patnaik, Boris Pioline, Siddhartha Sahi, Pierre Vanhove

I. Motivation

Fourier coefficients of automorphic forms



- Fourier coefficients of **classical modular forms** encode deep number-theoretic information (counting points on elliptic curves etc..)
- **Moonshine:** relations with finite sporadic groups and CFT/string theory
- **Enumerative geometry:** rational curves on K3, GW-theory...
- Higher rank groups: **Langlands program** (automorphic L-functions, functoriality...)

See **Sol Friedberg's** talk in this series!
- The Fourier coefficients of Eisenstein series also encode **string theory scattering amplitudes**

QFT amplitudes


Compute the probability of a scattering process between elementary particles

$$\mathcal{A}(p_1, p_2, p_3, p_4) \in \mathbb{C}$$

QFT amplitudes

Compute the probability of a scattering process between elementary particles


$$\mathcal{A}(p_1, p_2, p_3, p_4) \in \mathbb{C}$$



QFT amplitudes

Compute the probability of a scattering process between elementary particles

$$\mathcal{A}(p_1, p_2, p_3, p_4) \in \mathbb{C}$$




Perturbation theory: expand for small coupling g

$$\mathcal{A}(p_1, p_2, p_3, p_4) = \sum_{n \geq 0} g^n \mathcal{A}_n(p_1, p_2, p_3, p_4) + \mathcal{A}^{\text{non-pert.}}(p_1, p_2, p_3, p_4)$$
$$\sim e^{-1/g}$$


String amplitudes

Understand the structure of **string interactions**



String amplitudes

Understand the structure of **string interactions**




String amplitudes

Understand the structure of **string interactions**



Strongly constrained by **symmetries!**

- supersymmetry
- U-duality



amplitudes have intricate
arithmetic structure $G(\mathbb{Z})$


String amplitudes

Understand the structure of **string interactions**



Strongly constrained by **symmetries!**


- supersymmetry
- U-duality



amplitudes have intricate
arithmetic structure $G(\mathbb{Z})$


Symmetry constrains interactions, leads to insights about:

- ultraviolet properties of gravity
- non-perturbative effects (black holes, instantons)
- novel mathematical predictions from physics



Toroidal compactifications yield the chain of **U-duality groups**

[Cremmer, Julia][Hull, Townsend]




D	G	K	$G(\mathbb{Z})$
10	$\text{SL}(2, \mathbb{R})$	$\text{SO}(2)$	$\text{SL}(2, \mathbb{Z})$
9	$\text{SL}(2, \mathbb{R}) \times \mathbb{R}^+$	$\text{SO}(2)$	$\text{SL}(2, \mathbb{Z})$
8	$\text{SL}(3, \mathbb{R}) \times \text{SL}(2, \mathbb{R})$	$\text{SO}(3) \times \text{SO}(2)$	$\text{SL}(3, \mathbb{Z}) \times \text{SL}(2, \mathbb{Z})$
7	$\text{SL}(5, \mathbb{R})$	$\text{SO}(5)$	$\text{SL}(5, \mathbb{Z})$
6	$\text{Spin}(5, 5, \mathbb{R})$	$(\text{Spin}(5) \times \text{Spin}(5))/\mathbb{Z}_2$	$\text{Spin}(5, 5, \mathbb{Z})$
5	$E_6(\mathbb{R})$	$\text{USp}(8)/\mathbb{Z}_2$	$E_6(\mathbb{Z})$
4	$E_7(\mathbb{R})$	$\text{SU}(8)/\mathbb{Z}_2$	$E_7(\mathbb{Z})$
3	$E_8(\mathbb{R})$	$\text{Spin}(16)/\mathbb{Z}_2$	$E_8(\mathbb{Z})$

Amplitudes are given by **functions** on

$$G(\mathbb{Z}) \backslash G(\mathbb{R}) / K$$


Toroidal compactifications yield the chain of **U-duality groups**

[Cremmer, Julia][Hull, Townsend]



D	G	K	$G(\mathbb{Z})$
10	$SL(2, \mathbb{R})$	$SO(2)$	$SL(2, \mathbb{Z})$
9	$SL(2, \mathbb{R}) \times \mathbb{R}^+$	$SO(2)$	$SL(2, \mathbb{Z})$
8	$SL(3, \mathbb{R}) \times SL(2, \mathbb{R})$	$SO(3) \times SO(2)$	$SL(3, \mathbb{Z}) \times SL(2, \mathbb{Z})$
7	$SL(5, \mathbb{R})$	$SO(5)$	$SL(5, \mathbb{Z})$
6	$Spin(5, 5, \mathbb{R})$	$(Spin(5) \times Spin(5))/\mathbb{Z}_2$	$Spin(5, 5, \mathbb{Z})$
5	$E_6(\mathbb{R})$	$USp(8)/\mathbb{Z}_2$	$E_6(\mathbb{Z})$
4	$E_7(\mathbb{R})$	$SU(8)/\mathbb{Z}_2$	$E_7(\mathbb{Z})$
3	$E_8(\mathbb{R})$	$Spin(16)/\mathbb{Z}_2$	$E_8(\mathbb{Z})$


Amplitudes



Action functional

Toroidal compactifications yield the chain of **U-duality groups**

[Cremmer, Julia][Hull, Townsend]



D	G	K	$G(\mathbb{Z})$
10	$SL(2, \mathbb{R})$	$SO(2)$	$SL(2, \mathbb{Z})$
9	$SL(2, \mathbb{R}) \times \mathbb{R}^+$	$SO(2)$	$SL(2, \mathbb{Z})$
8	$SL(3, \mathbb{R}) \times SL(2, \mathbb{R})$	$SO(3) \times SO(2)$	$SL(3, \mathbb{Z}) \times SL(2, \mathbb{Z})$
7	$SL(5, \mathbb{R})$	$SO(5)$	$SL(5, \mathbb{Z})$
6	$Spin(5, 5, \mathbb{R})$	$(Spin(5) \times Spin(5))/\mathbb{Z}_2$	$Spin(5, 5, \mathbb{Z})$
5	$E_6(\mathbb{R})$	$USp(8)/\mathbb{Z}_2$	$E_6(\mathbb{Z})$
4	$E_7(\mathbb{R})$	$SU(8)/\mathbb{Z}_2$	$E_7(\mathbb{Z})$
3	$E_8(\mathbb{R})$	$Spin(16)/\mathbb{Z}_2$	$E_8(\mathbb{Z})$

Amplitudes



Action functional

Einstein-Hilbert
action

$$\int d^4x \sqrt{G} R$$



Einstein's
equations

Higher-derivative action in type II string theory

Action functional:

$$\int d^{10-n}x \sqrt{G} \left[(\alpha')^3 f_0(g) \mathcal{R}^4 + (\alpha')^5 f_4(g) \partial^4 \mathcal{R}^4 + \dots \right]$$

Higher-derivative action in type II string theory

Action functional:


$$\int d^{10-n}x \sqrt{G} \left[(\alpha')^3 f_0(g) \mathcal{R}^4 + (\alpha')^5 f_4(g) \partial^4 \mathcal{R}^4 + \dots \right]$$

contraction of four Riemann tensors

Higher-derivative action in type II string theory


Action functional:

$$\int d^{10-n}x \sqrt{G} \left[(\alpha')^3 f_0(g) \mathcal{R}^4 + (\alpha')^5 f_4(g) \partial^4 \mathcal{R}^4 + \dots \right]$$



Higher-derivative action in type II string theory

Action functional:



Higher-derivative action in type II string theory

Action functional:

$$\int d^{10-n}x \sqrt{G} \left[(\alpha')^3 f_0(g) \mathcal{R}^4 + (\alpha')^5 f_4(g) \partial^4 \mathcal{R}^4 + \dots \right]$$

- $f_0(g), f_4(g)$ are functions of $g \in E_{n+1}(\mathbb{R})/K$
- must be **invariant** under U-duality $E_{n+1}(\mathbb{Z})$
- supersymmetry requires that they are
Laplacian eigenfunctions
- well-defined **weak-coupling expansions** as $g_s \rightarrow 0$

Higher-derivative action in type II string theory

Action functional:

$$\int d^{10-n}x \sqrt{G} \left[(\alpha')^3 f_0(g) \mathcal{R}^4 + (\alpha')^5 f_4(g) \partial^4 \mathcal{R}^4 + \dots \right]$$

- $f_0(g), f_4(g)$ are functions of $g \in E_{n+1}(\mathbb{R})/K$
- must be **invariant** under U-duality $E_{n+1}(\mathbb{Z})$
- supersymmetry requires that they are
Laplacian eigenfunctions
- well-defined **weak-coupling expansions** as $g_s \rightarrow 0$

defining properties
of an
**automorphic
form!**

Example: type IIB in D=10

$$\int d^{10}x \sqrt{G} f_0(\tau) \mathcal{R}^4$$

Example: type IIB in D=10

$$\int d^{10}x \sqrt{G} f_0(\tau) \mathcal{R}^4$$

→ $f_0(\tau)$ function of $\tau = x + iy \in \mathbb{H} \cong SL(2, \mathbb{R})/SO(2)$

“axio-dilaton”

Example: type IIB in D=10

$$\int d^{10}x \sqrt{G} f_0(\tau) \mathcal{R}^4$$

→ $f_0(\tau)$ function of $\tau = x + iy \in \mathbb{H} \cong SL(2, \mathbb{R})/SO(2)$ “axio-dilaton”

→ $f_0\left(\frac{a\tau + b}{c\tau + d}\right) = f_0(\tau)$ $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$ S-duality

Example: type IIB in D=10

$$\int d^{10}x \sqrt{G} f_0(\tau) \mathcal{R}^4$$

- $f_0(\tau)$ function of $\tau = x + iy \in \mathbb{H} \cong SL(2, \mathbb{R})/SO(2)$ “axio-dilaton”
- $f_0\left(\frac{a\tau + b}{c\tau + d}\right) = f_0(\tau)$ $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$ S-duality
- $\Delta_{\mathbb{H}} f_0(\tau) = -\frac{3}{4} f_0(\tau)$ $\Delta_{\mathbb{H}} = y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right)$ Supersymmetry

Example: type IIB in D=10

$$\int d^{10}x \sqrt{G} f_0(\tau) \mathcal{R}^4$$

- $f_0(\tau)$ function of $\tau = x + iy \in \mathbb{H} \cong SL(2, \mathbb{R})/SO(2)$ “axio-dilaton”
- $f_0\left(\frac{a\tau + b}{c\tau + d}\right) = f_0(\tau)$ $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$ S-duality
- $\Delta_{\mathbb{H}} f_0(\tau) = -\frac{3}{4} f_0(\tau)$ $\Delta_{\mathbb{H}} = y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right)$ Supersymmetry
- $f_0(\tau) \sim 2\zeta(3)y^{3/2}$ as $g_s = e^\phi = y^{-1} \rightarrow 0$ weak string coupling limit

Example: type IIB in D=10

$$\int d^{10}x \sqrt{G} f_0(\tau) \mathcal{R}^4$$

- $f_0(\tau)$ function of $\tau = x + iy \in \mathbb{H} \cong SL(2, \mathbb{R})/SO(2)$ “axio-dilaton”
- $f_0\left(\frac{a\tau + b}{c\tau + d}\right) = f_0(\tau)$ $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$ S-duality
- $\Delta_{\mathbb{H}} f_0(\tau) = -\frac{3}{4} f_0(\tau)$ $\Delta_{\mathbb{H}} = y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right)$ Supersymmetry
- $f_0(\tau) \sim 2\zeta(3)y^{3/2}$ as $g_s = e^\phi = y^{-1} \rightarrow 0$ weak string coupling limit

unique solution!

$$f_0(\tau) = \sum_{(m,n) \neq (0,0)} \frac{y^{3/2}}{|m + n\tau|^3}$$

[Green, Gutperle]
 [Green, Sethi]
 [Pioline]

Non-holomorphic Eisenstein series

Consider the sum:

$$E_s(\tau) = \sum_{(c,d)=1} \frac{y^s}{|c\tau + d|^{2s}}$$

non-holomorphic
Eisenstein series
 $s \in \mathbb{C}$

→ a function on

$$\mathbb{H} = \{\tau = x + iy \in \mathbb{C} \mid y > 0\}$$

→ invariant under

$$\tau \mapsto \gamma \cdot \tau = \frac{a\tau + b}{c\tau + d}$$

$$\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$$

Non-holomorphic Eisenstein series

Consider the sum:

$$E_s(\tau) = \sum_{(c,d)=1} \frac{y^s}{|c\tau + d|^{2s}}$$

non-holomorphic
Eisenstein series
 $s \in \mathbb{C}$

- a function on $\mathbb{H} = \{\tau = x + iy \in \mathbb{C} \mid y > 0\}$
- invariant under $\tau \mapsto \gamma \cdot \tau = \frac{a\tau + b}{c\tau + d}$
- converges absolutely for $\Re s > 1$
- $\Delta_{\mathbb{H}} E_s = s(s-1)E_s$

Invariance under $\tau \mapsto \tau + 1$ yields the **Fourier expansion**

$$E_s(\tau) = \underbrace{C(y; s)}_{\substack{\text{constant term} \\ \text{zero mode}}} + \underbrace{\sum_{n \neq 0} F_n(y; s) e^{2\pi i n x}}_{\text{non-zero mode}}$$

Invariance under $\tau \mapsto \tau + 1$ yields the **Fourier expansion**

$$E_s(\tau) = \underbrace{C(y; s)}_{\substack{\text{constant term} \\ \text{zero mode}}} + \underbrace{\sum_{n \neq 0} F_n(y; s) e^{2\pi i n x}}_{\text{non-zero mode}}$$

→ $C(y; s) = y^s + \frac{\xi(2s - 1)}{\xi(2s)} y^{1-s}$ completed zeta-function:
 $\xi(s) = \pi^{-s/2} \Gamma(s/2) \zeta(s)$

Invariance under $\tau \mapsto \tau + 1$ yields the **Fourier expansion**

$$E_s(\tau) = \underbrace{C(y; s)}_{\substack{\text{constant term} \\ \text{zero mode}}} + \underbrace{\sum_{n \neq 0} F_n(y; s) e^{2\pi i n x}}_{\text{non-zero mode}}$$

$\rightarrow C(y; s) = y^s + \frac{\xi(2s - 1)}{\xi(2s)} y^{1-s}$ completed zeta-function:
 $\xi(s) = \pi^{-s/2} \Gamma(s/2) \zeta(s)$

$\rightarrow F_n(y; s) = \frac{2\sqrt{y}}{\xi(2s)} |n|^{s-1/2} \mu_{1-2s}(n) K_{s-1/2}(2\pi |n| y)$

Invariance under $\tau \mapsto \tau + 1$ yields the **Fourier expansion**

$$E_s(\tau) = \underbrace{C(y; s)}_{\substack{\text{constant term} \\ \text{zero mode}}} + \underbrace{\sum_{n \neq 0} F_n(y; s) e^{2\pi i n x}}_{\text{non-zero mode}}$$

$\rightarrow C(y; s) = y^s + \frac{\xi(2s - 1)}{\xi(2s)} y^{1-s}$ completed zeta-function:
 $\xi(s) = \pi^{-s/2} \Gamma(s/2) \zeta(s)$

$\rightarrow F_n(y; s) = \frac{2\sqrt{y}}{\xi(2s)} |n|^{s-1/2} \mu_{1-2s}(n) K_{s-1/2}(2\pi |n| y)$

divisor sum: $\mu_s(n) = \sum_{d|n} d^s$

modified Bessel function

Example: $G = SL(2, \mathbb{R})$

$$\int d^{10}x \sqrt{G} f_0(\tau) \mathcal{R}^4$$

$$f_0(\tau) = \underbrace{2\zeta(3)y^{3/2} + 4\zeta(2)y^{-1/2}}_{\text{perturbative terms}} + \underbrace{2\pi \sum_{m \neq 0} \sqrt{|m|} \sigma_{-2}(m) e^{-S_{\text{inst}}(z)} [1 + \mathcal{O}(y^{-1})]}_{\text{non-perturbative terms}}$$

tree-level one-loop



amplitudes in the presence of instantons


Example: $G = SL(2, \mathbb{R})$

$$\int d^{10}x \sqrt{G} f_0(\tau) \mathcal{R}^4$$

$$f_0(\tau) = \underbrace{2\zeta(3)y^{3/2}}_{\text{tree-level}} + \underbrace{4\zeta(2)y^{-1/2}}_{\text{one-loop}} + 2\pi \sum_{m \neq 0} \sqrt{|m|} \sigma_{-2}(m) e^{-S_{\text{inst}}(z)} [1 + \mathcal{O}(y^{-1})]$$



amplitudes in the presence of instantons



Example: $G = SL(2, \mathbb{R})$

$$\int d^{10}x \sqrt{G} f_0(\tau) \mathcal{R}^4$$

perturbative terms

$$f_0(\tau) = \underbrace{2\zeta(3)y^{3/2}}_{\text{tree-level}} + \underbrace{4\zeta(2)y^{-1/2}}_{\text{one-loop}} + 2\pi \sum_{m \neq 0} \sqrt{|m|} \sigma_{-2}(m) e^{-S_{\text{inst}}(z)} [1 + \mathcal{O}(y^{-1})]$$

non-perturbative terms

amplitudes in the presence of instantons

instanton action

$$S_{\text{inst}}(z) := 2\pi |m| y - 2\pi i m x$$

Example: $G = SL(2, \mathbb{R})$

$$\int d^{10}x \sqrt{G} f_0(\tau) \mathcal{R}^4$$

$$f_0(\tau) = \underbrace{2\zeta(3)y^{3/2} + 4\zeta(2)y^{-1/2}}_{\text{perturbative terms}} + \underbrace{2\pi \sum_{m \neq 0} \sqrt{|m|} \sigma_{-2}(m) e^{-S_{\text{inst}}(z)} [1 + \mathcal{O}(y^{-1})]}_{\text{non-perturbative terms}}$$

tree-level one-loop



instanton action

$$S_{\text{inst}}(z) := 2\pi |m| y - 2\pi i m x$$

amplitudes in the presence of instantons

instanton measure

$$\sigma_{-2}(m) = \sum_{d|m} d^{-2}$$

Example: $G = SL(2, \mathbb{R})$

$$\int d^{10}x \sqrt{G} f_0(\tau) \mathcal{R}^4$$

$$f_0(\tau) = \underbrace{2\zeta(3)y^{3/2} + 4\zeta(2)y^{-1/2}}_{\text{perturbative terms}} + \underbrace{2\pi \sum_{m \neq 0} \sqrt{|m|} \sigma_{-2}(m) e^{-S_{\text{inst}}(z)} [1 + \mathcal{O}(y^{-1})]}_{\text{non-perturbative terms}}$$

tree-level one-loop

The diagram shows two Feynman-like diagrams. The left diagram, labeled 'tree-level', consists of two vertical lines connected by a horizontal line. The right diagram, labeled 'one-loop', consists of two vertical lines connected by a horizontal line that contains a small loop.

amplitudes in the presence of instantons

Fourier coefficients of automorphic forms encode information about scattering amplitudes!

2. Automorphic forms and representation theory

Data:

- ▶ $G(\mathbb{R})$ real simple Lie group (e.g. $SL(n, \mathbb{R})$)
- ▶ $G(\mathbb{Z}) \subset G$ arithmetic subgroup (e.g. $SL(n, \mathbb{Z})$)

Definition:

An **automorphic form** is a smooth function $\varphi : G \longrightarrow \mathbb{C}$ satisfying

1. Automorphy: $\forall \gamma \in G(\mathbb{Z}), \varphi(\gamma g) = \varphi(g)$
2. φ is an eigenfunction of the ring of inv. diff. operators on G
3. φ has well-behaved growth conditions

Sometimes we also add a condition involving the maximal compact subgroup
(K -finiteness)

Eisenstein series on semi-simple Lie groups

The **Langlands Eisenstein series** on a semi-simple Lie group is defined by:

$$E(\lambda, g) = \sum_{\gamma \in B(\mathbb{Z}) \backslash G(\mathbb{Z})} e^{\langle \lambda + \rho | H(\gamma g) \rangle}$$

Eisenstein series on semi-simple Lie groups

The **Langlands Eisenstein series** on a semi-simple Lie group is defined by:

$$E(\lambda, g) = \sum_{\gamma \in B(\mathbb{Z}) \backslash G(\mathbb{Z})} e^{\langle \lambda + \rho | H(\gamma g) \rangle}$$

Iwasawa decomposition: $G = BK = NAK$

$$A \sim \begin{pmatrix} * & & \\ & * & \\ & & * \end{pmatrix}$$

$$N \sim \begin{pmatrix} 1 & * & * \\ & 1 & * \\ & & 1 \end{pmatrix}$$

Eisenstein series on semi-simple Lie groups

The **Langlands Eisenstein series** on a semi-simple Lie group is defined by:

$$E(\lambda, g) = \sum_{\gamma \in B(\mathbb{Z}) \backslash G(\mathbb{Z})} e^{\langle \lambda + \rho | H(\gamma g) \rangle}$$

Iwasawa decomposition: $G = BK = NAK$

Logarithm map: $H : G \rightarrow \mathfrak{h} = \text{Lie } A \quad H(nak) = \log a$

Eisenstein series on semi-simple Lie groups

The **Langlands Eisenstein series** on a semi-simple Lie group is defined by:

$$E(\lambda, g) = \sum_{\gamma \in B(\mathbb{Z}) \backslash G(\mathbb{Z})} e^{\langle \lambda + \rho | H(\gamma g) \rangle}$$

Iwasawa decomposition: $G = BK = NAK$

Logarithm map: $H : G \rightarrow \mathfrak{h} = \text{Lie } A \quad H(nak) = \log a$

Weight: $\lambda \in \mathfrak{h}^* \otimes \mathbb{C}$

Weyl vector: $\rho = \frac{1}{2} \sum_{\alpha > 0} \alpha$

Eisenstein series on semi-simple Lie groups

The **Langlands Eisenstein series** on a semi-simple Lie group is defined by:

$$E(\lambda, g) = \sum_{\gamma \in B(\mathbb{Z}) \backslash G(\mathbb{Z})} e^{\langle \lambda + \rho | H(\gamma g) \rangle}$$

- Converges absolutely on a subspace of $\mathfrak{h}^* \otimes \mathbb{C}$ Godement's domain
 $\{\lambda \mid \langle \lambda, \alpha \rangle > 1, \forall \alpha \in \Pi\}$
- Can be continued to a meromorphic function on all of $\mathfrak{h}^* \otimes \mathbb{C}$ [Langlands]

Eisenstein series on semi-simple Lie groups

The **Langlands Eisenstein series** on a semi-simple Lie group is defined by:

$$E(\lambda, g) = \sum_{\gamma \in B(\mathbb{Z}) \backslash G(\mathbb{Z})} e^{\langle \lambda + \rho | H(\gamma g) \rangle}$$

- Converges absolutely on a subspace of $\mathfrak{h}^* \otimes \mathbb{C}$ Godement's domain
 $\{\lambda \mid \langle \lambda, \alpha \rangle > 1, \forall \alpha \in \Pi\}$
- Can be continued to a meromorphic function on all of $\mathfrak{h}^* \otimes \mathbb{C}$ [Langlands]
- Invariant: $E(\lambda, \gamma g k) = E(\lambda, g)$ $\gamma \in G(\mathbb{Z})$ $k \in K$
- Eigenfunction of the Laplacian: $\Delta_{G/K} E(\lambda, g) = \frac{1}{2}(\langle \lambda | \lambda \rangle - \langle \rho | \rho \rangle) E(\lambda, g)$
- Functional relation: $E(\lambda, g) = M(w, \lambda) E(w\lambda, g), \quad \forall w \in W(\mathfrak{g})$

Eisenstein series on semi-simple Lie groups

The **Langlands Eisenstein series** on a semi-simple Lie group is defined by:

$$E(\lambda, g) = \sum_{\gamma \in B(\mathbb{Z}) \backslash G(\mathbb{Z})} e^{\langle \lambda + \rho | H(\gamma g) \rangle}$$

Example: $G = SL(2, \mathbb{R})$

$$E(s, g) = \sum_{(c,d)=1} \frac{y^s}{|c\tau + d|^{2s}} = \sum_{\gamma \in B(\mathbb{Z}) \backslash SL(2, \mathbb{Z})} e^{\langle \lambda + \rho | H(\gamma g) \rangle}$$

$$\lambda + \rho = 2s\Lambda \quad (\text{fundamental weight: } \Lambda = \alpha/2)$$

$$H(a) = H(e^{yH_\alpha}) = yH_\alpha \quad \langle \Lambda | H_\alpha \rangle = 1$$

Automorphic representations

Eisenstein series are attached to the (non-unitary) **principal series**:

$$I(\lambda) = \text{Ind}_B^G \chi = \{f : G \rightarrow \mathbb{C} \mid f(bg) = \chi(b)f(g), b \in B\}$$

Automorphic representations

Eisenstein series are attached to the (non-unitary) **principal series**:

$$I(\lambda) = \text{Ind}_B^G \chi = \{f : G \rightarrow \mathbb{C} \mid f(bg) = \chi(b)f(g), b \in B\}$$

The theory of Eisenstein series then defines a map

$$E : I(\lambda) \rightarrow \mathcal{A}(G(\mathbb{Z}) \backslash G(\mathbb{R}))$$

from the principal series to **the space of automorphic forms on $G(\mathbb{R})$**

Automorphic representations

Eisenstein series are attached to the (non-unitary) **principal series**:

$$I(\lambda) = \text{Ind}_B^G \chi = \{f : G \rightarrow \mathbb{C} \mid f(bg) = \chi(b)f(g), b \in B\}$$

The theory of Eisenstein series then defines a map

$$E : I(\lambda) \rightarrow \mathcal{A}(G(\mathbb{Z}) \backslash G(\mathbb{R}))$$

from the principal series to **the space of automorphic forms on $G(\mathbb{R})$**

G acts on $\mathcal{A}(G(\mathbb{Z}) \backslash G(\mathbb{R}))$ by **right-translation**:

$$[\rho(h)f](g) = f(gh)$$

The irreducible constituents in the decomposition of $\mathcal{A}(G(\mathbb{Z}) \backslash G(\mathbb{R}))$ under this action are called **automorphic representations**

Toy model: Fourier analysis on $\mathbb{Z} \setminus \mathbb{R} \cong S^1$

Any function $f \in C^\infty(\mathbb{Z} \setminus \mathbb{R})$ can be decomposed into a Fourier series:

$$f(x) = \sum_{k \in \mathbb{Z}} c_k \psi_k(x)$$

$$\psi_k : \mathbb{Z} \setminus \mathbb{R} \rightarrow U(1) \quad \psi_k(x) = e^{2\pi i k x} \quad k \in \mathbb{Z}, x \in \mathbb{R}$$

Toy model: Fourier analysis on $\mathbb{Z} \backslash \mathbb{R} \cong S^1$

Any function $f \in C^\infty(\mathbb{Z} \backslash \mathbb{R})$ can be decomposed into a Fourier series:

$$f(x) = \sum_{k \in \mathbb{Z}} c_k \psi_k(x)$$

$$\psi_k : \mathbb{Z} \backslash \mathbb{R} \rightarrow U(1) \quad \psi_k(x) = e^{2\pi i k x} \quad k \in \mathbb{Z}, x \in \mathbb{R}$$

Moderate growth: restrict to **square integrable functions**

$$L^2(\mathbb{Z} \backslash \mathbb{R}) = \{f \in C^\infty(\mathbb{Z} \backslash \mathbb{R}) \mid \sum_{k \in \mathbb{Z}} |c_k|^2 < \infty\}$$

$G = \mathbb{R}$ acts on $L^2(\mathbb{Z} \backslash \mathbb{R})$ via the regular representation

$$(\rho(y)f)(x) = f(x + y)$$

Toy model: Fourier analysis on $\mathbb{Z} \setminus \mathbb{R} \cong S^1$

Any function $f \in C^\infty(\mathbb{Z} \setminus \mathbb{R})$ can be decomposed into a Fourier series:

$$f(x) = \sum_{k \in \mathbb{Z}} c_k \psi_k(x)$$

$$\psi_k : \mathbb{Z} \setminus \mathbb{R} \rightarrow U(1) \quad \psi_k(x) = e^{2\pi i k x} \quad k \in \mathbb{Z}, x \in \mathbb{R}$$

Moderate growth: restrict to **square integrable functions**

$$L^2(\mathbb{Z} \setminus \mathbb{R}) = \{f \in C^\infty(\mathbb{Z} \setminus \mathbb{R}) \mid \sum_{k \in \mathbb{Z}} |c_k|^2 < \infty\}$$

$$L^2(\mathbb{Z} \setminus \mathbb{R}) = \bigoplus_{k \in \mathbb{Z}} \mathbb{C} \psi_k$$

Toy model: Fourier analysis on $\mathbb{Z} \backslash \mathbb{R} \cong S^1$

Any function $f \in C^\infty(\mathbb{Z} \backslash \mathbb{R})$ can be decomposed into a Fourier series:

$$f(x) = \sum_{k \in \mathbb{Z}} c_k \psi_k(x)$$

$$\psi_k : \mathbb{Z} \backslash \mathbb{R} \rightarrow U(1) \quad \psi_k(x) = e^{2\pi i k x} \quad k \in \mathbb{Z}, x \in \mathbb{R}$$

Moderate growth: restrict to **square integrable functions**

$$L^2(\mathbb{Z} \backslash \mathbb{R}) = \{f \in C^\infty(\mathbb{Z} \backslash \mathbb{R}) \mid \sum_{k \in \mathbb{Z}} |c_k|^2 < \infty\}$$

$$L^2(\mathbb{Z} \backslash \mathbb{R}) = \bigoplus_{k \in \mathbb{Z}} \mathbb{C} \psi_k$$

“automorphic representation”

Automorphic representations

There is an important notion of “size” of an automorphic representation, called the **Gelfand-Kirillov dimension**.

GKdim = “smallest number of variables on which the functions depend”

Automorphic representations

There is an important notion of “size” of an automorphic representation, called the **Gelfand-Kirillov dimension**.

$\text{GKdim} =$ “smallest number of variables on which the functions depend”

Example: $\text{GKdim}(L^2(\mathbb{R}^n)) = n$

Automorphic representations

There is an important notion of “size” of an automorphic representation, called the **Gelfand-Kirillov dimension**.

$\text{GKdim} =$ “smallest number of variables on which the functions depend”

Example: $\text{GKdim}(L^2(\mathbb{R}^n)) = n$

For the principal series we have:

$$\text{GKdim}(I(\lambda)) = \dim_{\mathbb{R}} B \backslash G = \dim_{\mathbb{R}} N$$

This is **important for physics**, since we have the rough correspondence:

Automorphic representations

There is an important notion of “size” of an automorphic representation, called the **Gelfand-Kirillov dimension**.

$\text{GKdim} =$ “smallest number of variables on which the functions depend”

Example: $\text{GKdim}(L^2(\mathbb{R}^n)) = n$

For the principal series we have:

$$\text{GKdim}(I(\lambda)) = \dim_{\mathbb{R}} B \backslash G = \dim_{\mathbb{R}} N$$

This is **important for physics**, since we have the correspondence:

number of independent **physical charges** (e.g. electric, magnetic)



Gelfand-Kirillov dimension of the associated automorphic representation

3. Fourier coefficients

Fourier coefficients

The periodicity $f(\tau + 1) = f(\tau)$ generalises to

$$E(\lambda, ng) = E(\lambda, g) \quad n \in N(\mathbb{Z})$$

Much more complicated since $N(\mathbb{Z})$ is **non-abelian**.

Fourier coefficients

The periodicity $f(\tau + 1) = f(\tau)$ generalises to

$$E(\lambda, ng) = E(\lambda, g) \quad n \in N(\mathbb{Z})$$

Much more complicated since $N(\mathbb{Z})$ is **non-abelian**.

General structure:

$$E(\lambda, g) = E^{\text{const}}(\lambda, g) + \sum_{\psi} W_{\psi}(\lambda, g) + \dots$$



constant term

(zero-mode)

**perturbative
effects**

Fourier coefficients

The periodicity $f(\tau + 1) = f(\tau)$ generalises to

$$E(\lambda, ng) = E(\lambda, g) \quad n \in N(\mathbb{Z})$$

Much more complicated since $N(\mathbb{Z})$ is **non-abelian**.

General structure:

$$E(\lambda, g) = E^{\text{const}}(\lambda, g) + \sum_{\psi} W_{\psi}(\lambda, g) + \dots$$



abelian coefficient

(non zero-modes)

**non-perturbative
effects**

Fourier coefficients

The periodicity $f(\tau + 1) = f(\tau)$ generalises to

$$E(\lambda, ng) = E(\lambda, g) \quad n \in N(\mathbb{Z})$$

Much more complicated since $N(\mathbb{Z})$ is **non-abelian**.

General structure:

$$E(\lambda, g) = E^{\text{const}}(\lambda, g) + \sum_{\psi} W_{\psi}(\lambda, g) + \dots$$



non-abelian coefficients
(non zero-modes)
non-perturbative
effects

Whittaker coefficients

$$E(\lambda, g) = E^{\text{const}}(\lambda, g) + \sum_{\psi} W_{\psi}(\lambda, g) + \dots$$

Whittaker coefficients

$$E(\lambda, g) = E^{\text{const}}(\lambda, g) + \sum_{\psi} W_{\psi}(\lambda, g) + \dots$$

$$W_{\psi}(g) = \int_{N(\mathbb{Z}) \backslash N(\mathbb{R})} E(\lambda, ng) \overline{\psi(n)} dn$$

**Whittaker
coefficient**

$$\psi : N(\mathbb{Z}) \backslash N(\mathbb{R}) \rightarrow U(1)$$

unitary character on $N(\mathbb{R})$
trivial on $N(\mathbb{Z})$

Whittaker coefficients

$$E(\lambda, g) = E^{\text{const}}(\lambda, g) + \sum_{\psi} W_{\psi}(\lambda, g) + \dots$$

$$W_{\psi}(g) = \int_{N(\mathbb{Z}) \backslash N(\mathbb{R})} E(\lambda, ng) \overline{\psi(n)} dn$$

Whittaker coefficient

$$\psi : N(\mathbb{Z}) \backslash N(\mathbb{R}) \rightarrow U(1) \quad \text{unitary character on } N(\mathbb{R})$$

→ $\psi(n) = e^{2\pi i \sum_j m_j x_j}$ (simple roots)

$$m_j \in \mathbb{Z}$$

if all $m_j \neq 0$ then ψ is **generic**

if some $m_j = 0$ then ψ is **degenerate**

$$x_j \in \mathbb{R}$$

$$\textbf{Holomorphic modular form} \quad f(\tau) \qquad \tau \in \mathbb{H} \cong SL(2, \mathbb{R})/U(1)$$

$$\psi\left(\begin{pmatrix} 1 & x \\ & 1 \end{pmatrix}\right) = \psi(e^{xE_\alpha}) = e^{2\pi imx}$$

$$x \in \mathbb{R} \qquad m \in \mathbb{Z}$$

$$\psi \text{ generic} \iff m \neq 0$$

$$\textbf{Holomorphic modular form} \quad f(\tau) \qquad \tau \in \mathbb{H} \cong SL(2, \mathbb{R})/U(1)$$

$$\psi \left(\begin{pmatrix} 1 & x \\ & 1 \end{pmatrix} \right) = \psi(e^{xE_\alpha}) = e^{2\pi i mx}$$

$$x \in \mathbb{R} \qquad m \in \mathbb{Z}$$

$$\psi \text{ generic} \iff m \neq 0$$

$$W_m(\tau) = \int_0^1 f(\tau + 1)e^{-2\pi i mu} du \\ = c(m)e^{2\pi i m\tau}$$

$$\textbf{Holomorphic modular form} \quad f(\tau) \qquad \tau \in \mathbb{H} \cong SL(2, \mathbb{R})/U(1)$$

$$\psi \left(\begin{pmatrix} 1 & x \\ & 1 \end{pmatrix} \right) = \psi(e^{xE_\alpha}) = e^{2\pi i mx}$$

$$x \in \mathbb{R} \qquad m \in \mathbb{Z}$$

$$\psi \text{ generic} \iff m \neq 0$$

$$W_m(\tau) = \int_0^1 f(\tau + 1)e^{-2\pi i mu} du \\ = c(m)e^{2\pi im\tau}$$

Non-holomorphic Eisenstein series

$$E(s, \tau) = \sum_{\substack{(m, n) \in \mathbb{Z}^2 \\ (m, n) = 1}} \frac{y^s}{|m + n\tau|^{2s}} \qquad s \in \mathbb{C} \qquad \tau = x + iy \in \mathbb{H}$$

$$\textbf{Holomorphic modular form} \quad f(\tau) \quad \tau \in \mathbb{H} \cong SL(2, \mathbb{R})/U(1)$$

$$\psi \left(\begin{pmatrix} 1 & x \\ & 1 \end{pmatrix} \right) = \psi(e^{xE_\alpha}) = e^{2\pi i mx}$$

$$x \in \mathbb{R} \qquad m \in \mathbb{Z}$$

$$\psi \text{ generic} \iff m \neq 0$$

$$W_m(\tau) = \int_0^1 f(\tau + 1)e^{-2\pi imu} du$$

$$= c(m)e^{2\pi im\tau}$$

Non-holomorphic Eisenstein series

$$E(s, \tau) = \sum_{\substack{(m, n) \in \mathbb{Z}^2 \\ (m, n) = 1}} \frac{y^s}{|m + n\tau|^{2s}}$$

$$s \in \mathbb{C}$$

$$\tau = x + iy \in \mathbb{H}$$

$$W_m(\tau) = \int_0^1 E(s, \tau + u)e^{-2\pi imu} du = \frac{\sqrt{y}}{\xi(2s)} \sigma_{1-2s}(m) K_{s-1/2}(2\pi|m|y) e^{2\pi imx}$$

$$\sigma_{1-2s}(m) = \sum_{d|m} d^{1-2s}$$



 (modified) Bessel function

Euler products

Before we proceed with the higher rank case we mention some further properties of the Fourier expansion, namely that it [decomposes into Euler products](#)

$$W_\psi(g) = W_\infty(g) \prod_{p < \infty} W_p(1)$$

Euler products

Before we proceed with the higher rank case we mention some further properties of the Fourier expansion, namely that it **decomposes into Euler products**

$$W_\psi(g) = W_\infty(g) \prod_{p < \infty} W_p(1)$$

archimedean Whittaker coefficient


$$W_\infty(g) = \frac{2\pi^s}{\Gamma(s)} y|m|^{s-1/2} K_{s-1/2}(2\pi|m|y) e^{2\pi i mx}$$

$$g = nak = \begin{pmatrix} 1 & x \\ & 1 \end{pmatrix} \begin{pmatrix} y^{1/2} & \\ & y^{-1/2} \end{pmatrix} k \in SL(2, \mathbb{R})$$

Euler products

Before we proceed with the higher rank case we mention some further properties of the Fourier expansion, namely that it **decomposes into Euler products**

$$W_\psi(g) = W_\infty(g) \prod_{p < \infty} W_p(1)$$



p -adic Whittaker coefficient

$$W_p(1) = (1 - p^{-2s}) \frac{1 - p^{-2s+1} |m|_p^{2s-1}}{1 - p^{-2s+1}}$$

Euler products

Before we proceed with the higher rank case we mention some further properties of the Fourier expansion, namely that it **decomposes into Euler products**

$$W_\psi(g) = W_\infty(g) \prod_{p < \infty} W_p(1)$$

\nearrow

$1 \in SL(2, \mathbb{Q}_p)$

p -adic Whittaker coefficient

$$W_p(1) = (1 - p^{-2s}) \frac{1 - p^{-2s+1} |m|_p^{2s-1}}{1 - p^{-2s+1}}$$

\searrow

This is the basis for the **adelic** formulation of automorphic forms

Adelic framework


An efficient, but abstract, way to approach the subject of automorphic forms is by the introduction of adeles, rather ungainly objects that nevertheless, once familiar, spare much unnecessary thought and many useless calculations.

— Robert P. Langlands

Adelic framework

An efficient, but abstract, way to approach the subject of automorphic forms is by the introduction of adeles, rather ungainly objects that nevertheless, once familiar, spare much unnecessary thought and many useless calculations.

— Robert P. Langlands



Adelic framework

arithmetic groups $G(\mathbb{Z}) \subset G(\mathbb{R}) \longrightarrow G(\mathbb{Q}) \subset G(\mathbb{A})$

space of
automorphic forms $\mathcal{A}(G(\mathbb{Z}) \backslash G(\mathbb{R})) \longrightarrow \mathcal{A}(G(\mathbb{Q}) \backslash G(\mathbb{A}))$

Adelic framework

$$\text{arithmetic groups} \quad G(\mathbb{Z}) \subset G(\mathbb{R}) \longrightarrow G(\mathbb{Q}) \subset G(\mathbb{A})$$

$$\text{space of automorphic forms} \quad \mathcal{A}(G(\mathbb{Z}) \backslash G(\mathbb{R})) \longrightarrow \mathcal{A}(G(\mathbb{Q}) \backslash G(\mathbb{A}))$$

$$\text{Eisenstein series} \quad \sum_{\gamma \in B(\mathbb{Z}) \backslash G(\mathbb{Z})} e^{\langle \lambda + \rho | H(\gamma g) \rangle} \longrightarrow \sum_{\gamma \in B(\mathbb{Q}) \backslash G(\mathbb{Q})} e^{\langle \lambda + \rho | H(\gamma g) \rangle}$$

$$\lambda \in \mathfrak{h}^* \otimes \mathbb{C} \qquad \qquad H : G \rightarrow \mathfrak{h}$$

Theorem [Jacquet, Langlands]: The generic Whittaker coefficient is Eulerian

$$W_\psi(\lambda, g) = \int_{N(\mathbb{A})} \chi(w_0 n g) \overline{\psi(n)} dn = \prod_p W_{\psi_p}$$

Theorem [Jacquet, Langlands]: The generic Whittaker coefficient is Eulerian

$$W_\psi(\lambda, g) = \int_{N(\mathbb{A})} \chi(w_0 n g) \overline{\psi(n)} dn = \prod_p W_{\psi_p}$$


$w_0 = \text{longest element of } W(\mathfrak{g})$

Theorem [Jacquet, Langlands]: The generic Whittaker coefficient is Eulerian

$$W_\psi(\lambda, g) = \int_{N(\mathbb{A})} \chi(w_0 n g) \overline{\psi(n)} dn = \prod_p W_{\psi_p}$$

$$W_{\psi_\infty} = \int_{N(\mathbb{R})} \chi_\infty(w_0 n a_\infty) \overline{\psi_\infty(n)} dn$$

$$W_{\psi_p} = \int_{N(\mathbb{Q}_p)} \chi_p(w_0 n a_p) \overline{\psi_p(n)} dn$$

Theorem [Jacquet, Langlands]: The generic Whittaker coefficient is Eulerian

$$W_\psi(\lambda, g) = \int_{N(\mathbb{A})} \chi(w_0 n g) \overline{\psi(n)} dn = \prod_p W_{\psi_p}$$

Theorem [Shintani, Casselman-Shalika]: When $p < \infty$ we have

W_{ψ_p} = Weyl character formula for ${}^L G$

Perturbative limit - choices of unipotent subgroups

$$P = LU$$

Levi subgroups:

→ **Decompactification limit**

- perturbative parameter: radius of decompactified circle
- non-perturbative effects: KK-instantons, BPS-instantons



$$L = E_7$$

→ **String perturbation limit**

- perturbative parameter: string coupling
- non-perturbative effects: D-instantons, NS5-instantons



$$L = D_7$$

→ **M-theory limit**

- perturbative parameter: volume of M-theory torus
- non-perturbative effects: M2- & M5-instantons



$$L = A_7$$

Example $SL(3, \mathbb{R})$

$$N = \left\{ \begin{pmatrix} 1 & * & * \\ & 1 & * \\ & & 1 \end{pmatrix} \right\}$$

$$U_{\alpha_1} = \left\{ \begin{pmatrix} 1 & * & * \\ & 1 & \\ & & 1 \end{pmatrix} \right\}, \quad U_{\alpha_2} = \left\{ \begin{pmatrix} 1 & & * \\ & 1 & * \\ & & 1 \end{pmatrix} \right\}$$

**In tomorrow's lecture we will
talk about:**

Automorphic forms on Kac-Moody groups

Small representations of exceptional groups

Open questions/future research

Thank you!