Assignment 2

02610 Optimization and Data Fitting – Anders Hørsted (s082382)

Question 1: Fitting to Air Pollution Data

In this question we are going to fit sums of sines and cosines to the Air Pollution Data given in the assignment. First a function NOfit that calculates the parameters and the residuals for the fit is created.

Question 1.1

The function should have the call [x_star, r_star] = NOfit(t,y,n), but for convenience it is written to also return the design matrix A. The function NOfit relies on get_A to calculate the actual design matrix. The function get_A is used later when the fits are plotted.

```
function [xstar, rstar, A] = NOfit(t, y, n)

    omega = 2*pi/24;
    m = length(t);

if m ~= length(y)
        error('The length of t and y should match')
    end

A = get_A(t, n);

xstar = (A'*A)\(A'*y);
    rstar = y - A*xstar;
end
```

Code Listing 1: NOfit function to fit sine, cosines of order n

```
function [A] = get_A(t, n)

if mod(n, 2) == 0
    error('Only odd number of basis functions');
end

omega = 2*pi/24;
m = length(t);
A = zeros(m, n);
fns = {@sin, @cos};
```

```
for i=1:m
    for j=1:n
        % Cycle between sine and cosine
        f = fns{mod(j,2)+1};
            A(i,j) = f(floor(j/2)*omega*t(i));
    end
end
end
```

Code Listing 2: Helper function get_A used to generate design matrix

Question 1.2

See appendix A.1 for code used in this exercise.

The NOfit function is now tested. Using the data given in the assignment text a 3rd order cosine is fitted which gives the model.

$$M(\mathbf{x}, t) = 186.81 - 44.94\sin(\omega t) - 93.43\cos(\omega t)$$

To confirm the implementation of N0fit the residual 2-norm is checked and it is the expected $||\mathbf{r}^*||_2 = 292.558$. Since the implementation seems to be right, the fitted model is plotted along with the data in figure 1.




Figure 1: CAPTION!!!

Question 1.3

See appendix A.2 for code used in this exercise.

The optimal order of the fit must be determined. Using the test for random signs and the test for correlation, for the orders n = 3, 5, 7, 9, 11, 13 gives the results shown in figure 2.




Figure 2: CAPTION!!!

A Appendices

All MATLAB source code is included in the appendices. All the source code including the LaTex code used for the report can also be found at https://github.com/alphabits/dtu-fall-2011/tree/master/02610/assignment-2.

A.1 Question 1.2

```
load_ex1;
n = 3;
[xstar, rstar] = NOfit(t, y, n);
file = fopen('../tables/3rd-order-fitted-model.tex', 'w');
fprintf(file, 'M(\\myvec{x}, t) = %.02f %+.02f \\sin(\\omega t) %+.02f \\cos(\\omega t)', xstar);
fclose(file);
plot_fit(t, y, n, xstar, rstar, '3rd order fit for Air Pollution Data');
saveeps('../media/3rd-order-fit.eps');
```

Code Listing 3: ex12.m

A.2 Question 1.3

```
load_ex1;

for n=3:2:13
    [x, r] = N0fit(t, y, n);
    plot_fit_with_res_analysis(t, y, n, x, r);
    saveeps(sprintf('../media/order-determination-%d.eps', n));
end
```

Code Listing 4: ex13.m

A.3 Helper functions

```
function [] = plot_fit(t, y, n, x, r, plot_title)
    tplot = linspace(1,24,100);
    A = get_A(tplot, n);
    fit = A*x;
    fs = 18;
    set(gca, 'fontsize', fs);
    plot(tplot, fit, '-b', t, y, 'or');
    title(plot_title, 'fontsize', fs);
```

end

Code Listing 5: plot_fit.m

Code Listing 6: plot_fit_with_res_analysis.m

References

- [1] Jorge Nocedal & Stephen J. Wright, *Numerical Optimization*. Springer Science+Business Media, 2nd Edition, 2006.
- [2] Kaj Madsen & Hans Bruun Nielsen, Introduction to Optimization and Data Fitting. DTU IMM, 1st Edition, 2010.
- [3] Hans Bruun Nielsen, *Checking Gradients*. DTU IMM, 1st Edition, 2000, http://www2.imm.dtu.dk/~hbn/Software/checkgrad.ps.