## Recursion relation

## December 27, 2023

```
[]: Find the value of T(2) for the recurrence relation T(n) = 3T(n-1) + 12n, given
      \hookrightarrowthat T(0) = 5.
     # Recurrence Relation:
     \# T(n) = 3T(n-1) + 12n
     # Initial Condition:
     # T(0) = 5
     # Calculate T(1):
     \#T(1) = 3 * T(0) + 12 * 1
          # = 3 * 5 + 12
         # = 27
     # Calculate\ T(2):
     \#T(2) = 3 * T(1) + 12 * 2
         # = 3 * 27 + 24
          #= 105
     # Therefore, the value of T(2) for the given recurrence relation is 105.
     #2 Given a recurrence relation, solve it using the substitution method:
     \# a. T(n) = T(n-1) + c
     # Substitution method
     \# T(n) = T(n-1) + c
     # Assumption: T(k) = T(k-1) + c
     # Substitute: T(n) = T(n-2) + 2c
     # Repeat until base case (assume base case is T(0))
     # T(n) = T(0) + nc
     # b. T(n) = 2T(n/2) + n
```

```
# Substitution method
\# T(n) = 2T(n/2) + n
# Assumption: T(k) = 2T(k/2) + k
# Substitute: T(n) = 2^2 T(n/2^2) + n + n
\# Repeat until base case (assume base case is T(1))
\#T(n) = 2^{(\log 2(n))} T(1) + n \log_{2}(n)
\# c. T(n) = 2T(n/2) + c
# Substitution method
\# T(n) = 2T(n/2) + c
# Assumption: T(k) = 2T(k/2) + c
# Substitute: T(n) = 2^2 T(n/2^2) + 2c
# Repeat until base case (assume base case is T(1))
\#T(n) = 2^{(\log 2(n))} T(1) + c (\log 2(n) - 1)
\# d. T(n) = T(n/2) + c
# Substitution method
# T(n) = T(n/2) + c
# Assumption: T(k) = T(k/2) + c
# Substitute: T(n) = T(n/2^2) + 2c
# Repeat until base case (assume base case is T(1))
#T(n) = T(1) + c \log_2(n)
# 3. Given a recurrence relation, solve it using the recursive tree approach:
# a. T(n) = 2T(n-1) + 1
# Recursive Tree Approach:
# Start with T(n).
# At each level, multiply by the coefficient of the recurrence term (2) and addu
\hookrightarrow the constant term (1).
# Continue this process until reaching the base case.
# Step 1:
\# T(n) = 2T(n-1) + 1
# Step 2:
```

```
# T(n) = 2[2T(n-2) + 1] + 1
# In general:
# T(n) = 2^(n-1) T(1) + (n-1)
# b. T(n) = 2T(n/2) + n

# Recursive Tree Approach:
# Start with T(n).
# At each level, multiply by the coefficient of the recurrence term (2) and add_____the constant term (n).
# Continue this process until reaching the base case.

# Step 1:
# T(n) = 2T(n/2) + n

# Step 2:
# T(n) = 2[2T(n/4) + n/2] + n

# In general:
# In general:
# T(n) = n log2(n) + n
```

[]: