

gcamreport: An R tool to process and standardize GCAM outputs

Clàudia Rodés-Bachs¹, Jon Sampedro¹, Dirk-Jan Van de Ven¹, Ryna Yiyun Cui², Alicia Zhao², Matthew Zwerling², and Zarrar Khan³

¹ Basque Centre for Climate Change (BC3), Leioa, Spain ² Center for Global Sustainability, School of Public Policy, University of Maryland, College Park, MD, USA ³ Pacific Northwest National Laboratory's Joint Global Change Research Institute, College Park, MD, USA

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))

Summary

There is an urgent need for multi-model studies to characterize uncertainty arising from model heterogeneity. These studies aim to build a more reliable and transparent framework, informing policymakers in the design and implementation of climate policies (Guivarch et al., 2022). In response to this challenge, multiple institutes and organizations have adopted the standardized data template developed by the Integrated Assessment Modeling Consortium (IAMC). This template is maintained by the International Institute for Applied Systems Analysis (IIASA) and aims to standardize and facilitate model intercomparison exercises. For the latest Assessment Report (AR6), the Intergovernmental Panel on Climate Change (IPCC) required all contributors to homogenize their data to enable comparisons and ensure full transparency (Krey et al., 2014). This practice has set the foundation for a new open management of the outputs in the area of global scenario analysis.

In the case of the Global Change Analysis Model (GCAM) (Calvin et al., 2019), a well-regarded model that has been extensively used for different international and national scenario analysis, the harmonization code has never been documented nor standardized, making it difficult to reproduce outputs and hindering the transparency of results. To overcome these limitations, we have developed gcamreport, an R package that systematizes the transformations of GCAM outputs, generates figures to facilitate the analysis of the results, and allows user interaction with the produced outputs. Furthermore, the tool can be used embedded in a Docker image, which allows users to use the package in a virtual environment without having to install any specific software or library. Finally, each gcamreport release is linked to either a version of GCAM or a study in which GCAM was used, ensuring reproducibility, interoperability, accessibility, and findability, which is in line with the well-known open science principles FAIR and TRUST (Lin et al., 2020; Wilkinson et al., 2016).




Figure 1: Structure of the gcamreport package.

Statement of need

Integrated assessment models (IAMs) are the dominant tools for global scenario analysis and for exploring system-wide implications of climate policies (Van Beek et al., 2020). They must be transparent and open in order to be an effective instrument for informing decision making. In addition, multi-model studies should be considered to strengthen the consistency and robustness of the policy recommendations (Nikas et al., 2021). However, it is not always easy to compare outputs from different models, as it involves a laborious task of bringing together modeling communities with different backgrounds. To overcome these handicaps, the IAMC developed a time-series data template (Huppmann et al., 2023) which has been used in prominent multi-IAM studies, such as the presentation of the Shared Socio-economic Pathways (SSPs) (O'Neill et al., 2014), the design of the Representative Concentration Pathways (RCPs) (Moss et al., 2010), and the last two IPCC Assessment Reports (AR5 (Masson-Delmotte et al., 2021) and AR6 (Shukla, J. et al., 2022)), as well as the IPCC Special Report on Global Warming of 1.5°C (Masson-Delmotte et al., 2018) and the recent climate scenarios of the Network for Greening the Financial System (NGFS) (Richters et al., 2022). For these kind of assessments, it is essential that the outputs from the different models meet strict requirements and are reported in a standardized way to facilitate the comparison and multi-analysis (Skea et al., 2021).

One of the most extensively applied integrated assessment model is the Global Change Analysis Model (GCAM) (Calvin et al., 2019). GCAM is an open-source multi-sector model developed at the Joint Global Change Research Institute (JGCRI) designed to explore the linkages between energy, water, land, climate, and economics within a single computational system. It enables users to explore potential what-if type futures by dividing the world in 32 geopolitical regions and running in a 5-year time step. GCAM is in continuous development, with each new version increasing its accuracy by better representing and detailing existing elements, while adding new ones and fixing old ones (Binsted et al., 2022; Sampedro et al., 2022). Hence, the code to transform GCAM outputs into suitable outputs for model intercomparison should be adapted to each particular version/study. Otherwise, it becomes difficult to reproduce, automatize, and track changes (Stodden et al., 2018).

Here we present `gcamreport`, a powerful R-tool aligned with the principles of open science (Lin et al., 2020; Wilkinson et al., 2016), to guarantee the transparency of the produced outcomes by transforming GCAM outputs to the IAMC template requirements, making the results directly applicable to multi-IAM studies (including upcoming IPCC reports). Moreover, to overcome the limitations to reproduce the execution environment (e.g., packages or libraries), `gcamreport` is also available within a Docker image, which already contains all the required packages and facilitates the usage, ensuring interoperability and reproducibility (Boettiger, 2015).

Finally, to support the analysis of the model results, `gcamreport` includes a user-friendly interface that allows to visualize the standardized outputs in tabular format, allowing the user to subset the results by regions, scenarios, models, variables, and years, along with the option to download the desired results. Moreover, the `gcamreport` user interface generates plots of the selected variables within the same category, aggregated by region and/or sub-variables, which can also be downloaded for more in-depth analysis.

Functionality

A brief description

The `gcamreport` package is accessible online at the public domain <https://github.com/bc3LC/gcamreport>. To run it to generate and save the dataset meeting the IAMC requirements and to launch the user interface, you can follow a detailed [step-by-step tutorial](#) or this simplified code once the package has been cloned:

```
# 1) open the gcamreport project
```

```
# 2) load the package  
devtools::load_all()
```

```
# 3) run the main function to produce the dataset report meeting the IAMC  
# requirements and launch the user interface  
db_path <- "path/to/my-gcam-database"  
db_name <- "my-gcam-database"  
prj_name <- "my-project.dat"  
scen_name <- "name of the GCAM scenarios"  
generate_report(db_path = db_path, db_name = db_name, prj_name = prj_name,  
scenarios = scen_name, final_year = 2100, save_output = TRUE, launch_ui = TRUE)
```

81 The additional instructions to run the package through a Docker image and using further
82 functionalities can be found in the [online documentation](#) for all the release versions.

83 A note on the architecture

84 The gcamreport package consists of a set of functions divided into two different blocks. All
85 functions are described in a specific [documentation](#) supplemented by detailed tutorials.

- 86 ■ Dataset generation block: Transformation and saving of outputs from a created or
87 pre-loaded GCAM database to meet the IAMC formatting requirements.
- 88 ■ Interactive user block: Dataset visualization and manipulation, and figures generation.

89 To read the raw GCAM outputs, gcamreport uses [rgcam](#). This R package is part of the
90 GCAM-ecosystem, which is a set of complementary tools to the GCAM model that extend its
91 functionality and facilitate data management. In particular, rgcam allows to both read and
92 create a manageable data file containing the desired model outputs selected from the GCAM
93 results database.

94 For the correct performance of gcamreport internal functions, the package includes user-
95 modifiable constant values and mapping files, such as regional and sectorial aggregations,
96 as well as external files to compute additional calculations to produce some results that are
97 beyond the regular GCAM reporting (e.g., installed capacity). These additional calculations
98 are necessary to meet the IAMC reporting requirements (since not all models provide the same
99 type of outputs, but often through back-on-the-envelope calculations, harmonizing outputs
100 in this way improves the comparability of model results). Mapping files are automatically
101 loaded and used, but they can be customized and adjusted by the user if required by a specific
102 study as described in a detailed [R vignette](#). If the mapping files or the reporting template are
103 modified, it is recommended to save and tag the final used version to allow reusability and
104 reproducibility.

105 The output generated by the functions within the *dataset generation block* consists of an
106 RData dataset, a Comma-Separated Values (CSV) file, and a Microsoft Excel Open XML
107 Spreadsheet (XLSX), all of which are automatically saved unless the user specifies otherwise
108 with the `save_output` parameter. This ensures portability and format compatibility of the
109 output ([Krey, 2023](#)). Besides, the output generated by the *interactive user block* are both
110 CSV files and Portable Network Graphic (PNG) files, corresponding to the processed tabular
111 data and the generated figures, which are only saved if the user clicks on the corresponding
112 *download* button. This last block can be directly run with the RData dataset generated by the
113 former block, avoiding the need to re-create the reporting dataset, as detailed in this [example](#).




Figure 2: (left) Interactive user interface visualization example: tabular example and plotting example. (right) Usage workflow scheme.

Despite the optimization, the package still requires some performing time, mostly in order to create the project file from the given database. If the project file is already given, it takes less than a minute to produce the reporting dataset.

Lastly, the package will be updated simultaneously with the official GCAM releases, promoting the reusability and transparency. To keep track of the versions, there will be a gcamreport release by each GCAM version, tagged and named as documented in this [protocol](#). In the same line, some new features are planned for next releases, such as the reporting of hydrogen or refinery investments, or the generation of other types of plots in the user interface, in order to make the package more flexible and better adaptable to the needs of the scientific community. There are also plans to integrate the package into the GCAM-ecosystem suite of tools. Combining gcamreport with other tools in the ecosystem (such as [gcamextractor](#)) (Waite et al., 2023) will allow GCAM users to obtain the model outputs in different formats, depending on their scientific needs.

Acknowledgements

C.R. and D.V. acknowledges financial support from the European Union's Horizon research program under grant agreement 101056306 (IAM COMPACT project).

References

- Binsted, M., Iyer, G., Patel, P., Graham, N. T., Ou, Y., Khan, Z., Kholod, N., Narayan, K., Hejazi, M., & Kim, S. (2022). GCAM-USA v5. 3_water_dispatch: Integrated modeling of subnational US energy, water, and land systems within a global framework. *Geoscientific Model Development*, 15(6), 2533–2559.
- Boettiger, C. (2015). An introduction to docker for reproducible research, with examples from the r environment. *ACM SIGOPS Operating Systems Review*, 49(1), 71–79. <https://doi.org/10.1145/2723872.2723882>

- Calvin, K., Patel, P., Clarke, L., Asrar, G., Bond-Lamberty, B., Cui, R. Y., Di Vittorio, A., Dorheim, K., Edmonds, J., Hartin, C., Hejazi, M., Horowitz, R., Iyer, G., Kyle, P., Kim, S., Link, R., McJeon, H., Smith, S. J., Snyder, A., ... Wise, M. (2019). GCAM v5.1: Representing the linkages between energy, water, land, climate, and economic systems. *Geoscientific Model Development*, 12(2), 677–698. <https://doi.org/10.5194/gmd-12-677-2019>
- Guivarch, C., Le Gallic, T., Bauer, N., Fragkos, P., Huppmann, D., Jaxa-Rozen, M., Keppo, I., Kriegler, E., Krisztin, T., Marangoni, G., Pye, S., Riahi, K., Schaeffer, R., Tavoni, M., Trutnevyte, E., Van Vuuren, D., & Wagner, F. (2022). Using large ensembles of climate change mitigation scenarios for robust insights. *Nature Climate Change*, 12(5), 428–435. <https://doi.org/10.1038/s41558-022-01349-x>
- Huppmann, D., Wienpahl, L., Hackstock, P., & Castella, L. (2023). *Nomenclature* (Version v0.9.1). Zenodo. <https://doi.org/10.5281/zenodo.7956229>
- Krey, V. (2023). *IAMC Webinar: Standardization of scenario data collection via the IAMC data template: Past developments and future challenges*. <https://www.iamconsortium.org/event/iamc-webinar-standardization-of-scenario-data-collection-via-the-iamc-data-template-past-dev>
- Krey, V., Masera, O., Blanford, G., Bruckner, T., Cooke, R., Fisher-Vanden, K., Haberl, H., Hertwich, E., Kriegler, E., & Mueller, D. (2014). *Annex 2-metrics and methodology*.
- Lin, D., Crabtree, J., Dillo, I., Downs, R. R., Edmunds, R., Giarretta, D., De Giusti, M., L'Hours, H., Hugo, W., & Jenkyns, R. (2020). The TRUST principles for digital repositories. *Scientific Data*, 7(1), 144.
- Masson-Delmotte, V., Pörtner, H.-O., Skea, J., Zhai, P., Roberts, D., Shukla, P. R., Pirani, A., Pidcock, R., Chen, Y., Lonnoy, E., Moufouma-Okia, W., Péan, C., Connors, S., Matthews, J. B. R., Zhou, X., Gomis, M. I., Maycock, T., Tignor, M., & Waterfield, T. (2018). *An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty*.
- Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., & Gomis, M. (2021). Climate change 2021: The physical science basis. *Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, 2.
- Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., Van Vuuren, D. P., Carter, T. R., Emori, S., Kainuma, M., & Kram, T. (2010). The next generation of scenarios for climate change research and assessment. *Nature*, 463(7282), 747–756.
- Nikas, A., Gambhir, A., Trutnevyte, E., Koasidis, K., Lund, H., Thellufsen, J. Z., Mayer, D., Zachmann, G., Miguel, L. J., Ferreras-Alonso, N., Sognaes, I., Peters, G. P., Colombo, E., Howells, M., Hawkes, A., Van Den Broek, M., Van De Ven, D. J., Gonzalez-Eguino, M., Flamos, A., & Doukas, H. (2021). Perspective of comprehensive and comprehensible multi-model energy and climate science in Europe. *Energy*, 215, 119153. <https://doi.org/10.1016/j.energy.2020.119153>
- O'Neill, B. C., Kriegler, E., Riahi, K., Ebi, K. L., Hallegatte, S., Carter, T. R., Mathur, R., & Van Vuuren, D. P. (2014). A new scenario framework for climate change research: The concept of shared socioeconomic pathways. *Climatic Change*, 122, 387–400.
- Richters, O., Bertram, C., Kriegler, E., Al Khourdajie, A., Cui, R., Edmonds, J., Hackstock, P., Holland, D., Hurst, I., & Kikstra, J. (2022). *NGFS Climate Scenarios Data Set Version 3.0*.
- Sampedro, J., Iyer, G., Msangi, S., Waldhoff, S., Hejazi, M., & Edmonds, J. A. (2022). Implications of different income distributions for future residential energy demand in the

- 190 US. *Environmental Research Letters*, 17(1), 014031.
- 191 Shukla, J., Skea, R., Slade, A., Al Khourdajie, R., R. van Diemen, D. McCollum, M. Pathak, S.
192 Some, P. Vyas, R. Fradera, M. Belkacemi, A. Hasija, G. Lisboa, S. Luz, J. Malley, & et al.
193 (2022). Climate change 2022 mitigation of climate change. *Cambridge University Press*,
194 Cambridge, UK and New York, NY, USA. <https://doi.org/10.1017/9781009157926>
- 195 Skea, J., Shukla, P., Al Khourdajie, A., & McCollum, D. (2021). Intergovernmental panel
196 on climate change: Transparency and integrated assessment modeling. *WIREs Climate*
197 *Change*, 12(5). <https://doi.org/10.1002/wcc.727>
- 198 Stodden, V., Seiler, J., & Ma, Z. (2018). An empirical analysis of journal policy effectiveness for
199 computational reproducibility. *Proceedings of the National Academy of Sciences*, 115(11),
200 2584–2589.
- 201 Van Beek, L., Hajer, M., Pelzer, P., Van Vuuren, D., & Cassen, C. (2020). Anticipating
202 futures through models: The rise of Integrated Assessment Modelling in the climate
203 science-policy interface since 1970. *Global Environmental Change*, 65, 102191. <https://doi.org/10.1016/j.gloenvcha.2020.102191>
- 204
- 205 Waite, T., Pressburger, L., Zhao, M., & Khan, Z. (2023). *JGCRI/gcamextractor: v1.1.0*
206 (Version v1.1.0). Zenodo. <https://doi.org/10.5281/zenodo.8132290>
- 207 Wilkinson, M. D., Dumontier, M., Aalbersberg, Ij. J., Appleton, G., Axton, M., Baak, A.,
208 Blomberg, N., Boiten, J.-W., Silva Santos, L. B. da, & Bourne, P. E. (2016). The FAIR
209 guiding principles for scientific data management and stewardship. *Scientific Data*, 3(1),
210 160018. <https://doi.org/10.1038/sdata.2016.18>