





Warning, missmatch of prediction

and supplier information

# **Methylation profiling report**

# **Supplier information**

Supplier diagnosis:

Sample identifier: sampleName1547108512 **Automatic prediction** Sentrix ID: **EPIC** 203064760015 R01C01 Array type: Material type: **KRYO DNA** Material type: **KRYO DNA** Gender: Gender: female NA

prediction not available

Brain tumor methylation classifier results (v11b4)

Methylation classes (MCs with score >= 0.3)  Calibra		Calibrated score	Interpretation		
methylation class family Glioblastoma	a, IDH wildtype	0.99	match	<b>~</b>	
MC family members with sco	MC family members with score >= 0.1				
methylation class glioblastoma	, IDH wildtype, subclass RTK II	0.82	match	•	
methylation class glioblastoma	, IDH wildtype, subclass RTK I	0.13			
Legend: ✓ Match (score >= 0.9) X No match (score < 0.9): possibly still relevant for low tumor content and low DNA					


#### Class descriptions

Methylation class family Glioblastoma, IDH wildtype: The methylation class family "Glioblastoma, IDH wildtype" comprises the methylation classes glioblastoma, IDH wildtype, subtype RTK I to III, glioblastoma, IDH wildtype, subtype mesenchymal, glioblastoma, IDH wildtype, subtype MYCN and glioblastoma, IDH wildtype, subtype midline.

Methylation class glioblastoma, IDH wildtype, subclass RTK II: The methylation class "glioblastoma, IDH wildtype, subclass RTK II" is comprised of tumors with a histological diagnosis of glioblastoma. IDH wildtype and rarely gliosarcoma. IDH wildtype. These tumors are typically located in the cerebral hemispheres. Median age is 61 years (range 36 to 86). Recurrent chromosomal alterations are gain of chromosome 7 with or without EGFR amplification (>90%), loss of 9p21 (CDKN2A/B; >70%) and chromosome 10 loss (>90%). Gin of chromosome 19 and 20 is also recurrently observed (40% of cases). Expression profiles often resemble the 'Classical' subgroup according to the TCGA classification.

Methylation class glioblastoma, IDH wildtype, subclass RTK I: The methylation class "glioblastoma, IDH wildtype, subclass RTK I" is comprised of tumors with a histological diagnosis of glioblastoma, IDH wildtype. The tumors are located in the cerebral hemispheres. Median age is 64 years (range 29 to 84). Recurrent chromosomal alterations are gain of chromosome 7 with or without EGFR amplification (>80%), loss of 9p21 (CDKN2A/B: >50%) and chromosome 10 loss (>70%). Amplifications of the PDGFRA oncogene are enriched in this class (present in 20-30% of cases). Expression profiles often resemble the 'Proneural' subgroup according to the TCGA classification.

## Copy number variation profile



Depiction of chromosome 1 to 22 (and X/Y if automatic prediction was successful). Gains/amplifications represent positive, losses negative deviations from the baseline. 29 brain tumor relevant gene regions are highlighted for easier assessment. (see Hovestadt & Zapatka, <a href="http://www.bioconductor.org/packages/devel/bioc/html/conumee.html">http://www.bioconductor.org/packages/devel/bioc/html/conumee.html</a>)

### **MGMT promotor methylation (MGMT-STP27)**



**Disclaimer** 

Classification using methylation profiling is a research tool under development, it is not verified and has not been clinically validated. Implementation of the results in a clinical setting is in the sole responsibility of the treating physician. Intended for non-commercial use only.

### **Run information**

Report: idat\_reportBrain\_v11b4\_sample Version 2.0 Task version:

(see Bady et al, J Mol Diagn 2016; 18(3):350-61)

Task	Versior
idat_qc	2.0
idat_predictBrain	2.1
idat_rs_gender	2.0
idat_predictMGMT	2.0
idat_cnvp	3.0
idat_reportBrain_v11b4	2.0