





Methylation profiling report GERMAN CANCER RESEARCH CENTER IN THE HELMHOLITZ ASSOCIATION

Supplier information

GCGR-E20 p13 Sample identifier:

Sentrix ID: 205715840058 R03C01

Material type: **DNA-KRYO**

Gender: NA

Supplier diagnosis: **GBM**

Automatic prediction				
Array type:		EPIC		
Material type:		DNA-KRYO		~
Gender:		female		
Legend:	✓ Ok	Supplier information or prediction not available	➤ Warning, missmat prediction and supplied information	

Brain tumor classifier results (11b4)

Methylation classes (MCs with score >= 0.3)		Score	Interpreta	tion
methylation	n class family Glioblastoma, IDH wildtype	0.98	match	~
ı	MC family members with score >= 0.1			
	mothylation class gliphlastoma, IDH wildtype, subclass DTK I	0.01	match	

methylation class gliobiastoma, IDH wildtype, subclass RTK I	0.81	match		
methylation class glioblastoma, IDH wildtype, subclass RTK II	0.17			
Match (score >= 0.9) X No match (score < 0.9): possibly still relevant for low tumor content and low DNA	Matc	h to MC family n	nember	

Legend:





quality cases.




(score >= 0.5)

Class descriptions

Methylation class family Glioblastoma, IDH wildtype: The methylation class family "Glioblastoma, IDH wildtype" comprises the methylation classes glioblastoma, IDH wildtype, subtype RTK I to III, glioblastoma, IDH wildtype, subtype mesenchymal, glioblastoma, IDH wildtype, subtype MYCN and glioblastoma, IDH wildtype, subtype


Methylation class glioblastoma, IDH wildtype, subclass RTK I: The methylation class "glioblastoma, IDH wildtype, subclass RTK I" is comprised of tumors with a histological diagnosis of glioblastoma, IDH wildtype. The tumors are located in the cerebral hemispheres. Median age is 64 years (range 29 to 84). Recurrent chromosomal alterations are gain of chromosome 7 with or without EGFR amplification (>80%), loss of 9p21 (CDKN2A/B; >50%) and chromosome 10 loss (>70%). Amplifications of the PDGFRA oncogene are enriched in this class (present in 20-30% of cases). Expression profiles often resemble the 'Proneural' subgroup according to the TCGA classification. Methylation class glioblastoma, IDH wildtype, subclass RTK II: The methylation class "glioblastoma, IDH wildtype, subclass RTK II" is comprised of tumors with a histological diagnosis of glioblastoma, IDH wildtype and rarely gliosarcoma, IDH wildtype. These tumors are typically located in the cerebral hemispheres. Median age is 61 years (range 36 to 86). Recurrent chromosomal alterations are gain of chromosome 7 with or without EGFR amplification (>90%), loss of 9p21 (CDKN2A/B; >70%) and chromosome 10 loss (>90%). Gin of chromosome 19 and 20 is also recurrently observed (40% of cases). Expression profiles often resemble the 'Classical' subgroup according to the TCGA classification.

Copy number variation profile



Depiction of chromosome 1 to 22 (and X/Y if automatic prediction was successful). Gains/amplifications represent positive, losses negative deviations from the baseline. 29 brain tumor relevant gene regions are highlighted for easier assessment. (see Hovestadt & Zapatka, http://www.bioconductor.org/packages/devel/bioc/html/conumee.html)

MGMT promotor methylation (MGMT-STP27)



(see Bady et al, J Mol Diagn 2016; 18(3):350-61)

Disclaimer

Classification using methylation profiling is a tool/website for research use only, it is not verified and has not been clinically validated and, therefore, must not be used for diagnostic purposes. This tool/website is not HIPAA compliant.

Run information

Report: report_website_mnp_brain_v11b4_sample (Version 2.1)

Task version:

Task	Version
idat_preprocess	2.0.1
idat_qc	2.0.1
idat_predictBrain	2.0.1
idat_rs_gender	2.0.1
idat_cnvp	3.0.1
idat_mgmt	2.0.1
report_website_mnp_brain_v11b4_research	2.1
report_website_mnp_brain_v11b4_sample	2.1
idat_predictBrain	12.5