

# Proposal for a GDR of the Intensity Frontiers

all names

August 2, 2016

# 1 Introduction

The remarkable success of the "standard model" (SM) of particle physics in describing the particles and their strong, electromagnetic and weak interactions has nevertheless certain limitations. For example, it does not explain dark matter, dark energy, the hierarchy of the fermion masses or the matter-antimatter asymmetry in the universe. There is a general consensus in the physics community that a theory more fundamental than the standard model should exist, which is sometimes referred to as "new physics" (NP).

There are broadly two categories of searches for new physics: the energy frontier and the intensity frontier. In the former, experiments are designed to try to produce (and detect) TeV-scale particles directly, i.e. via collisions at high energy. This is the main approach currently followed by the general purpose detectors, ATLAS and CMS, at the LHC. In order to discover new particles, one is required to run at higher energies; such experiments are therefore said to be probing the energy frontier. Instead, in particle physics at the intensity frontier one probes new physics not by pushing the energy scale but rather the experiment's luminosity. This could provide signs of new physics in two ways. Either one measures SM processes for which theoretical predictions with uncertainties well under control exist, and can be compared with the experimental measurements: observing a significant discrepancy between the two would be the sign of new physics. This technique is often applied to study processes which are mediated at leading order by loop diagrams. This allows yet undiscovered particles, with masses beyond the energy of the collisions, to intervene, modifying the rates and the properties of the decay respect to the SM predictions. These measurements need to be extremely precise, so they require a large quantity of data. Apart from being a way to discover new physics, this approach also provides constraints and highlights on the nature of the new physics and eventually on its flavour structure. Alternatively one searches for processes which don't exist in the SM, and therefore a measurement automatically signifies NP. This could either probe (effective) couplings which do not exist in the SM, or particles at scales much below the energy frontier but which have not so far been seen due to the fact that they are very weakly interacting with SM particles. For example lepton flavour violating decays, axion searches or neutrinoless double beta decay.

Experimentally, the challenge with this approach is to collect a large and pure enough data sample in order to obtain evidence of NP interactions. Theoretically, it is crucial to have the description of the processes in the SM under control. Theory and experiment then need to come together to correctly interpret the experimental results in terms of theoretical predictions, to combine all the bounds produced in the different searches which hopefully eventually will lead towards the discovery of the new physics.

Historically the French community has been very active in this effort. From the experimental point of view, the focus today is on the LHCb experiment, dedicated to flavor physics and currently challenging the standard model predictions with many precise measurements; it's scope extends well beyond the realm of  $B$  physics. Worldwide, other experiments currently adopt a similar approach (NA62, MEG), some will start their data taking soon (Belle2) and other are in the preparatory phase (SHIP). From the theoretical side, those working in this field are very active both in the interpretation of current data in terms of new physics models and in the improving the precision on theoretical predictions for key observables. Given the need to compare the theoretical predictions with the experimental measurements, the interplay between theory and experiment in this field is essential. As a natural need of sharing competences and knowledge, during past years some collaborations have already risen between members of the two communities. A well known example of a fruitful exchange is the CKMfitter collaboration, which originated as a result of a French initiative. More recently, in the context of the study of rare  $B$  meson decays, three CNRS PEPS-PTI (Projet Exploratoire Premier Soutien de Physique Theorique et ses Interfaces) of one year each were proposed and accepted: NouvPhyLHCb in 2014 and PhenoBas in 2015 and 2016. These grants permitted the organization of three fruitful workshops, allowing to establish first connections and collaborations between LHCb experimentalists and theorists working on  $b \rightarrow sll$  transitions.

Following discussions among people active in the field, the need for a GDR in physics at the intensity frontier has been established. In fact, we believe that the more stable framework of a GDR would be beneficial to those working on high energy physics who are focused on the intensity frontier, in order to bring this community together and reinforce the interplay between the different research lines in the field.

The role of the GDR would be to facilitate the collaboration between different laboratories and between theorists and experimentalists, with the purpose of keeping the community in touch and informed about the latest advancements in the field, exchanging ideas and spreading knowledge. In this way the GDR will stimulate the emergence of common projects within the French community and allow it to grow. It will be a way to provide greater visibility of this large community on a national and international level. In addition, we believe that there is a real need to come together in order to discuss how research plans for the future should be shaped; including the decision of which experiments to become involved in.

We envisage the GDR to be divided into several working groups, which would both function independently and together. We have identified the following topics where there is currently activity and interest in the French community:

- **CP violation.** Since the  $B$ -factories, CP violation in the quark sector has also been proven to be a precise test of the Standard Model, through the measurement of the Unitarity Triangle. This measurement has room for improvement, and LHCb and Belle2 will provide further insight on it, as well as additional tests involving the  $B_s$  and  $b$  baryons.
- **Rare, radiative and semi-leptonic decays.** Generally mediated by loops, these decays are a powerful probe of new physics, provided that precise theoretical predictions can be made for experimentally clean observables. The large dataset collected by the LHCb experiment is currently showing the most exciting signs of slight deviations from the theoretical predictions that certainly deserves to be further analysed and deeply understood.
- **Heavy flavour production and spectroscopy.** Not only is this field an ideal framework to test the QCD predictions, but it provides crucial input needed for other measurements and interpretations in the search for physics beyond the SM. It further has recently revealed that quarks can form more complex structures than previously believed, i.e. tetraquarks and pentaquarks; the existence of these bound states has now been established though they are not yet fully understood.
- **Charm and kaon physics.** The study of kaons and charmed mesons has been at the origin of the flavor physics. Given the present experimental opportunities, a renewed interest in the analysis of their decays is emerging, as they provide complementary ways to search for new physics effects. Although for the charm physics there is already a large production of data, for the kaons some experimental challenges need to be faced and additional theoretical observables are being proposed.
- **Lepton flavour and the interplay with quark flavour.**
- **Future experiments.** It will be very beneficial for our community to discuss the future of our field, at a time where future upgrades of the LHCb experiment as well as new experiments are being proposed. This GDR could play a role in identifying the priorities for French involvement in order to continue to play an active role in the future.

In the subsequent sections of this document we will provide a brief description to highlight the interest of the working group topics, summarising their current status and the proposed near-future work of the French community.

The GDR would function through carefully planned workshops. More specifically, we plan to organize a general kick-off meeting, to bring the whole community together in order to define and consolidate collaborations and goals. This would be followed by a series of working group meetings, more intimate and focused on specific themes. These smaller meetings would allow detailed discussions and brainstorming within the specific topic of the working group, allowing close collaborations to emerge by really working together. Regularly, global workshops involving all the members of the GDR will be organized, where more general talks and discussions will be held and where we will ensure to share the advancements of the working groups and to address the connexions between them. This is particularly important since there is a clear interplay between the different working groups. For example the charm and kaon physics will have to address specific experimental and theoretical issues of the field, but the results obtained will

99 certainly have to be interpreted in the global CP violation picture and in relation with the other rare  
100 decays studies. One of the purpose of the GDR will be also to have a wider look into what is done in  
101 the same field in other countries or in experiments where the French community is not currently directly  
102 involved, but which still represent an interest for the field. Presenting ourselves as a unified community,  
103 we will aim in establishing productive interactions inviting occasionally speakers from other experiments  
104 and other particle physics GDRs, ensuring in this way that we keep the connection with the whole field.  
105 The format of these meetings would be free, and decided according to the specific needs and objectives,  
106 and it would be encouraged for younger members of the community to participate in the organisation and  
107 the discussion. In fact, we further hope to use the GDR as an opportunity to put the younger members  
108 of the community in the spotlight. One way to do so will be by giving to the postdocs the responsibility  
109 of organising and chairing the meetings. In addition, we will create an environment where PhD students  
110 would feel confident to present their work and interact with physicists from other laboratories.

## 2 CP Violation

Since its discovery in the kaon system, CP violation has been an intriguing field. A lot of efforts have been put in understanding it, as, a part from the interest of the phenomenon by itself and of its relations with the matter/antimatter asymmetry, it is a very powerful probe for NP. In fact, CP violation is particularly interesting because, in the SM, it originates from a single parameter. All the CP violating observables in the  $K$ ,  $D$  and  $B$  meson sectors are thus directly related and their combined study provides a highly powerful test of the whole SM dynamics. Instead, most model of New Physics are far less restrictive and allow for a plethora of new CP-violating sources. None of the delicate interplays between observables expected in the SM should survive to the presence of new dynamics at the TeV scale.

When experimentally testing SM predictions, it is fundamental that the theoretical precision matches the experimental accuracy. A priori, this looks challenging because CP-violation is a purely hadronic phenomenon in the SM, originating from the quark couplings. However, dedicated strategies have been designed and CP-violating observables are actually among our best windows to look through when searching for NP. For example, in CP-violating asymmetries, most of the uncertainties cancel between the numerator and the denominator, so that we can construct measurable quantities with small uncertainties. Alternatively, some observables are predicted to be so small that they can be considered forbidden in the SM, and simply observing a non-zero value would unequivocally signal the presence of New Physics.

The French community has been deeply involved in CP-violation dedicated experiments for many years (CPLEAR, NA48, BaBar, LHCb). It has been designing, building and maintaining key elements of the detectors, like trigger and calorimeters. Expertise has been developed in several key techniques needed to study CP-violation: amplitude analyses, tagged-time-dependent angular analyses, flavour tagging, neutral objects reconstruction. From the analysis point of view, the French community has been involved more specifically in the measurements of the Unitarity angles  $\alpha$ ,  $\gamma$  and  $\phi_s$ . Among other decay channels, the following have been studied in detail:  $B_{(s,d)} \rightarrow D^0 K^{*0}$ ,  $B_{(s,d)} \rightarrow \bar{D}^0 hh$ ,  $B_s^0 \rightarrow D_s K$ ,  $B_s^0 \rightarrow J/\psi \phi$ ,  $B_s^0 \rightarrow J/\psi \bar{K}^{*0}$ . One of the outcome of the work done is  $\phi_s$  is illustrated on Figure 2.

CP-violation related studies of charmless  $b$ -hadron decays are another field in which the experimental French groups have been very active in the B-factories era, and still are nowadays. These decays have a number of theoretical applications and provide a probe to new physics. For instance, the decays  $B^0 \rightarrow K_s^0 \pi^+ \pi^-$  and  $B^0 \rightarrow K_s^0 K^+ K^-$  are dominated by  $b \rightarrow q \bar{q} s$  ( $q = u, d, s$ ) loop transitions. Mixing-induced CP asymmetries in such decays are predicted to be approximately equal to those in  $b \rightarrow c \bar{c} s$  transitions, *e.g.*  $B^0 \rightarrow J/\psi K_s^0$ , by the Cabibbo-Kobayashi-Maskawa mechanism. However, the loop diagrams that dominate the charmless decays can have contributions from new particles in several extensions of the Standard Model, which could introduce additional weak phases. A time-dependent analysis of the three-body Dalitz plot allows measurements of the mixing-induced CP-violating phase. The current experimental measurements of  $b \rightarrow q \bar{q} s$  decays show fair agreement with the results from  $b \rightarrow c \bar{c} s$  decays (measuring the weak phase  $\beta$ ) for each of the scrutinised CP eigenstates. There is, however, a global trend towards lower values than the weak phase measured from  $b \rightarrow c \bar{c} s$  decays. The interpretation of this deviation is made complicated by QCD corrections, which depend on the final state and are difficult to handle. An analogous extraction of the mixing-induced CP-violating phase in the  $B_s^0$  system ( $\beta_s$ ) will, with a sufficiently large dataset, also be possible with the  $B_s^0 \rightarrow K_s^0 K^\pm \pi^\mp$  decay, which can be compared with that from, *e.g.*  $B_s^0 \rightarrow J/\psi \phi$ .

The charmless three-body analyses provide a long-term physics programs that can profit from the LHCb upgrade. In fact, these analyses proceed in increasingly complex steps, which become more and more sensitive to new physics observables with the growing dataset, and with more observed decay modes. One of the long-term goals is to perform full flavor- and time-dependent Dalitz-plot analyses of the  $B_s^0 \rightarrow K_s^0 h^+ h'^-$  modes to measure the weak phases  $\beta$  and  $\beta_s$ . Recent theoretical and experimental activity has focused on the determination of the CKM angle  $\gamma$  from charmless  $B$  meson decays using and refining the methods proposed in Refs. Moreover, with the upgrade of LHCb, more modes, eventually with more neutral hadrons, are being considered.

In the coming years, most of the experimental effort will go into the analysis of data collected by LHCb during run2 and in its upgrade plans.

From the theory side ... (CKMfitter, QCD challenge, precision challenge, sub-leading diagram estimation, penguin pollution) **This paragraph should be developed ...**

In the next five years, the GDR will provide the opportunity to continue the measurements started many years ago using the full run2 dataset of the LHC, and to explore new routes. In summary:

- the effort on the measurement of  $\phi_s$  will continue, with a focus on controlling the sub-leading penguins contributions;
- several additional ways to measure  $\gamma$  and  $\alpha$  will be explored, using for example charmless  $B$  decays;
- we will have the opportunity to explore the CP violation in baryons, largely produced at the LHC, and this is currently a mostly unexplored field, complementary to the meson field;
- CP violation in charm, still to be discovered, will be searched for;
- the theoretical advancement in CP violation in kaon, like lattice studies of the  $\varepsilon'$  matrix elements, will be followed, together with the achievements of the NA62 experiment, currently in data taking and aiming to measure the  $K^+ \rightarrow \pi^+ \nu \nu$  branching fraction.
- EDM: should we include the nEDM experiment? It is among the IN2P3 projects **Yes, this sentence should be developed.**

For all these items, the synergy between experimental and theoretical communities is essential, because a major discovery can not come if the uncertainties are not under control in both places. The new data coming from the LHC and soon from Belle2, as well as from NA62 will certainly allow to make important advancements in the CP violation field, and we need to ensure to provide our contribution and to correctly interpret the measurements in the global CP violation picture.




Figure 1: Left plot: 68% CL regions in  $B_s^0$  width difference  $\Delta\Gamma_s$  and weak phase  $\phi_s$  obtained from individual and combined CDF, DØ, ATLAS, CMS and LHCb likelihoods of  $B_s^0 \rightarrow J/\psi\phi$ ,  $B_s^0 \rightarrow J/\psi KK$ ,  $B_s^0 \rightarrow J/\psi\pi\pi$  and  $B_s^0 \rightarrow D_s^+ D_s^-$ . The expectation within the Standard Model is shown as the black rectangle. Right plot: the current status of the CKM unitarity triangle test from the CKMfitter collaboration.

### 3 Rare, radiative and semileptonic $B$ decays

Rare decays of  $B$  mesons are natural candidates to be studied within the indirect approach: being dominated by loop and box diagrams, they are very suppressed in the Standard Model, and so the most sensitive to variations due to new physics. Among them, radiative  $B$  decays, with the emission of a virtual or real photon, as well as semileptonic  $B$  decays are extremely interesting, as the theory has the instruments to perform very precise predictions. A joint work between experimentalists and theorists has allowed to identify an ensemble of observables which are at the same time sensitive to the couplings to different possible sources of New Physics (NP) and as immune as possible to non factorizable QCD effects. With the abundance of data produced by the LHC collisions, for the first time some of these decays can be observed, their properties measured and the prediction tested with better precision than ever.

For example, the LHCb collaboration has produced a large set of results, related to the exclusive  $b \rightarrow s \ell \ell$  decay modes, which are currently dominating the field. In the special case of the  $\mathcal{B}(B_s \rightarrow \mu\mu)$ , the CMS collaboration is also significantly contributing and after combining the results of the two experiments, it turned out that the long searched  $\mathcal{B}(B_s \rightarrow \mu\mu)$  is only slightly lower than, but compatible with, the value predicted in the Standard Model (SM). The  $B_d \rightarrow \mu\mu$  decay has also been seen, its branching ratio turning out to be compatible with the value predicted by the SM.

On the other hand, after comparing the experimental values of  $B \rightarrow K^* \mu\mu$  angular observables, as well as of  $\mathcal{B}(B \rightarrow K \mu\mu)$  and  $\mathcal{B}(B_s \rightarrow \phi \mu\mu)$ , with the theoretical estimates derived in the SM, one finds considerable discrepancies, of the order of 3 to 3.5 standard deviations. The LHCb results of the  $B^0 \rightarrow K^* \mu\mu$  angular analysis were also recently confirmed by the BELLE collaboration. It appears, however, that the most significant discrepancies occur near the charm production threshold, a region notoriously difficult for the theoretical description of these decays. In fact in this region it is required an accurate estimate of the hadronic matrix element of a non-local operator corresponding to disconnected  $c\bar{c}$ -diagrams, which cannot be computed by means of numerical simulations of QCD on the lattice. For that reason, as of now, it is not clear whether the current discrepancies are due to the lack of theoretical control of the  $c\bar{c}$  contributions, or they indicate the presence of NP couplings. If the second option is adopted, the angular observables of  $B \rightarrow K^* \mu\mu$  and  $B_s \rightarrow \phi \mu\mu$  decay modes provide very stringent constraints on the scenarios of NP.

On top of the very rich set of results involving muons, LHCb has also performed an angular analysis of the  $B^0 \rightarrow K^* e e$  decay mode in the low dilepton invariant mass region. The results found are in agreement with SM but currently quite limited in statistics. With more data coming, the electron channels will be more and more competitive and their measurements more precise. Among other observables, they actually provide a measurement of the photon polarization, as the di-electron are emitted by a virtual photon in some part of the kinematic space. This is a complementary measurement to the one of the decays involving real photons, like  $B_s \rightarrow \phi \gamma$  and  $B \rightarrow K^* \gamma$ . In the SM the photon polarization in  $b \rightarrow s \gamma$  transition is known to be left (right) for a  $b$  ( $\bar{b}$ ) quark, modulo effect of the order of 4% due to the quark masses and the emission of soft gluons. Any deviation from this precise expectation would be a clear sign of NP. In the next years precise measurements of the photon polarization in  $b \rightarrow s \gamma$  transitions are expected to come, but some challenges need to be faced, like for example the study of the resonant  $K^*$  structure, which needs a close collaboration between theorists and experimentalists.

Another experimental result which has also provoked some interest in the flavor physics community is that  $R_K = \mathcal{B}(B \rightarrow K \mu\mu)/\mathcal{B}(B \rightarrow K e e)_{\text{low-}q^2}$  was found to be  $2.6\sigma$  smaller than predicted in the SM, which suggests the violation of universality of the coupling to leptons (LFUV). Such a puzzling phenomenon should be scrutinized with higher statistics and tested in other similar situations, such as  $R_K$  at high- $q^2$ 's,  $R_{K^*}$ ,  $R_{\Lambda^{(*)}}$  at both low- and high- $q^2$ . This observation is adding to an already noted problem of  $R_{D^{(*)}} = \mathcal{B}(B \rightarrow D^* \tau \nu_\tau)/\mathcal{B}(B \rightarrow D^* \mu \nu_\mu)$  for which the experimental result, first measured at the  $B$ -factories and then confirmed at LHCb, is  $(2 \div 4)\sigma$  larger than predicted in the SM. There are very few phenomenologically viable theoretical scenarios of NP which can simultaneously explain that  $R_K^{\text{exp}} < R_K^{\text{SM}}$  and that  $R_{D^{(*)}}^{\text{exp}} > R_{D^{(*)}}^{\text{SM}}$ . To further understand the origin of the LFUV one can envisage doing the angular analysis of all the mentioned decay modes, and from the ratios of angular observables check whether or not a similar size of the LFUV is indeed observed. Furthermore, to facilitate a comparison

with theory it is more sound to compare  $B_s \rightarrow D_s^{(*)} \ell \nu_\ell$  because the theoretical uncertainty related to the chiral extrapolation in the light valence quark on the lattice is completely avoided in this way. Moreover, the emission of soft photons can differently affect  $B^- \rightarrow D^{0(*)} \ell^- \bar{\nu}_\ell$  and  $B^0 \rightarrow D^{-(*)} \ell^+ \nu_\ell$ , the modes which are usually averaged. Such a problem is much less significant if one works with  $B_s \rightarrow D_s^{(*)} \ell \nu_\ell$  decays.

Most of the models pretending to describe the LFUV effects allow for the lepton flavor violation (LFV) too. For that reason it is of great interest to measure the LFV modes such as  $B_s \rightarrow \mu \tau$ ,  $B \rightarrow K^{(*)} \mu \tau$ ,  $B_s \rightarrow \phi \mu \tau$ , which can now be probed thanks to the large statistics achievable at the LHC. The experimental bounds on  $\mathcal{B}(B_s \rightarrow \mu e)$  and  $\mathcal{B}(B_s \rightarrow K^{(*)} \mu e)$  can also be greatly improved. These results can be very useful for phenomenology of the LFV decays and for the bigger picture that could ultimately lead to a theory of flavor of quarks and leptons.

The French community is largely active both on the experimental and theoretical side on rare, radiative and semileptonic decays. In the need of exchanging results, it has been promoting in the past years the aforementioned PEPS-PTI projects and the related workshops, successfully followed by the community. They clearly establishing the need of a GDR as a regular framework were the discussions and the collaboration could take place on a more regular basis. The following topics should be addresses:

- For the LFUV, we should elaborate a general scenario of NP, in an effective field theory approach, allowing to isolate the observables which are most sensitive to the couplings to the vector (scalar) and/or axial (pseudoscalar) operators. Experimenters and theorists will elaborate on the feasibility of the distribution of  $B_{(s)} \rightarrow D_{(s)}^{*} \ell \nu_\ell$  according to the polarization of the outgoing vector meson. Furthermore, a contact with other leptonic observables should be made in order to test several plausible scenarios of NP which result in LFUV. New experimental results on the ratio measurements will be provided by LHCb, but are also expected from BelleII.
- For the angular analyses of  $b \rightarrow s \ell \ell$  transitions, with the new and more accurate experimental data, it becomes mandatory to assess the hadronic uncertainties on the theory side. Lattice QCD and the QCD sum rule practitioners will try and evaluate the size of theoretical errors and discuss the appropriate methodology on how to account for various sources of systematic uncertainties. We should explore the  $B^0 \rightarrow K^{*} \tau \tau$  channel and define new observables taking into account the direct access to the *tau* polarization. The study of  $b \rightarrow s \ell \ell$  transition in b-baryons is quite new and the identification of interesting observables for the  $\Lambda_b \rightarrow \Lambda^{*} \mu \mu$  may be interesting. Possible phenomenological ideas on how to relate the hadronic quantities in several decay modes will be discussed as they might be helpful in canceling a large part of hadronic uncertainties. Ideas on how to treat the non-resonant  $c\bar{c}$ -contributions would be very welcome. The phenomenological interpretation of the experimental results will allow to shed lights on which physics beyond standard model best fits the data.
- The LFV direct searches will soon provide new results. It will be crucial to work on a package that could include all the possible constraints relevant to LFV at low and high energy and see what are the lessons one can learn about the Yukawa sector from the data.
- Finally we should assess the current situation concerning the extraction of the Yukawa couplings from experimental data, clarifying in which way those data can be related to the low-energy physics observables and *b*-decay observables in particular. In order to address the issue of “*Which theory of flavor?*” we will try and combine the searches made at Belle II with those made at NA62 and KOTO experiments. Address the issue of (in)compatibility of the conclusions found in the Yukawa sector through the low-energy experiments with the LHC findings at the TeV-scale.






Figure 3 – Left: Bin-by-bin fit of the one-parameter scenario with a single coefficient  $C_9^{\text{NP}}$ . Right: Fit with independent coefficients  $C_9^{\text{NP}}_\mu$  and  $C_9^{\text{NP}}_e$ .

Figure 2: From arXiv:1605.06059v1 will develop the caption

## 4 Charm and Kaon Physics

In the LHC collisions, a large part of the proton-proton cross section goes into charm and strange quarks production. Indeed, the  $c\bar{c}$  cross-section is roughly 10% of the total inelastic cross-section, so that charm hadrons are produced extremely copiously. Their short but measurable decay times make them relatively simple to reconstruct and separate from background. The LHCb detector has developed dedicated triggers registering large samples of charm hadron decays and is currently exploring the best strategy to collect a large quantity of kaon decays. In fact, charm and kaon physics provides complementary insights into flavor physics to this obtained from the  $b$  sector. This working group will assess specific questions concerning the charm and kaon experimental and theoretical issues.

Precise studies of the properties and decays of charmed hadrons are motivated as almost-null tests of the Standard Model. In terms of CP violation, charm hadron decays involve only the first two generations of quarks, and the CP violation in decay is therefore expected to occur at below the per mil level in the SM. Additionally, compared to beauty or strange hadrons, the mixing of neutral charm hadrons is slow, with both the  $x = \Delta m/\Gamma$  and  $y = \Delta\Gamma/(2\Gamma)$  parameters at around the percent level. The CP violation in charm decays has not been observed so far, and the existing experimental limits are at the few per mil level. The theoretical predictions of charm CPV are difficult as long distance contributions dominate; CPV in decay close to the present experimental limits could be accommodated within the SM or could be signs of NP, and a progress on the theory side is required to disentangle the two. Similarly, in the case of mixing, or CP violation in the interference of decay and mixing, more precise experimental results are needed to stimulate progress on the theoretical predictions.

In addition, charmed hadrons are also an interesting place to study rare and forbidden transitions, for example FCNS or lepton number violating decays. The most recent studies of charmed hadron decays within the French community were performed on the rare decays  $D^0 \rightarrow \pi\mu\mu$  (with same sign muons) and  $D^0 \rightarrow K\pi\mu\mu$ . The former is of interest because the copious production rate of charmed hadrons allows effective limits to be placed on Majorana neutrinos. The latter is the charmed counterpart of  $B \rightarrow K^*\mu\mu$  and has now been observed for the first time by LHCb, albeit within a dimuon  $q^2$  region dominated by the  $\omega$  and  $\rho$  resonances. It should in principle share much of the same phenomenology of  $B \rightarrow K^*\mu\mu$ , with the complication of much higher backgrounds from decays to hadronic resonances (such as  $K\pi\rho$ ) which subsequently decay to dimuon pairs. Once that a large signal yields become available, an angular analysis will be of prior interest.

In the upcoming period, the most critical work will be the following.

- Improve the limits on CPV, both in decay and the interference of mixing and decay, as well as to make ever more precise measurements of charm mixing parameters using both the  $D \rightarrow hh$  and  $D \rightarrow K_s hh$  decay modes with the full Run II LHCb dataset.
- LHCb should obtain large samples of FCNC decays such as  $D^0 \rightarrow K\pi\mu\mu$ . Potentially this will allow for an observation of the nonresonant (in the dimuon spectrum) decay and a measurement of angular observables similar to the ones which characterise  $B \rightarrow K^*\mu\mu$ .
- Make more precise measurements of charm hadron lifetimes, in particular in the less well understood baryon sector. This could aid the development of HQE tools and techniques required to eventually obtain precise SM predictions for mixing and CPV in the charm sector.

Concerning kaon physics, it is worth to remember that kaon mixing and decays belong traditionally to the most constraining processes for physics beyond the SM. Nowadays, the discrepancies found in recent LHCb and  $B$ -factory data, in particular in the quantity known as  $R_K$ ?, provide motivations for searches of certain  $K$  decays, in particular lepton-flavor violating (LFV) ones of the kind  $K \rightarrow (\pi)e\mu$ . Interestingly, the  $R_K$  effect is consistent in magnitude and size with other discrepancies observed in the flavor sector. Without further assumptions, LFNU at a non-SM level implies LFV at a non-SM level. In fact, to account for  $R_K$  one needs to invoke new interactions distinguishing between leptons of different generations, for example lepton-lepton couplings with a new vector boson or quark-lepton couplings with a scalar leptoquark. The fermions involved in such interactions are generally not in the mass eigenbasis

– this basis doesn’t even exist at the scale of these interactions, usually above the EWSB scale. After EWSB, rotation of the quark and lepton fields to the mass eigenbasis generates LFV effects along with the LFNU ones.

One theoretically appealing way to generate new-physics shifts of the required size is to invoke an effective interaction involving dominantly quarks and leptons of the third generation ?. Then, the amount of LNU pointed to by  $R_K$  actually allows to quantify rather generally the expected amount of LFV to be in the ballpark of  $10^{-8}$ , which happens to be within reach at LHCb’s run 2. This argument, reported in Refs. ??, motivates searches of LFV  $B$  decays as a promising direction at LHCb.

This very argument has implications in  $K$  physics as well, in decays of the kind  $K \rightarrow (\pi)\ell\ell'$ , such as  $K_L \rightarrow e^\pm\mu^\mp$  and  $K^+ \rightarrow \pi^+e^\pm\mu^\mp$ . Experimental limits on these modes are more than ten years old:  $\mathcal{B}(K_L \rightarrow e^\pm\mu^\mp) < 4.7 \times 10^{-12}$  ?,  $\mathcal{B}(K^+ \rightarrow \pi^+e^-\mu^+) < 1.3 \times 10^{-11}$  ?,  $\mathcal{B}(K^+ \rightarrow \pi^+e^+\mu^-) < 5.2 \times 10^{-10}$  ?. Theoretical expectations for the above decays are straightforwardly calculable after suitably normalising the decay modes of interest in order to cancel phase-space factors. Defining  $\beta^{(K)}$  as the ratio of the new-physics Wilson coefficient responsible for the decay in the numerator over the SM Wilson coefficient responsible for the normalising decay, we get

$$\frac{\Gamma(K_L \rightarrow e^\pm\mu^\mp)}{\Gamma(K^+ \rightarrow \mu^+\nu_\mu)} = |\beta^{(K)}|^2 , \quad (1)$$

$$\frac{\Gamma(K^+ \rightarrow \pi^+\mu^\pm e^\mp)}{\Gamma(K^+ \rightarrow \pi^0\mu^+\nu_\mu)} = 4|\beta^{(K)}|^2 . \quad (2)$$

To get a numerical idea of the effects to be expected, we need a model predicting  $|\beta^{(K)}|^2$ . For the sake of definiteness, here we use “model A” of Ref. ? (any other motivated model, for example Ref. ?, will do), thereby obtaining  $|\beta^{(K)}|^2 = 2.15 \times 10^{-14}$ . Use of eqs. (1) then implies

$$\mathcal{B}(K_L \rightarrow e^\pm\mu^\mp) \approx 6 \times 10^{-14} , \quad (3)$$

where we have used  $\mathcal{B}(K^+ \rightarrow \mu^+\nu_\mu) \approx 64\%$  and  $\Gamma(K^+)/\Gamma(K_L) \approx 4.2$  ?. In addition

$$\mathcal{B}(K^+ \rightarrow \pi^+e^\pm\mu^\mp) \approx 3 \times 10^{-15} . \quad (4)$$

after use of  $\mathcal{B}(K^+ \rightarrow \pi^0\mu^+\nu_\mu) \approx 3\%$ .

While the  $K^+$  LFV mode is clearly too suppressed (within the considered model!), the  $K_L$  one, eq. (3), has a branching ratio close to  $10^{-13}$ . Such a rate may actually well be reachable at the NA62 experiment. As concerns LHCb, it should be noted that, although  $K$  mesons are produced copiously, their lifetimes are typically too long for the detector size – with the exception of the  $K_S$ . A dedicated study is thus necessary to understand the actual LHCb capabilities for the above decays.

## 5 Heavy Flavor Production and Spectroscopy

Quantum Chromodynamics (and the quark model out of which it grew) is one of the fundamental building blocks of the Standard Model. It has been extensively validated over the decades, and is very well understood. However, due to the non-perturbative behaviour that follows from the large self-coupling of low-energy gluons, the practical implications of QCD are still very much an active subject of research. This was vividly illustrated in the realm of spectroscopy recently, when the first compelling observation of pentaquark ( $qqqq\bar{q}$ ) states was made by LHCb [?] just over fifty years after their existence was predicted [?]. This discovery came as a surprise to experimentalists and theorists alike: the possible existence of such states was known, but the quark composition of quasi-stable pentaquark resonances (let alone their masses, widths, and production mechanisms) was not.

More broadly, results from QCD and strong physics are frequently needed as inputs to other measurements or to their interpretation. For example, there is considerable interest in the decays  $\bar{B} \rightarrow D^{(*)}\ell^-\bar{\nu}_\ell$ : the ratio of branching fractions (in a restricted region of phase space) for  $\ell = \mu$  and  $\ell = \tau$  can be used to test lepton universality. The current world average, combining results from LHCb, BABAR and Belle, is in tension at the  $4\sigma$  level with Standard Model expectations ?. One of the important systematic uncertainties in this measurement is associated with the spectrum and properties of excited charm resonances  $D^{**}$ , which could contaminate the final state with feed-down from  $\bar{B} \rightarrow D^{**}\ell^-\bar{\nu}_\ell$ : here, input from spectroscopy is needed for the measurement itself. There are numerous instances in which QCD input is needed for the interpretation of measurements, notably for  $B^0 \rightarrow K^{*0}\mu^+\mu^-$  in which local tensions of  $3\sigma$  were seen by LHCb in two regions of phase space [?], corresponding to an overall tension of  $3.4\sigma$  with the SM prediction of ?. The significance of the tension depends strongly upon the SM theory prediction and its uncertainties. To take a third and final example, QCD processes are an inherent background to all physics at the LHC, and in some cases Monte Carlo predictions of their spectrum need to be included in the fit itself. Tuning of the Monte Carlo models requires not only work from the phenomenology community but also measurements of production cross-sections across a range of transverse momentum and pseudorapidity.

The HEP community in France is engaged in this field, both on the experimental and theory sides. For practical reasons most experimental measurements have come from LHCb in recent years. LHCb-France has been involved on multiple fronts: spectroscopy of exotica (LAL, LAPP, LPNHE), spectroscopy of non-exotic resonances (LAL, LPNHE), and measurements of production rates (CPPM, LAL, LAPP, LPNHE). This list is not exhaustive, and there are far too many results to discuss them individually; purely by way of illustration we point to recent contributions by French groups to studies of exotic 4- and 5-quark resonances [?, ?], discoveries of two  $\Xi_b$  resonances and precise measurements of their mass splittings [?, ?], and measurements of the  $J/\psi$  production cross-section with the new 13 TeV LHC data [?]. Numerous theory groups are also actively involved, and we do not dare attempt an exhaustive list<sup>1</sup>. As well as hadron production, there is substantial French expertise in spectroscopy theory. For example, the opening theory review talk at the 2014 Workshop on Heavy Quark Baryons at LHCb<sup>2</sup>, a workshop organised by LHCb to which external experts were invited, was given by a member of IPNL. This is both recognition of this expertise and an illustration of the demand for productive theory-experiment crosstalk.

During the coming years, several analyses in this area are planned by LHCb-France and will be followed in the framework of the GDR.

- studies in beauty baryon spectroscopy following on from the observations of three  $\Xi_b$  resonances;
- searches for the doubly heavy  $\Xi_{cc}$  baryons, and measurements of production cross-sections at new centre-of-mass energies (including the 13 TeV Run-2 data and in heavy-ion collisions).
- Assuming that the  $\Xi_{cc}$  searches are successful, they in particular will lead to fruitful exchanges with theory: their masses and properties will have immediate implications for QCD models, and theory

<sup>1</sup> In the author list of one review paper of heavy flavour production alone [?], we counted eight French laboratories: IPNO, IRFU, LAL, LAPTh, LLR, LPC, LPSC, and SUBATECH.

<sup>2</sup> <https://indico.cern.ch/event/317758/>

input will be very useful for the next step, namely observing and studying their excitations.



## 7 Future Experiments

There are a large variety of future experiments at the intensity frontier planned in the coming years. Here we have divided them into two categories: those searching directly for physics beyond the Standard Model, and flavour physics experiments which can indirectly probe high energy scales through precision measurements.

### 7.1 Weakly interacting new light particles searches

In this subsection we focus on searches for weakly interacting new light particles, commonly called WISPs, of which the two canonical candidates are hidden photons and axion-like particles. Experiments to search for these are in many cases very cheap, and can often recycle older experiments.

The best motivated WISP is the QCD axion itself, which is expected to solve the strong CP problem but is associated with new physics above  $10^9$  GeV. Its mass may lie anywhere in the sub-eV range, and it is a very well-motivated dark matter candidate.

Axion-like particles (ALPs) are (pseudo)-scalars, perhaps cousins of the QCD axion but which do not obtain their masses from QCD. They are characterised by their coupling to photons in a Lagrangian term  $\mathcal{L} \supset -\frac{1}{4}g_{a\gamma\gamma}aF_{\mu\nu}\tilde{F}^{\mu\nu}$ . These are highly motivated from top-down constructions as generically arising when symmetries are broken at high scales, and also make attractive dark matter candidates. On the other hand, and perhaps most importantly, there have recently been several studies indicating possible discoveries of such particles in various astronomical observations: either as an explanation for excessive white dwarf cooling or anomalous transparency of the universe to gamma rays, and most excitingly as an explanation for the soft excess of X-rays from the coma cluster (at 200 eV) and/or the oscillatory modulation of X-rays from the Perseus cluster (and even, perhaps, an explanation for an observed 3.55 keV X-ray line). These hints all point to a very light ALP ( $< 10^{-12}$  eV) with a coupling  $g_{a\gamma\gamma} \sim \mathcal{O}(10^{-11} \div 10^{-12})\text{GeV}^{-1}$ . However, while this is a very interesting region to probe, such a particle could have a wide range of masses and couplings.

Hidden photons are new (massive) gauge bosons which mix kinetically with the visible photon via a dimensionless kinetic-mixing parameter  $\chi$ . While one motivation of these is as a possible explanation for the  $3\sigma$  discrepancy between the measured and calculated value of the muon dipole moment – this would require a hidden photon in the  $\mathcal{O}(100)$  MeV range with  $\chi \sim \mathcal{O}(10^{-3})$  – they also appear generically in top-down constructions of beyond-the-Standard Model physics. They have been proposed as perhaps the most natural force carriers for light dark matter particles, or could even make up the dark matter themselves. Intensity frontier experiments searching for these either search for the particles as dark matter or attempt to directly produce them. In the dark matter case, the assumption that there is a large abundance of particles all around us greatly enhances the reach; on the other hand, for the very light ALPs this is unlikely to be the case. The dark matter searches consist of resonant cavities, helioscopes, and now many more exotic suggestions. Direct searches are broadly photon regeneration experiments (light shining through a wall), electron colliders or beam dumps.

Important upcoming experiments include:

- Axion haloscopes (magnetic resonant cavity experiments) ADMX-HF, YMCE and WISPDMMX at the University of Washington, Yale and Hamburg respectively are all expected to report results soon, probing axion masses in the  $\mu\text{eV}$  range.
- The FUNK experiment in Karlsruhe uses a dish antenna to search for dark matter hidden photons.
- The helioscope SHIPS at Hamburg searches for hidden photons produced in the sun.
- The REAPR and ALPS-II photon regeneration experiments at Fermilab and DESY respectively will attempt to directly produce ALPs or hidden photons in the lab.
- The IAXO helioscope at CERN uses a magnetic field to search for ALPs produced in the sun. It is expected to operate over the next decade and there are significant synergies with the French community.

- The SHIP beam dump experiment using the SPS proton beam at CERN has a substantial input from French theorists and experimentalists. It has the potential to search for particles in the GeV range, which included heavier ALPs, as well as other candidates such as right-handed neutrinos. Its development will encompass much of the lifetime of the proposed GDR.
- There will be electron beam-dump experiments HPS, DarkLight and MESA at SLAC, JLab and Mainz respectively, with the latter running in about 2020. These will provide high intensity competition to hidden-photon searches in the 10 – 1000 MeV range.
- The BMV experiment at Toulouse received ANR funding in 2014 to build phase two and complement its 2007 results. It can perform photon regeneration searches; it also included an X-ray regeneration experiment.
- There is also a proposal to use the Tore Supra tokamak at Cadarache to search for ALPs. This would be particularly interesting to unite with the plasma physics community.

## 7.2 Future experimental programs related to $CP$ violation, rare decays of heavy flavours and lepton flavour violating processes

As far as  $CP$  violation and rare  $b$ -flavoured hadrons or  $\tau$  decays are concerned, the two main players at the horizon of 2025 are the upgraded LHCb experiment at CERN and the Belle II experiment at KEK. The synergy and complementarity between the two projects has been assessed clearly in the past and we should ensure within the GDR to follow the progress in both the collaborations. In this respect, we can profit of the connexions established already by members of the GDR with the KEK colleagues in the framework of the TYL/FJPPL (Franco-Japan Particle Physics Laboratory) .

Several large or medium scale projects related to Flavour Physics are envisaged to probe Beyond Standard Model Physics. Among them, there are prospective studies to educate the possibility to run the LHCb spectrometer in the High Luminosity phase of the LHC or to make use of high intensity beam lines (*e.g.* SPS and FCC injectors) with fixed target experiments.

A possible long-term strategy for high-energy physics at colliders, after the exploitation of the LHC and its High Luminosity upgrade, considers a tunnel of about 100 km circumference, which takes advantage of the present CERN accelerator complex. The Future Circular Collider (FCC) concept follows on the successful experience and outcomes of the LEP-LHC complex of experiments. A possible first step of the project is to fit in the tunnel a high-luminosity  $e^+e^-$  collider aimed at studying comprehensively the electroweak scale with centre-of-mass energies ranging from the  $Z$  pole up to beyond the  $t\bar{t}$  production threshold. A 100 TeV proton proton collider is considered as the ultimate goal of the project. Future Circular Collider study groups have been formed in a design study hosted by CERN, aiming at a Conceptual Design Report and a review cost in time for next European Strategy milestone (2018-2019). The unprecedented statistics at the  $Z$  pole ( $\mathcal{O}(10^{12-13})$   $Z$  decays) potentially delivered by the high-luminosity  $e^+e^-$  collider can be studied in particular to explore further the Flavour Physics case at large.

In that framework, several French teams, gathering small groups of experimentalists and phenomenologists, are contributing to the design study in Flavour studies. There is a Physics potential of the measurements of rare decays of  $b$ -hadrons, which can complement the anticipated results from the current and foreseen  $b$ -Physics programs (LHCb upgrade and SuperKEKB  $B$ -factory). In that respect, French contributions are mainly focused on rare electroweak penguins which are likely unique to the FCC:  $B^0 \rightarrow K^*(892)\tau^+\tau^-$  and  $B_s \rightarrow \tau^+\tau^-$ . The large statistics at the  $Z$  pole can be used as well to scrutinize in particular Lepton Flavour Violating (LFV)  $Z$  decays, which would serve as an indisputable evidence for New Physics if seen. Heavy right-handed neutrals are natural candidates to explain LFV phenomena. They can be as well searched for directly at FCC- $ee$ . A number of low energy experiment are addressing this very question through the search for LFV by muon capture on nuclei (*e.g.* COMET in Japan) or the radiative decay of large ensemble of muons (*e.g.* MEG in Europe).

One objective of the GDR is to address the complementarity of these high intensity machines, at large scale apparatus or low-energy experiments.



## 493 8 Conclusion

494 The intensity frontier is a strategic approach to search for new physics. Its validity is recognized at an  
495 international level and it is a domain in which the French particle physics community is traditionally very  
496 active. Interesting and puzzling results are being produced by the experiments currently taking data, and  
497 further more are expected to come from the new generation of experiments that are starting or being  
498 planned. Theoretical progress are ensuring the precision of the currently performed tests, and additional  
499 clean observables are under investigation.

500 The French community working in this field recognizes the need of coming together to pursue these  
501 searches together with a renewed enthusiasm and with a stronger collaboration between the experimental  
502 and the theoretical laboratories in France. The GDR intensity frontier will be the place where we will be  
503 able to put together our experience, share our knowledge, reinforce bounds and inspire new collaborations,  
504 ensuring that the French community continues to be competitive and focused on the most appealing topics  
505 of the field. It will additionally provide a forum to discuss the future of the field, and naturally promote  
506 the emergence of a young and dynamic generation of physicists. All this will allow to keep the current  
507 involvement and acquire an even higher visibility at a national and international level.