
KEGG enrichment anchored on >=2 sc_10x celltypes top30 (up_in_KO)
Bile secretion
Glycosphingolipid biosynthesis − lacto and neolacto ser
Ferroptosis
IgSF CAM signaling
Pancreatic secretion
Antigen processing and presentation
Adherens junction
NOD−like receptor signaling pathway
Motor proteins
Salmonella infection
Spliceosome
p53 signaling pathway
Transcriptional misregulation in cancer
DNA replication
Type I diabetes mellitus
Proteoglycans in cancer
Viral myocarditis
Graft−versus−host disease
Allograft rejection
Autoimmune thyroid disease
Tight junction
Cytokine−cytokine receptor interaction
Hematopoietic cell lineage
Leukocyte transendothelial migration
ECM−receptor interaction
Virion − Hepatitis viruses
Cornified envelope formation
Cytoskeleton in muscle cells
Staphylococcus aureus infection
Cell adhesion molecule (CAM) interaction
sc10x · Absorptive enterocytes
sc10x · Dedifferentiating Paneth−like cells
sc10x · Early Differentiating Enterocytes
sc10x · Enteroendocrine Cells
sc10x · Goblet cells
sc10x · Intestinal Stem Cells
sc10x · Paneth Cells
sc10x · Paneth Cells Progenitors
sc10x · TA Cells − G2/M
sc10x · TA Cells − S
GSE109280 (MEF)
GSE28585 (MEF array)
GSE13207 (MEF array)
GSE64642 early (hMSC)
GSE64642 late (hMSC)
−log10(p.adj)
10
5
rows = anchor terms ordered by anchor ranking; grey = not enriched in that unit