

2. A curve has equation

$$y = 2x^3 - 4x + 5$$

Find the equation of the tangent to the curve at the point $P(2, 13)$.

Write your answer in the form $y = mx + c$, where m and c are integers to be found.

Solutions relying on calculator technology are not acceptable.

(5)

3

A curve has equation

$$y = 3x^2 + \frac{24}{x} + 2 \quad x > 0$$

(a) Find, in simplest form, $\frac{dy}{dx}$

(3)

(b) Hence find the exact range of values of x for which the curve is increasing.

(2)

4. The curve C has equation $y = 6 - 3x - \frac{4}{x^3}$, $x \neq 0$

(a) Use calculus to show that the curve has a turning point P when $x = \sqrt{2}$ (4)

(b) Find the x -coordinate of the other turning point Q on the curve. (1)

(c) Find $\frac{d^2y}{dx^2}$.

(d) Hence or otherwise, state with justification, the nature of each of these turning points P and Q .

(3)

5. The curve with equation

$$y = x^2 - 32\sqrt[3]{x} + 20, \quad x > 0$$

has a stationary point P .

Use calculus

(a) to find the coordinates of P ,

(6)

(b) to determine the nature of the stationary point P .

(3)

6

A curve has equation $y = g(x)$.

Given that

- $g(x)$ is a cubic expression in which the coefficient of x^3 is equal to the coefficient of x
- the curve with equation $y = g(x)$ passes through the origin
- the curve with equation $y = g(x)$ has a stationary point at $(2, 9)$

(a) find $g(x)$,

(7)

(b) prove that the stationary point at $(2, 9)$ is a maximum.

(2)

7. The curve C has equation $y = f(x)$ where

$$f(x) = ax^3 + 15x^2 - 39x + b$$

and a and b are constants.

Given

- the point $(2, 10)$ lies on C
- the gradient of the curve at $(2, 10)$ is -3

(a) (i) show that the value of a is -2

(ii) find the value of b .

(4)

(b) Hence show that C has no stationary points.

(3)

(c) Write $f(x)$ in the form $(x - 4)Q(x)$ where $Q(x)$ is a quadratic expression to be found.

(2)

(d) Hence deduce the coordinates of the points of intersection of the curve with equation

$$y = f(0.2x)$$

and the coordinate axes.

(2)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

8. The curve C has equation

$$y = 3x^4 - 8x^3 - 3$$

(a) Find (i) $\frac{dy}{dx}$

$$\text{(ii)} \quad \frac{d^2 y}{dx^2} \tag{3}$$

(b) Verify that C has a stationary point when $x = 2$ (2)

(c) Determine the nature of this stationary point, giving a reason for your answer. (2)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

The curve C has equation

$$y = 5x^4 - 24x^3 + 42x^2 - 32x + 11 \quad x \in \mathbb{R}$$

(a) Find

$$(i) \quad \frac{dy}{dx}$$

$$(ii) \frac{d^2 y}{dx^2}$$

(3)

(b) (i) Verify that C has a stationary point at $x = 1$

(ii) Show that this stationary point is a point of inflection, giving reasons for your answer.

(4)

10.




Figure 1

Figure 1 shows a sketch of a curve C with equation $y = f(x)$ where $f(x)$ is a cubic expression in x .

The curve

- passes through the origin
- has a maximum turning point at $(2, 8)$
- has a minimum turning point at $(6, 0)$

(a) Write down the set of values of x for which

$$f'(x) < 0$$

(1)

The line with equation $y = k$, where k is a constant, intersects C at only one point.

(b) Find the set of values of k , giving your answer in set notation.

(2)

(c) Find the equation of C . You may leave your answer in factorised form.

(3)

11. The curve C has equation

$$y = (2x - 3)^5$$

The point P lies on C and has coordinates $(w, -32)$.

Find

- (a) the value of w ,

(2)

- (b) the equation of the tangent to C at the point P in the form $y = mx + c$, where m and c are constants.

(5)