

2. A curve C has parametric equations

$$x = 4t + 3, \quad y = 4t + 8 + \frac{5}{2t}, \quad t \neq 0$$

- (a) Find the value of $\frac{dy}{dx}$ at the point on C where $t = 2$, giving your answer as a fraction in its simplest form.

(3)

- (b) Show that the cartesian equation of the curve C can be written in the form

$$y = \frac{x^2 + ax + b}{x - 3}, \quad x \neq 3$$

where a and b are integers to be determined.

(3)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

3. A curve C has parametric equations

$$x = \frac{t^2 + 5}{t^2 + 1} \quad y = \frac{4t}{t^2 + 1} \quad t \in \mathbb{R}$$

Show that all points on C satisfy

$$(x-3)^2 + y^2 = 4$$

(3)

A diagram illustrating a closed curve C in a 2D Cartesian coordinate system. The horizontal axis is labeled x and the vertical axis is labeled y . The origin is labeled O . The curve C is an ellipse-like shape, tilted at an angle, intersecting both the x -axis and the y -axis.

Figure 3 shows a sketch of the curve C with parametric equations

(a) Show that

(b) Show that a cartesian equation of C is

where a and b are integers to be determined. (2)

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

5. The curve C has parametric equations

$$x = 3t - 4, \quad y = 5 - \frac{6}{t}, \quad t > 0$$

(a) Find $\frac{dy}{dx}$ in terms of t

(2)

The point P lies on C where $t = \frac{1}{2}$

(b) Find the equation of the tangent to C at the point P . Give your answer in the form $y = px + q$, where p and q are integers to be determined.

(3)

(c) Show that the cartesian equation for C can be written in the form

$$y = \frac{ax + b}{x + 4}, \quad x > -4$$

where a and b are integers to be determined.

(3)

This image shows a full page of blank, lined paper. It features approximately 20 evenly spaced horizontal grey lines across its entire width, providing a template for handwriting practice or general note-taking. The margins are consistent on all sides.

6.




Figure 2

The curve C_1 with parametric equations

$$x = 10\cos t, \quad y = 4\sqrt{2}\sin t, \quad 0 \leq t < 2\pi$$

meets the circle C_2 with equation

$$x^2 + y^2 = 66$$

at four distinct points as shown in Figure 2.

Given that one of these points, S , lies in the 4th quadrant, find the Cartesian coordinates of S .

(6)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Figure 2 shows a sketch of the curve C with parametric equations

The point P lies on C and has coordinates $\left(4\sqrt{3}, \frac{15}{2}\right)$.

- Give your answer as a simplified surd.

The point Q lies on the curve C , where $\frac{dy}{dx} = 0$

- (2)

8. The curve C has parametric equations

$$x = t^2 + 6t - 16 \qquad y = 6 \ln(t + 3) \qquad t > -3$$

(a) Show that a Cartesian equation for C is

$$y = A \ln(x + B) \quad x > -B$$

where A and B are integers to be found.

(3)

The curve C cuts the y -axis at the point P

(b) Show that the equation of the tangent to C at P can be written in the form

$$ax + by = c \ln 5$$

where a , b and c are integers to be found.

(4)

9. A curve C has parametric equations

$$x = 2\sin t, \quad y = 1 - \cos 2t, \quad -\frac{\pi}{2} \leq t \leq \frac{\pi}{2}$$

(a) Find $\frac{dy}{dx}$ at the point where $t = \frac{\pi}{6}$

(4)

(b) Find a cartesian equation for C in the form

$$y = f(x), \quad -k \leq x \leq k,$$

stating the value of the constant k .

(3)

(c) Write down the range of $f(x)$.

(2)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

A diagram showing a circle C in the xy -plane. The circle is tangent to the x -axis at the origin O . The y -axis is vertical and the x -axis is horizontal. The circle is centered on the y -axis.

Figure 2 shows a sketch of the curve C with parametric equations

(a) Sh that $\frac{dy}{dx} = k(\sqrt{3})\tan 2t$, where k is a constant to be determined. (5)

- Give your answer in the form $y = ax + b$, where a and b are constants.

11. The curve C has parametric equations

$$x = \sin 2\theta \quad y = \operatorname{cosec}^3 \theta \quad 0 < \theta < \frac{\pi}{2}$$

- (a) Find an expression for $\frac{dy}{dx}$ in terms of θ (3)
- (b) Hence find the exact value of the gradient of the tangent to C at the point where $y = 8$ (3)

12. The curve C has parametric equations

$$x = 2 \cos t, \quad y = \sqrt{3} \cos 2t, \quad 0 \leq t \leq \pi$$

(a) Find an expression for $\frac{dy}{dx}$ in terms of t .

(2)

The point P lies on C where $t = \frac{2\pi}{3}$

The line l is the normal to C at P .

(b) Show that an equation for l is

$$2x - 2\sqrt{3}y - 1 = 0$$

(5)

The line l intersects the curve C again at the point Q .

(c) Find the exact coordinates of Q .

You must show clearly how you obtained your answers.

(6)

13.


**Figure 6**

Figure 6 shows a sketch of the curve C with parametric equations

$$x = 2 \tan t + 1 \quad y = 2 \sec^2 t + 3 \quad -\frac{\pi}{4} \leq t \leq \frac{\pi}{3}$$

The line l is the normal to C at the point P where $t = \frac{\pi}{4}$

(a) Using parametric differentiation, show that an equation for l is

$$y = -\frac{1}{2}x + \frac{17}{2} \quad (5)$$

(b) Show that all points on C satisfy the equation

$$y = \frac{1}{2}(x-1)^2 + 5 \quad (2)$$

The straight line with equation

$$y = -\frac{1}{2}x + k \quad \text{where } k \text{ is a constant}$$

intersects C at two distinct points.

(c) Find the range of possible values for k .

(5)

14. The curve C has parametric equations

$$x = \ln t, \quad y = t^2 - 2, \quad t > 0$$

Find

- (a) an equation of the normal to C at the point where $t = 3$,

(6)

- (b) a cartesian equation of C .

(3)




Figure 1

The finite area R , shown in Figure 1, is bounded by C , the x -axis, the line $x = \ln 2$ and the line $x = \ln 4$.

- (c) Use calculus to find the exact area of the region R .

(6)

[illegible]

The figure shows a Cartesian coordinate system with a horizontal x-axis and a vertical y-axis. The origin is labeled O . A shaded region, labeled S , is located in the first quadrant. The boundary of S consists of the y-axis, the x-axis, and a curve labeled C . A straight line labeled l passes through the first quadrant, intersecting the curve C at point P and the x-axis at point Q . The region S is the area bounded by the y-axis, the x-axis, and the curve C from the y-axis to point P .

Figure 4 shows a sketch of part of the curve C with parametric equations

The point P lies on C and has coordinates $(3, 2)$.

(a) Find the x coordinate of the point Q .

The finite region S , shown shaded in Figure 4, is bounded by the curve C , the x -axis, the y -axis and the line l .

(b) Using algebraic integration, find the exact area of S .

(5)

[illegible]

16.




Figure 3

Figure 3 shows part of the curve C with parametric equations

$$x = \tan \theta, \quad y = \sin \theta, \quad 0 \leq \theta < \frac{\pi}{2}$$

The point P lies on C and has coordinates $\left(\sqrt{3}, \frac{1}{2}\sqrt{3}\right)$.

(a) Find the value of θ at the point P .

(2)

The line l is a normal to C at P . The normal cuts the x -axis at the point Q .

(b) Show that Q has coordinates $(k\sqrt{3}, 0)$, giving the value of the constant k .

(6)

The finite shaded region S shown in Figure 3 is bounded by the curve C , the line $x = \sqrt{3}$ and the x -axis.

(c) Find the exact area of region S .

(7)

17.




Figure 2

Figure 2 shows a sketch of part of the curve C with parametric equations

$$x = 1 - \frac{1}{2}t, \quad y = 2^t - 1$$

The curve crosses the y -axis at the point A and crosses the x -axis at the point B .

(a) Show that A has coordinates $(0, 3)$. (2)

(b) Find the x coordinate of the point B . (2)

(c) Find an equation of the normal to C at the point A . (5)

The region R , as shown shaded in Figure 2, is bounded by the curve C , the line $x = -1$ and the x -axis.

(d) Use integration to find the exact area of R . (6)

18.


Diagram not
drawn to scale**Figure 4**

Figure 4 shows a sketch of part of the curve C with parametric equations

$$x = 3\theta \sin \theta, \quad y = \sec^3 \theta, \quad 0 \leq \theta < \frac{\pi}{2}$$

The point $P(k, 8)$ lies on C , where k is a constant.

(a) Find the exact value of k .

(2)

The finite region R , shown shaded in Figure 4, is bounded by the curve C , the y -axis, the x -axis and the line with equation $x = k$.

(b) Show that the area of R can be expressed in the form

$$\lambda \int_{\alpha}^{\beta} (\theta \sec^2 \theta + \tan \theta \sec^2 \theta) d\theta$$

where λ , α and β are constants to be determined.

(4)

(c) Hence use integration to find the exact value of the area of R .

(6)

19.




Figure 3

The curve shown in Figure 3 has parametric equations

$$x = 6 \sin t \quad y = 5 \sin 2t \quad 0 \leq t \leq \frac{\pi}{2}$$

The region R , shown shaded in Figure 3, is bounded by the curve and the x -axis.

(a) (i) Show that the area of R is given by $\int_0^{\frac{\pi}{2}} 60 \sin t \cos^2 t \, dt$

(3)

(ii) Hence show, by algebraic integration, that the area of R is exactly 20

(3)




Figure 4

Part of the curve is used to model the profile of a small dam, shown shaded in Figure 4. Using the model and given that

- x and y are in metres
- the vertical wall of the dam is 4.2 metres high
- there is a horizontal walkway of width MN along the top of the dam

(b) calculate the width of the walkway.

(5)

Figure 6 shows a sketch of the curve C with parametric equations

The region R , shown shaded in Figure 6, is bounded by C , the x -axis and the line with equation $x = 4$

$$\int_0^a (8 - 8 \cos 4t + 48 \sin^2 t \cos t) dt$$

(5)

(4)