

1. Find

$$\int \left(8x^3 - \frac{3}{2\sqrt{x}} + 5 \right) dx$$

giving your answer in simplest form.

(4)

2. Find

$$\int \left(\frac{2}{3}x^3 - 6\sqrt{x} + 1 \right) dx$$

giving your answer in its simplest form.

(4)

3. Find

$$\int \frac{3x^4 - 4}{2x^3} \, dx$$

writing your answer in simplest form.

(4)

4. Use integration to find

$$\int_1^{\sqrt{3}} \left(\frac{x^3}{6} + \frac{1}{3x^2} \right) dx$$

giving your answer in the form $a + b\sqrt{3}$, where a and b are constants to be determined.

(5)

Figure 3 shows a sketch of part of the curve with equation

The finite region S , bounded by the x -axis and the curve, is shown shaded in Figure 3.

$$\int \left(3x - x^{\frac{3}{2}} \right) dx \tag{3}$$

(3)

Figure 3

Figure 3 shows a sketch of part of the curve C with equation

$$y = x(x + 4)(x - 2)$$

The curve C crosses the x -axis at the origin O and at the points A and B .

- (a) Write down the x -coordinates of the points A and B .

(1)

The finite region, shown shaded in Figure 3, is bounded by the curve C and the x -axis.

- (b) Use integration to find the total area of the finite region shown shaded in Figure 3.

(7)

7. (a) Find

$$\int 10x(x^{\frac{1}{2}} - 2)dx$$

giving each term in its simplest form.

(4)




Figure 2

Figure 2 shows a sketch of part of the curve C with equation

$$y = 10x(x^{\frac{1}{2}} - 2), \quad x \geq 0$$

The curve C starts at the origin and crosses the x -axis at the point $(4, 0)$.

The area, shown shaded in Figure 2, consists of two finite regions and is bounded by the curve C , the x -axis and the line $x = 9$

(b) Use your answer from part (a) to find the total area of the shaded regions.

(5)

8. A curve C has equation $y = f(x)$ where

$$f(x) = -3x^2 + 12x + 8$$

(a) Write $f(x)$ in the form

$$a(x + b)^2 + c$$

where a , b and c are constants to be found.

(3)

The curve C has a maximum turning point at M .

(b) Find the coordinates of M .

(2)




Figure 3

Figure 3 shows a sketch of the curve C .

The line l passes through M and is parallel to the x -axis.

The region R , shown shaded in Figure 3, is bounded by C , l and the y -axis.

(c) Using algebraic integration, find the area of R .

(5)

9.

A Cartesian coordinate system is shown with a horizontal x -axis and a vertical y -axis. The origin is labeled O . A parabola, labeled $y = x^2 + 2x + 2$, opens upwards. A horizontal line is labeled $y = 10$. The parabola intersects the horizontal line at two points, labeled A and B . The region between the parabola and the horizontal line, bounded by A and B , is shaded gray and labeled R .

Figure 1

The line with equation $y = 10$ cuts the curve with equation $y = x^2 + 2x + 2$ at the points A and B as shown in Figure 1. The figure is not drawn to scale.

- (a) Find by calculation the x -coordinate of A and the x -coordinate of B .

(2)

The shaded region R is bounded by the line with equation $y = 10$ and the curve as shown in Figure 1.

- (b) Use calculus to find the exact area of R .

(7)

This image shows a single sheet of white paper with horizontal blue or grey ruling lines. The lines are evenly spaced and run across the width of the page. There are approximately 20 lines visible. The paper appears to be a standard notebook page or a sheet of stationery.

10.

$$g(x) = 2x^3 + x^2 - 41x - 70$$

- (a) Use the factor theorem to show that $g(x)$ is divisible by $(x - 5)$. (2)

- (b) Hence, showing all your working, write $g(x)$ as a product of three linear factors. (4)

The finite region R is bounded by the curve with equation $y = g(x)$ and the x -axis, and lies below the x -axis.

- (c) Find, using algebraic integration, the exact value of the area of R . (4)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

11.




Figure 2

Figure 2 shows the line with equation $y = 10 - x$ and the curve with equation $y = 10x - x^2 - 8$

The line and the curve intersect at the points A and B , and O is the origin.

- (a) Calculate the coordinates of A and the coordinates of B .

(5)

The shaded area R is bounded by the line and the curve, as shown in Figure 2.

- (b) Calculate the exact area of R .

(7)

12.




Figure 3

Figure 3 shows a sketch of part of the curve with equation

$$y = 2x^3 - 17x^2 + 40x$$

The curve has a minimum turning point at $x = k$.

The region R , shown shaded in Figure 3, is bounded by the curve, the x -axis and the line with equation $x = k$.

Show that the area of R is $\frac{256}{3}$

(Solutions based entirely on graphical or numerical methods are not acceptable.)

(7)



Figure 2 shows a sketch of part of the curve with equation

$$y = 4x^3 + 9x^2 - 30x - 8, \quad -0.5 \leq x \leq 2.2$$

The curve has a turning point at the point A .

- (a) Using calculus, show that the x coordinate of A is 1

(3)

The curve crosses the x -axis at the points $B(2, 0)$ and $C\left(-\frac{1}{4}, 0\right)$

The finite region R , shown shaded in Figure 2, is bounded by the curve, the line AB , and the x -axis.

- (b) Use integration to find the area of the finite region R , giving your answer to 2 decimal places.

(7)

14.




Figure 2

In this question you must show all stages of your working.

Solutions relying on calculator technology are not acceptable.

Figure 2 shows a sketch of part of the curve C with equation

$$y = \frac{1}{3}x^2 - 2\sqrt{x} + 3 \quad x \geq 0$$

The point P lies on C and has x coordinate 4

The line l is the tangent to C at P .

(a) Show that l has equation

$$13x - 6y - 26 = 0 \quad (5)$$

The region R , shown shaded in Figure 2, is bounded by the y -axis, the curve C , the line l and the x -axis.

(b) Find the exact area of R . (5)

Not to scale




Figure 4 shows a sketch of part of the curve C with equation

$$y = \frac{32}{x^2} + 3x - 8, \quad x > 0$$

The line l is the normal to C at the point P .

The region R , shown shaded in Figure 4, is bounded by the line l , the curve C , the line with equation $x = 2$ and the x -axis.

(Solutions based entirely on graphical or numerical methods are not acceptable.)

(10)