Probability and Statistics

Example Problems

Example 1

Three balls are selected at random without replacement from an urn containing four green balls and six red balls. Let the random variable X denote the number of green balls drawn.

- (b) Find the value assigned to each outcome of the experiment by the random variable X. {3, 2, 2, 2, 1, 1, 1, 0}
- (c) Find the event consisting of the outcomes to which the value of 0 has been assigned by X. {RRR}

Example 2

Let X denote the random variable that gives the sum of the faces that fall uppermost when two fair dice are rolled. Find P(X = 2).

We know that there are 36 total outcomes and only 1 of those results in X = 2 (a roll of 1 and 1).

$$\frac{1}{36} = 0.03$$

Example 3

Determine whether the table gives the probability distribution of the random variable X. Explain your answer.

No, because the sum of the probabilities is less than 1.

Example 4

Find the expected value E(X) of a random variable X having the following probability distribution.

Example 5

Use the formula $C(n,x)p^xq^{n-x}$ to determine the probability of the given event.

The probability of exactly **zero** successes in **nine** trials of a binomial experiment in which $p=\frac{1}{2}$

$$C(9,0)*(\frac{1}{4})^0*(\frac{3}{4})^9=0.0751$$

Example 6

The scores on an economics examination are normally distributed with a mean of **68** and a standard deviation of **14**. If the instructor assigns a grade of A to **12**% of the class, what is the lowest score a student may have and still obtain an A?

$$100\% - 12\% = 88\%$$

Then, find 88% on the Appendix of Tables which ends up being ≈ 1.17 Next, add the multiply by the standard deviation and add the mean.

$$68 + (1.175 * 14) = 84.45$$

Distribution of Random Variables

Flip a coin three times and let X denote the number of heads.

Outcome	HHH	HHT	HTH	HTT	THH	THT	TTH	TTT
Value(x)	3	2	2	1	2	1	1	0

Binomial Distribution

$$C(n,x) * p^x * q^{n-x}$$

where...

n: Number of trials x: Number of successes

p: Chance of success

q: Chance of failure (1 - p)

Calculator Info

Given the mean, the standard deviation, find the percent in range min-max

2nd \rightarrow DISTR \rightarrow (2) normalcdf

lower, upper, μ (mean), σ (standard deviation)

Find the mean, standard deviation, mode, and median

 $STAT \rightarrow 1$ (Edit)

Fill in L1 with list and L2 with frequency list (if applicable, otherwise blank)

STAT \rightarrow (Right Arrow) CALC \rightarrow (1) 1-Var Stats

Set List: to L1 2nd \rightarrow LIST \rightarrow (1) L1 Repeat with FreqList and L2 if applicable

Mean: \overline{x}

Standard Deviation: σx

Median: Med