

Flat limit of artificial cosmology for scalar waves in Schwarzschild–de Sitter

Anil Zenginoğlu^{1,*} and Govind Arun Kumar^{1,†}

¹*Institute for Physical Science and Technology, University of Maryland, College Park, Maryland 20742, USA*

How do cosmological waveforms approximate waveforms from isolated systems for a small cosmological constant? This question is motivated by the observed small value of the cosmological constant, and also by Misner’s idea to regulate the formally singular terms introduced into the Einstein equations by conformal compactification. We investigate this question quantitatively using scalar waves on Schwarzschild–de Sitter backgrounds. A suitable flat limit requires the foliation to remain regular at the future conformal boundary. We compare bridge foliations extending from the black hole horizon to the cosmological horizon with a regular asymptotically flat limit. Comparisons of cosmological and asymptotically flat waveforms on geometrically matched data show that the flat limit is numerically well behaved and recovers the essential features of the asymptotically flat waveform. We extend Misner’s artificial cosmology by introducing a cosmological source that leaves the near-zone geometry unchanged and show that it improves agreement of the ringdown waveform but fails to capture the intermediate polynomial tail behavior accurately.

I. INTRODUCTION

Gravitational radiation from isolated systems is unambiguously defined at future null infinity, $\mathcal{I}^+$. It is therefore important to include null infinity in the computational domain when calculating waveforms. In hyperboloidal compactification, we use spacelike slices that approach cuts of null infinity and solve for conformal variables rescaled by suitable powers of the conformal factor, Ω . A difficulty in this approach is the treatment of formally singular terms at infinity. Misner proposed regulating such terms with a small positive cosmological constant [1–3]. At finite cosmological length L , we replace null infinity where $\Omega = 0$ with a cosmological horizon where $\Omega_L > 0$. This raises the question how such a construction can approximate the asymptotically flat waveform. Another motivation for this work comes from the fact that cosmological observations are consistent with a small positive cosmological constant in our Universe [4]. This gives a physical motivation for numerically investigating how well asymptotically flat waveforms approximate those in a spacetime with small positive Λ . We address this question using scalar waves in Schwarzschild–de Sitter (SdS) spacetimes.

There are various approaches to include the asymptotic boundary in a computational domain. Characteristic evolution and Cauchy–characteristic matching can now successfully calculate nonlinear binary-black-hole mergers and tails [5, 6]. The hyperboloidal approach uses spacelike slices instead [7]. Early calculations exposed difficulties related to constraints, gauge conditions, and moving boundaries [8–12]. More recently, generalized conformal field equations based on a Weyl connection and conformal geodesics have demonstrated global calculations and nonlinear black-hole scattering [13–21]. While the coordinate location of the conformal boundary is a

priori known in this approach, the problem of moving boundaries and the associated loss of resolution remains.

Scri fixing maps null infinity to a time-independent boundary by using a suitable conformal compactification on stationary hyperboloidal foliations [22], which proved very useful in black-hole perturbation theory. Time-domain Schwarzschild and Kerr calculations demonstrated accurate long evolutions, including power-law tails [23–32]. Applications also include particle-source evolutions [33, 34], second-order Kerr perturbations [35], ringdown analysis [36], and free 3+1 evolutions [37–42]. This idea has also been applied in the frequency domain to quasinormal-mode calculations [43–48], pseudospectrum analyses [49–51], and self-force calculations [52–55].

Applying scri-fixing to nonlinear Einstein equations is a difficult problem due to formally singular terms [56, 57]. Implementations have been demonstrated in spherical symmetry using conformal BSSN/Z4 systems [58–61] and dual-foliation generalized harmonic gauge [62–66]. Beyond spherical symmetry, the only implementations of hyperboloidal scri-fixing use constrained evolution [67–71]. Building a robust, free-evolution hyperboloidal code that fixes $\mathcal{I}^+$ for generic three-dimensional spacetimes remains an open problem.

Misner’s idea of using a small positive cosmological constant to regulate formally singular terms is an attractive alternative approach. This idea has only been tested in de Sitter spacetime with a Minkowski limit by Misner and collaborators [1–3]. The Minkowski case is special due to the Huygens’ principle according to which the support of the solution is confined to the light cone. In this paper, we study Misner’s proposal in the presence of a black hole. We investigate the flat limit of artificial cosmology in the context of SdS spacetimes using scalar waves as a toy problem. For $\Lambda > 0$, the future conformal boundary is spacelike, with different asymptotic symmetries, radiative degrees of freedom, and fluxes from those at $\mathcal{I}^+$ [72–74]. We ask: how does a cosmological calculation approximate an asymptotically flat solution? We compare minimal and conformal scalar coupling, $\xi = 0$ and $1/6$, separating common regulator behavior from de-

* Corresponding author: anil@umd.edu

† govind02@umd.edu

pendence on the asymptotic field equation [75].

The flat limit changes the global causal and conformal structure. Writing $\Lambda = 3/L^2$ and holding M fixed, the SdS metric approaches Schwarzschild as $L \rightarrow \infty$. The asymptotic limit, however, is singular: the cosmological horizon moves to infinite areal radius, and the space-like future conformal boundary becomes null [72, 76–78]. Comparing waveforms at the outer boundary therefore requires a controlled limit of the foliation and the extraction surfaces. A regular hyperboloidal flat limit requires the limiting slices to meet distinct cuts of future null infinity, with a smooth, nondegenerate conformal spatial metric [78, 79].

As a dynamical demonstration of this singular limit, consider late-time decay. Schwarzschild fields have inverse-power tails [80], whereas SdS fields decay exponentially [75, 81, 82]. Recovering a Schwarzschild signal over a finite interval is therefore different from recovering its asymptotic decay. We test how far finite-time agreement extends as L increases. Building on the bridge construction of [79] and the SdS limiting geometry of [78], we give a quantitative time-domain test of artificial cosmology. To potentially improve Misner’s proposal, we also test a varying cosmological source supported in the asymptotic domain based on a construction by Dymnikova [83, 84].

Section II reviews the bridge foliations of SdS and their flat limit, Sec. III defines the scalar comparison, and Secs. IV and V give the uniform and exterior-supported results. The appendices present numerical details.

II. FLAT LIMIT OF SDS BRIDGES

A. SdS static patch

The SdS metric in static coordinates is

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2 d\omega^2, \quad (1)$$

$$f(r) = 1 - \frac{2M}{r} - \frac{r^2}{L^2}, \quad d\omega^2 = d\theta^2 + \sin^2 \theta d\varphi^2. \quad (2)$$

We use geometrized units $G_N = c = 1$. The coordinates are given in units of M . For $0 < M < L/(3\sqrt{3})$, f has a black-hole root r_b , a cosmological root r_c , and a negative root $r_u = -(r_b + r_c)$. They satisfy

$$L^2 = r_b^2 + r_b r_c + r_c^2, \quad 2ML^2 = r_b r_c (r_b + r_c), \quad (3)$$

$$f(r) = \frac{(r - r_b)(r_c - r)(r - r_u)}{L^2 r}. \quad (4)$$

For $i = b, c$, the surface gravities follow from (5). We use the positive algebraic coefficient $\kappa_u := |f'(r_u)|/2$ at the negative root:

$$\kappa_i = \frac{1}{2} |f'(r_i)| = \frac{1}{2} \left| \frac{1}{r_i} - \frac{3r_i}{L^2} \right|, \quad i \in \{b, c, u\}. \quad (5)$$

Using a reference radius $r_0 \in (r_b, r_c)$, the normalized tortoise coordinate is

$$r_{*,L}(r) = \frac{1}{2\kappa_b} \ln \frac{r - r_b}{r_0 - r_b} - \frac{1}{2\kappa_c} \ln \frac{r_c - r}{r_c - r_0} + \frac{1}{2\kappa_u} \ln \frac{r - r_u}{r_0 - r_u}. \quad (6)$$

The tortoise coordinate goes to $-\infty$ at r_b and to $+\infty$ at r_c . In the large- L limit, the static patch expands to the Schwarzschild exterior [78] with

$$r_b \rightarrow 2M, \quad r_c = L - M + \mathcal{O}(M^2/L). \quad (7)$$

B. Global bridge foliation

For a regular stationary foliation, we introduce a stationary height function [22, 79],

$$\tau = t + h(r), \quad B(r) := f(r)h'(r), \quad (8)$$

where B is the boost relative to the tortoise coordinate r_* . For a radial coordinate $\rho = g(r)$ and $G = d\rho/dr$, the metric becomes

$$ds^2 = -f d\tau^2 + \frac{2B}{G} d\tau d\rho + \frac{1 - B^2}{fG^2} d\rho^2 + r^2 d\omega^2. \quad (9)$$

A future-directed spacelike bridge within the static patch satisfies

$$|B| < 1, \quad B(r_b) = +1, \quad B(r_c) = -1, \quad (10)$$

At each horizon, the radial metric coefficient $(1 - B^2)/(fG^2)$ must approach a finite positive value. For each finite L , our radial map has nonzero boundary values of G , so this is equivalent to requiring a finite positive limit of $(1 - B^2)/f$. The horizons give $h \sim +r_*$ at $\mathcal{H}_b^+$ and $h \sim -r_*$ at $\mathcal{H}_c^+$ [79]. Note that these are local conditions at the horizons, and do not determine the foliation beyond $\mathcal{H}_c^+$. On the future branch of the expanding region, where $f < 0$, spacelike continuation beyond the cosmological horizon requires $B < -1$. How B continues from its horizon value $B(r_c) = -1$ determines how the leaves approach the spacelike future boundary.

Two height functions demonstrate this remaining freedom. Both are normalized by $h(r_0) = 0$ and have the same horizon data:

$$h_{\min}(r) = \frac{1}{2\kappa_b} \ln \left| \frac{r - r_b}{r_0 - r_b} \right| + \frac{1}{2\kappa_c} \ln \left| \frac{r_c - r}{r_c - r_0} \right|, \quad (11)$$

$$h_{\text{mg}}(r) = \frac{1}{2\kappa_b} \ln \left| \frac{(r - r_b)r_0}{(r_0 - r_b)r} \right| + \frac{1}{2\kappa_c} \ln \left| \frac{(r_c - r)r_0}{(r_c - r_0)r} \right| - \frac{1}{2\kappa_u} \ln \left| \frac{(r - r_u)r_0}{(r_0 - r_u)r} \right|. \quad (12)$$

The two logarithms in the minimum height function $h_{\min}$ are sufficient for horizon regularity. The second height function comes from the minimal gauge which was first

discussed in hyperboloidal QNM calculations [44–46]. Its derivation for static spherical metrics uses the in-out and out-in strategies of [85]. For SdS, the out-in construction adds the regular third-root contribution and gives a regular Schwarzschild limit [78].

Figure 1 shows how these horizon-regular slices continue toward the conformal boundary at finite L . The minimum-height slices meet at the asymptotic end. The minimal-gauge slices reach distinct cuts of the future conformal boundary. A regular hyperboloidal flat limit requires this boundary foliation to remain smooth as $L \rightarrow \infty$, with a nondegenerate conformal spatial metric [78, 79]. Below, we express this geometric requirement as an asymptotic condition on the boost.

C. Compactification and the flat limit

In addition to (11) and (12), we consider the following foliations as determined by the boost functions [79]:

$$B_{\text{lin}} = -1 + 2 \frac{r_c - r}{r_c - r_b}, \quad (13)$$

$$B_{\text{fl}} = -1 + 2 \frac{r_c - r}{r_c - r_b} \frac{r_b^2}{r^2}, \quad (14)$$

$$B_{\text{Mav}} = -\frac{1 - 3M/r}{\sqrt{1 - 9M^2\Lambda}} \sqrt{1 + \frac{6M}{r}}, \quad (15)$$

$$B_{\text{sr}} = -\gamma r + \frac{\beta}{r^2}; \quad \gamma = \frac{r_c^2 + r_b^2}{r_c^3 - r_b^3}, \quad \beta = \frac{r_c^2 r_b^2 (r_c + r_b)}{r_c^3 - r_b^3}. \quad (16)$$

The linear boost B_{lin} is a simple interpolation between the required horizon values $+1$ and -1 from (10). The modified-linear boost B_{fl} adds the factor r_b^2/r^2 to obtain a flat Schwarzschild limit to the minimal gauge as $L \rightarrow \infty$ [79]. The boost B_{Mav} was used in wave estimates on SdS by Mavrogiannis [86, 87]. The slow-roll boost B_{sr} comes from coordinates adapted to slow-roll scalar evolution around a black hole [88], also used in quantum-field correlation calculations [89, 90]. Its constants γ and β are fixed by the two horizon conditions. All these families are regular across both SdS horizons and are suitable for numerics with finite L . However, not all of them have a regular flat limit across future null infinity, which imposes an additional condition on their asymptotic behavior. To see this, map the horizons to fixed coordinate locations using

$$\rho = \frac{1 - r_b/r}{1 - r_b/r_c}, \quad r(\rho) = \frac{r_b}{1 - (1 - r_b/r_c)\rho}, \quad (17)$$

$$G(r) = \frac{d\rho}{dr} = \frac{r_b}{(1 - r_b/r_c)r^2}. \quad (18)$$

The black-hole and cosmological horizons are at $\rho = 0$ and 1 . We use subscript 0 to denote the Schwarzschild limit, $\Lambda \rightarrow 0$ ($L \rightarrow \infty$) at fixed M . At fixed r ,

$$\rho \rightarrow \rho_0 := 1 - \frac{2M}{r}. \quad (19)$$

For the compactification (17), we choose the conformal factor as

$$\Omega_L := \frac{r_b}{r} = 1 - \left(1 - \frac{r_b}{r_c}\right) \rho. \quad (20)$$

At the cosmological horizon for each finite L ,

$$\Omega_L(1) = \frac{r_b}{r_c} > 0. \quad (21)$$

In the Schwarzschild limit, $\Omega_L \rightarrow \Omega_0 = 2M/r = 1 - \rho_0$, and the outer boundary becomes $\mathcal{I}^+$, where $\Omega_0 = 0$.

To determine which horizon-regular SdS bridges approach outgoing hyperboloidal slices in Schwarzschild, we compare their fixed-radius flat limits and behavior at large radius. For each family, define $B_0(r) := \lim_{L \rightarrow \infty} B(r; L)$ with M and r held fixed. With $x := M/r$, the table below lists the limits for the six families.

bridge	$B_0(r)$	$\lim_{r \rightarrow \infty} B_0(r)$
minimum height	$3x - \frac{1}{2}$	$-\frac{1}{2}$
minimal gauge	$-1 + 8x^2$	-1
linear	$+1$	$+1$
modified linear	$-1 + 8x^2$	-1
Mavrogiannis	$-(1 - 3x)\sqrt{1 + 6x}$	-1
slow roll	$4x^2$	0

An outgoing Schwarzschild hyperboloidal slice must satisfy $B_0 \rightarrow -1$ as $r \rightarrow \infty$. Regularity of its conformal spatial metric fixes the rate of approach. With $G_0 = 2M/r^2$ and $f_0 = 1 - 2M/r$, the radial coefficient is

$$\bar{g}_{0,\rho\rho} = \Omega_0^2 \frac{1 - B_0^2}{f_0 G_0^2} = \frac{r^2(1 - B_0^2)}{f_0}. \quad (23)$$

A finite, positive boundary value requires $B_0 = -1 + aM^2/r^2 + o(M^2/r^2)$ with $a > 0$, giving $\bar{g}_{0,\rho\rho} \rightarrow 2aM^2$. This falloff also makes $h_0 + r_{*,0}$ approach a finite constant; the retarded time $t - r_{*,0} = \tau - (h_0 + r_{*,0})$ therefore labels distinct cuts of $\mathcal{I}^+$ as τ varies. These conditions exclude the minimum-height, linear, and slow-roll families. The minimal, modified-linear, and Mavrogiannis families give regular hyperboloidal limiting slices. We use the minimal gauge for its simple analytic form and regular hyperboloidal Schwarzschild limit [78, 85]. The Mavrogiannis foliation provides another suitable family.

Misner's regulator uses the positive finite- L boundary value (21) to keep the cosmological horizon at a nonvanishing conformal factor. To demonstrate the relevant scale, suppose that g satisfies $G_{ab}[g] + \Lambda g_{ab} = 0$ and define the conformal metric $\bar{g} = \Omega_L^2 g$. Then

$$G_{ab}[\bar{g}] = \mathcal{T}_{ab}[\Omega_L] - \frac{\Lambda}{\Omega_L^2} \bar{g}_{ab}, \quad (24)$$

where $\mathcal{T}_{ab}$ includes the derivative terms of Ω_L [57]. Evaluated at the cosmological horizon for each finite L , the coefficient Λ/Ω_L^2 approaches $3/(4M^2)$ as $L/M \rightarrow \infty$, while the outer boundary of the compactified domain becomes Schwarzschild future null infinity. At fixed areal radius, by contrast, this coefficient goes to zero.




FIG. 1. Conformal diagrams of two SdS bridge families that are regular at both horizons, given by (11) and (12). Curves show constant-time surfaces for each family. The dashed lines use the chart regular beyond the black-hole horizon; the solid lines use the chart regular beyond the cosmological horizon. The figure is taken from Fig. 11 of [79].

III. SCALAR WAVES

A. Curvature coupling and first-order system

We study the massless curvature-coupled equation

$$(\square_g - \xi R[g]) \Phi = 0, \quad \xi \in \left\{0, \frac{1}{6}\right\}, \quad (25)$$

where g denotes the physical SdS or Schwarzschild metric. Minimal coupling has $\xi = 0$; conformal coupling in four spacetime dimensions has $\xi = 1/6$. Write

$$\Phi(t, r, \theta, \varphi) = \frac{1}{r} \sum_{\ell m} u_{\ell m}(t, r) Y_{\ell m}(\theta, \varphi). \quad (26)$$

Each mode satisfies

$$(-\partial_t^2 + \partial_{r_*}^2 - V_{\ell, \xi}) u = 0, \quad (27)$$

$$V_{\ell, \xi} = f \left[\frac{\ell(\ell+1)}{r^2} + \frac{f'(r)}{r} + \xi R[g] \right]. \quad (28)$$

For uniform SdS, $R[g] = 12/L^2$ and therefore

$$V_{\ell, \xi} = f \left[\frac{\ell(\ell+1)}{r^2} + \frac{2M}{r^3} + \frac{2(6\xi - 1)}{L^2} \right]. \quad (29)$$

There is a $-2/L^2$ term at $\xi = 0$ which cancels at $\xi = 1/6$. In Schwarzschild, $R[g] = 0$, so both couplings give the same equation and the same reference waveform.

To see the role of curvature coupling, we write the conformal transformation for the scalar wave equation. Let $\bar{g} = \Omega_L^2 g$ and $\phi = \Omega_L^{-1} \Phi$. For the sourced wave equation, conformal covariance gives

$$\left(\square_{\bar{g}} - \frac{R[\bar{g}]}{6} \right) \phi + \left(\frac{1}{6} - \xi \right) \frac{R[g]}{\Omega_L^2} \phi = \Omega_L^{-3} S. \quad (30)$$

The curvature-coupling coefficient vanishes at $\xi = 1/6$. Because $\Omega_L = r_b/r$, the evolved field is proportional to its conformal counterpart, $\phi_{\ell m} = (u_{\ell m}/r_b) Y_{\ell m}$.

The scalar equations contain no tensorial conformal Einstein source terms, and therefore do not fully test the regularization idea. Nevertheless, minimal scalar coupling has a lower-order term with the same Λ/Ω_L^2 scaling as (24), whereas conformal coupling cancels it. The two cases provide, respectively, an algebraic stress test and a conformally regular radiative analogue. They test whether matched scalar observables recover their Schwarzschild counterparts in a controlled large- L limit.

Next, we discuss the first-order reduction used in the numerical implementation. We suppress the mode labels below. For the time evolution, define

$$p := \frac{d\rho}{dr_*} = fG, \quad A := \frac{p}{1 - B^2}, \quad (31)$$

and the potential coefficient

$$P_{\ell, \xi} := \frac{V_{\ell, \xi}}{p}. \quad (32)$$

We write the equation in first-order symmetric hyperbolic form using auxiliary variables defined as

$$\psi := \partial_\rho u, \quad \pi := \frac{1 - B^2}{p} \partial_\tau u - B\psi. \quad (33)$$

The evolution system becomes

$$\partial_\tau u = A(B\psi + \pi), \quad (34)$$

$$\partial_\tau \psi = \partial_\rho [A(B\psi + \pi)], \quad (35)$$

$$\partial_\tau \pi = \partial_\rho [A(\psi + B\pi)] - P_{\ell, \xi} u. \quad (36)$$

To check the accuracy of the evolution, we monitor the constraint

$$C := \psi - \partial_\rho u. \quad (37)$$

The radial characteristic speeds are

$$c_- = -A(1 + B) = -\frac{p}{1 - B}, \quad c_+ = A(1 - B) = \frac{p}{1 + B}. \quad (38)$$

The boundary speeds are $(c_+, c_-) = (0, c_- < 0)$ at $\rho = 0$ and $(c_+, c_-) = (c_+ > 0, 0)$ at $\rho = 1$. The characteristic flow determines all boundary values. For a simple crossing with $B'(r_h) \neq 0$, analytic cancellation gives

$$A_h = \frac{f'(r_h)G(r_h)}{-2s_h B'(r_h)}, \quad s_b = +1, \quad s_c = -1. \quad (39)$$

We cancel every height-function pole algebraically and assign (39) at each horizon.

The evolution domain ends at the cosmological horizon, $\mathcal{H}_c^+$. No data from beyond the horizon is needed. This is similar to the excision procedure in numerical relativity, which motivates the title of Misner's paper [3]. We use the analytic continuation of the height function only as a geometric diagnostic of the foliation family and of the asymptotic end it approaches as $L/M \rightarrow \infty$. The finite- L evolutions test the regular horizon coefficients and extract the outer waveform on $\mathcal{H}_c^+$.

For the minimal gauge, the limiting Schwarzschild coefficients are

$$B_0 = -1 + 2(1 - \rho_0)^2, \quad A_0 = \frac{1}{8M(2 - \rho_0)}, \quad (40)$$

$$P_{\text{Schw}} = \frac{\ell(\ell + 1) + (1 - \rho_0)}{2M}. \quad (41)$$

The Schwarzschild code implements these expressions separately from the finite- L geometry. For uniform SdS at fixed interior ρ , both finite- L potentials approach this result. At the cosmological horizon, however, the limits are

$$\begin{aligned} \lim_{L \rightarrow \infty} P_{\ell,0}(1) &= \frac{\ell(\ell + 1) - 2}{2M}, \\ \lim_{L \rightarrow \infty} P_{\ell,1/6}(1) &= \frac{\ell(\ell + 1)}{2M} = P_{\text{Schw}}(1). \end{aligned} \quad (42)$$

B. Matched data, sources, clocks, and observables

To isolate the effect of artificial cosmology, we compare waveforms using matched data and a common time normalization. We match the physical data and source geometrically so that changes in their coordinate representation do not introduce a different physical solution. For the data choices below, we use the peak-normalized bump

$$b(x) = \begin{cases} \exp[1 - 1/(1 - x^2)], & |x| < 1, \\ 0, & |x| \geq 1. \end{cases} \quad (43)$$

a. Fixed-window data. We first test prompt and ringdown waveform recovery on retarded-time intervals held fixed as L varies, separating finite-time agreement from the eventual cosmological decay. The finite-time comparison uses the same areal-radius profile on every background,

$$u(0, r) = b\left(\frac{r - 4M}{1.5M}\right), \quad \psi = D_\rho u, \quad \pi = -B\psi, \quad (44)$$

with support $2.5M < r < 5.5M$. Here D_ρ is the spectral differentiation operator; $\partial_\tau u = \partial_t u = 0$ initially, so the data are stationary with respect to Killing time.

b. Tail data. To probe weak late-time decay separately, we use an initially dynamical pulse with zero displacement and nonzero velocity. The Schwarzschild solution has the Price-law decay. We prescribe

$$\begin{aligned} u = \psi = 0, \quad G_v(r) &= b\left(\frac{r - 6M}{3M}\right), \\ \partial_\tau u &= G_v, \quad \pi = G_v/A, \end{aligned} \quad (45)$$

with support $3M < r < 9M$. In Eqs. (44) and (45), the common areal profiles and prescribed Killing-time derivatives define data on L -dependent slices approaching the chosen Schwarzschild slice.

c. Retarded time. Matched waveforms require a common retarded-time origin. We set $r_0 = 4M$ and normalize

$$h_L(r_0) = 0, \quad r_{*,L}(r_0) = 0. \quad (46)$$

For a bridge with this normalization, the offset between τ and the outgoing null coordinate at the cosmological horizon is

$$q_B(L; r_0) := \lim_{r \rightarrow r_c} (h_L + r_{*,L}) = \int_{r_0}^{r_c} \frac{1 + B}{f} dr. \quad (47)$$

The integrand is regular at r_c , where $U = \tau - q_B = t - r_{*,L}$. For the minimal gauge, cancellation of the horizon logarithms and the identity $(2\kappa_b)^{-1} - (2\kappa_c)^{-1} + (2\kappa_u)^{-1} = 0$ give

$$q_L = \frac{1}{\kappa_b} \ln \frac{r_0(r_c - r_b)}{r_c(r_0 - r_b)} \rightarrow q_0. \quad (48)$$

The finite Schwarzschild limit is

$$q_0 = 4M \ln \frac{r_0}{r_0 - 2M}, \quad (49)$$

which reduces to $q_0 = 4M \ln 2$ for the reference radius $r_0 = 4M$. Waveforms are compared at the geometric retarded time

$$U = \tau - q_L. \quad (50)$$

At $\mathcal{H}_c^+$, U is the outgoing retarded-time coordinate. The same normalization reaches Schwarzschild $\mathcal{I}^+$ in the limit. At finite radius, U is a stationary clock normalized at the outer boundary. Its relation to the local outgoing coordinate is

$$U - (t - r_{*,L}) = h_L(r) + r_{*,L}(r) - q_L. \quad (51)$$

An additive clock offset preserves the asymptotic decay exponent but changes the finite-time logarithmic slope used to estimate it. We therefore use the common geometric time origin in (50) when comparing local power indices and transition times.

d. Localized source. To test artificial cosmology beyond single-mode radial evolution, we consider a spatially localized scalar response with nontrivial angular structure. This tests Schwarzschild recovery for multidimensional wave propagation, including focusing and caustic echoes [38, 91, 92]. We exploit the spherical background to evolve independent radial modes and reconstruct the three-dimensional field. We use a normalized source localized in radius, time, and angle as a smooth approximation to a Dirac delta function. The resulting field is the retarded Green function integrated against this source profile. We compare the full localized response across backgrounds. With zero initial data,

$$(\square_g - \xi R[g])\Phi = S, \quad \Phi|_{\tau=0} = \partial_\tau \Phi|_{\tau=0} = 0. \quad (52)$$

The same physical source S is used on every background; its conformal representation is $\Omega_L^{-3}S$. Let γ be the angle from the source direction (θ_s, φ_s) and set

$$S = \mathcal{A}T(t)R(r)\mathcal{P}_\kappa(\gamma), \quad (53)$$

$$T(t) = \frac{b[(t-t_s)/\sigma_t]}{\sigma_t\beta_0}, \quad R(r) = \frac{b[(r-r_s)/\sigma_r]}{\beta_2},$$

$$\mathcal{P}_\kappa(\gamma) = \frac{\kappa e^{\kappa(\cos\gamma-1)}}{2\pi(1-e^{-2\kappa})}. \quad (54)$$

Here $\beta_0 = \int_{-1}^1 b(x)dx$ and $\beta_2 = \int b[(r-r_s)/\sigma_r]r^2dr$, so each factor has unit integral and $\int \sqrt{-g}S d^4x = \mathcal{A}$. The parameter choices are given in Sec. IV B.

The real-spherical-harmonic coefficients are

$$a_{\ell m} = \frac{i_\ell(\kappa)}{i_0(\kappa)} Y_{\ell m}(\theta_s, \varphi_s), \quad (55)$$

where i_ℓ is a modified spherical Bessel function. The m dependence factors out, so one response per retained ℓ solves

$$(-\partial_t^2 + \partial_{r_*}^2 - V_{\ell,\xi})v_\ell = fr\mathcal{A}T(t)R(r). \quad (56)$$

In the first-order system the source is evaluated explicitly at the Killing time $t = \tau - h_L(r)$, ensuring the same emitter on every background. With $u_{\ell m} = a_{\ell m}v_\ell$, the reconstructed waveform is

$$u(U, \theta, \varphi) = \sum_{\ell m} a_{\ell m} v_\ell(U) Y_{\ell m}(\theta, \varphi), \quad (57)$$

at the extraction radius, using the common retarded time (50).

At the outer boundary, the $\xi = 0$ observable is the reduced field $W_L(U) \equiv W_{L,0}(U) := u_2(U, \rho = 1)$ on $\mathcal{H}_c^+$; its Schwarzschild counterpart $W_{\text{Schw}}(U) := u_2(U, \rho_0 = 1)$ is evaluated at $\mathcal{J}^+$. Thus W is the relevant spherical-harmonic coefficient of the radiation-scaled field $r\Phi$. For a common time interval $\mathcal{J}$, the direct fixed-mode waveform error is

$$E_2(L; \mathcal{J}) = \frac{\|W_L - W_{\text{Schw}}\|_{L^2(\mathcal{J})}}{\|W_{\text{Schw}}\|_{L^2(\mathcal{J})}}, \quad (58)$$

where W_{Schw} is computed with the separate Schwarzschild implementation. For the localized source, W denotes the outer-boundary fields $\{u_{\ell m}\}$ and the primary comparison uses the sphere-integrated Parseval norm

$$\|W\|_{\mathcal{J}, S^2}^2 = \int_{\mathcal{J}} dU \sum_{\ell m} |u_{\ell m}(U)|^2. \quad (59)$$

The corresponding relative error is

$$E_{2,S^2}(L; \mathcal{J}) = \frac{\|W_L - W_{\text{Schw}}\|_{\mathcal{J}, S^2}}{\|W_{\text{Schw}}\|_{\mathcal{J}, S^2}}. \quad (60)$$

Besides comparing individual finite- L waveforms with Schwarzschild, we test whether combining several cosmological lengths improves the recovery. For minimal coupling, we model the large- L dependence at each common retarded time U as

$$W_{L,0} = W_{\text{Schw}} + c_{1,0} \frac{M}{L} + c_{2,0} \left(\frac{M}{L}\right)^2 + \mathcal{O}((M/L)^3). \quad (61)$$

At each U , the coefficients $c_{1,0}$ and $c_{2,0}$ are independent of L . Waveforms at cosmological lengths L , $2L$, and $4L$ then give the extrapolated estimate

$$W_\infty^{(L)} = \frac{W_L - 6W_{2L} + 8W_{4L}}{3}, \quad (62)$$

which cancels the terms proportional to M/L and $(M/L)^2$ without using the Schwarzschild waveform as input. Repeating the construction with $(2L, 4L, 8L)$ gives a second, overlapping extrapolant. We compare the two estimates with each other and with the separately evolved Schwarzschild control to test the extrapolation. A conformally coupled sequence requires a separate test of its finite- L expansion.

The fixed-mode comparison uses cumulative intervals that begin at $U = 0$ and end at $U/M = 40, 80$, and 160 , together with the disjoint intervals $40 \leq U/M \leq 80$ and $80 \leq U/M \leq 160$. The localized-source comparison uses the common simulated interval, from the first nonnegative output time through the last time reached by every evolution. The uniform waveform and localized-source sequences use linear interpolation onto a common U grid; the exterior ringdown and matched-tail comparisons use cubic interpolation.

C. Numerical methods and error control

We tested artificial cosmology using different numerical methods. The uniform waveform sequence and the large- L tail benchmark use Dedalus 3 [93] with a Chebyshev- T basis and RK222. The localized source uses eighth-order finite differences on a uniform radial grid and classical RK4, with source evaluation at each stage. The exterior ringdown and matched-tail comparisons use continuous Legendre-Gauss-Lobatto spectral elements [94]

and third-order Radau IIA [95], evolving u and $\partial_\tau u$ directly. All calculations evolve the characteristic boundaries without prescribed boundary data and use the geometric clock normalization defined above.

Each simulation sequence is tested at three resolutions. Uniform and localized-source comparisons use matched Schwarzschild controls. We distinguish discretization error from sensitivity to the estimator, comparison window, and sampling interval, and from physical source-width dependence. Appendix A gives the resolution sequences and independent checks; Appendix B gives the tail-specific criteria and convergence results.

IV. RECOVERING SCHWARZSCHILD SOLUTIONS WITH ARTIFICIAL COSMOLOGY

In this section, we quantify the waveform error when a cosmological horizon replaces null infinity at finite L . The same comparisons also test the isolated-system approximation in a cosmological spacetime, motivated by the small positive cosmological constant. Increasing L/M , we study how scalar waveforms approach their asymptotically flat limits over finite observation intervals. We also check different aspects of the regulator, including the convergence of extrapolated waveforms and the recovery of a localized multimode response. The results show that the Misner regulator is effective for both minimal and conformal coupling, and that the finite- L error can be reduced to a few percent with $L/M \sim 10^2$.

A. Direct and extrapolated fixed-mode recovery

Figure 2 shows how the boundary waveforms approach the Schwarzschild solution as L/M increases from 20 to 640. The prompt-pulse and ringdown structure is maintained throughout the sequence. In the large- L regime, the dominant residual amplitude falls by about a factor of two with each doubling of L .

Table I quantifies this convergence and gives the benchmark for the regulator: the direct waveform error E_2 against the separately evolved Schwarzschild solution, measured over the same interval $0 \leq U/M \leq 80$ for every L . The error decreases from 63.96% at $L/M = 20$ to 1.34% at $L/M = 640$. The numerical medium-to-fine changes are much smaller, showing that the finite- L difference significantly dominates over discretization error.

For these data and this norm, the first tested lengths to meet 5% and 2% tolerances are $L/M = 320$ and 640, respectively. The approximate M/L falloff is encoded in the extrapolation model (61). At fixed outgoing retarded time $\hat{U} = t - r_{*,0}$, a smooth Schwarzschild field can have the expansion $u = F(\hat{U}) + G(\hat{U})/r + \mathcal{O}(r^{-2})$, with $2dG/d\hat{U} = \ell(\ell+1)F$. Since $r_c \sim L$, such corrections show how an $\mathcal{O}(M/L)$ boundary discrepancy can coexist with an $\mathcal{O}(L^{-2})$ local metric deformation.




FIG. 2. Minimally coupled ($\xi = 0$), pure- $\ell = 2$ sequence at the outer boundary. Upper panel: waveforms for $L/M = 20$ to 640 against the separately evolved Schwarzschild reference. Lower panel: the residual $W_L - W_{\text{Schw}}$ on the same retarded time $U = \tau - q_L$, with no other fitting or rescaling. The residual amplitude falls with L . Compare with Table I.

TABLE I. Direct waveform error for the minimally coupled pure- $\ell = 2$ sequence on $0 \leq U/M \leq 80$. E_2 is the fine-grid discrepancy from Schwarzschild [Eq. (58)]; δ_{mf} is the numerical medium-to-fine change of the paired residual [Eq. (A1)]. Both columns are percentages of the Schwarzschild norm.

L/M	E_2 (%)	δ_{mf} (%)
20	63.96	0.033
40	26.08	0.094
80	11.75	0.069
160	5.57	0.041
320	2.72	0.064
640	1.34	0.024

To test sensitivity to higher-order finite- L corrections, we use extrapolation and combine triples (80, 160, 320) and (160, 320, 640) in units of L/M . The largest fine-grid residual among $W_\infty^{(80)}$, $W_\infty^{(160)}$, and the separately evolved Schwarzschild waveform is 0.0313%. The largest numerical medium-to-fine change among these residuals is 0.409%, and the largest propagated Richardson estimate is 0.119%. We can conclude that the extrapolated waveforms agree with the separately evolved Schwarzschild waveform at 1%. Appendix A gives the numerical refinement and extrapolation checks.




FIG. 3. Minimally coupled localized source on the common simulated interval. (a) Direct sphere-integrated errors E_{2,S^2} in percent; the dashed line marks 1%. (b) Instantaneous absolute $L^2(S^2)$ norms of the Schwarzschild field, the finite- L residuals against Schwarzschild, and the difference between the two nested extrapolants. The norms include all retained modes and integrate over the entire observation sphere.

B. Localized multimode response

To extend the $\xi = 0$ benchmark beyond a single spherical harmonic, we solve for the localized source family described in Section III B. We use $r_s = 6M$, $\sigma_r = 0.75M$, $t_s = 30M$, $\sigma_t = 2M$, $\kappa = 64$, and $(\theta_s, \varphi_s) = (\pi/2, 0)$. These choices give radial support $5.25M < r < 6.75M$, temporal support $28M < t < 32M$, and angular scale $\kappa^{-1/2} = 0.125$ radians. Modes with $|a_{\ell m}|$ below 10^{-13} of the maximum are omitted. We extract waveforms at $r = 8M$, $r = 12M$, and the outer boundary; the comparison uses the outer-boundary field on the common interval $0 \lesssim U/M \leq 57.2$, before the late-time signal.

In the sphere-integrated norm (59), the direct differences are $E_{2,S^2}(320) = 5.27\%$ and $E_{2,S^2}(640) = 2.63\%$. The $W_\infty^{(80)}$ and $W_\infty^{(160)}$ extrapolants differ by 0.0517%; their Schwarzschild differences are 0.0584% and 0.00672%, respectively. Figure 3 shows how the error decreases with L and how the residuals vary over the signal.

C. Large- L Price-law benchmark

Tails provide a stricter test than the prompt waveform. To test Schwarzschild decay-rate at the asymptotic boundary with artificial cosmology, we evolve matched $\ell = 1$, $\xi = 0$ data at $L/M = 3072$, extracting the SdS waveform at $\mathcal{H}_c^+$, and we compare to an independently evolved Schwarzschild reference at $\mathcal{I}^+$.

For each extracted waveform $W(U)$, we define the centered root-mean-square envelope

$$A(U) = \left[\frac{1}{\Delta U} \int_{U-\Delta U/2}^{U+\Delta U/2} |W(s)|^2 ds \right]^{1/2}, \quad (63)$$

with $\Delta U = 10M$, evaluated as an average of squared waveform samples. We estimate the local power index $p_{\text{eff}} = -d \ln A / d \ln U$ by fitting $\ln A$ against $\ln U$ over a centered $40M$ window. For a slowly varying power-law tail, $A \simeq |W|$, so this index approximates the decay exponent.

The SdS local power index remains within 5% of $p = 3$ on $232 \lesssim U/M \lesssim 382$. Spatial and timestep checks give a spread of about $2M$ (see Appendix B 1 for details). These checks indicate that an intermediate interval is compatible with the Schwarzschild Price-law rate.

Figure 4 shows the Price-law interval within the longer waveform. The common evolution extends to $U/M \simeq 1984$, or $\kappa_c U = 0.645$. In the experiments that demonstrate the polynomial decay, the cosmological exponential regime is not reached before the numerical floor. This is reasonable and supports the well-known idea that the isolated system approximation is valid for long times before cosmological effects become relevant. Appendix B 1 gives further details.

D. Cosmological decay

At smaller L , the exponential cosmological decay becomes accessible within the resolved evolution. A separate minimally coupled $\ell = 1$ study at $r = 8M$ measures both departure from the Schwarzschild decay rate and entry into the cosmological regime. For $L/M = 80$ and 160, the median departure times are $U = 115.9M$ and $133.5M$, respectively, whereas cosmological decay begins at $\kappa_c U = 2.83$ and 2.94. Doubling L therefore changes the departure time only modestly in black-hole units, while entry occurs at nearly the same time in cosmological units. These two cases provide qualitative evidence for distinct timescales.

Both departures precede a settled Schwarzschild Price tail: at the same radius, the reference decay index does not remain within 5% of its $\ell = 1$ Price-law value until $U = 220.3M$. These runs resolve a transition from a Schwarzschild-like transient to cosmological decay. The large- L boundary test above resolves an intermediate Price-law interval; its later cosmological regime remains




FIG. 4. Intermediate, polynomial tail decay rate in minimally coupled $\ell = 1$ SdS at $L/M = 3072$. Upper panel: the local power indices at $\mathcal{H}_c^+$ (solid orange) and Schwarzschild $\mathcal{J}^+$ (dashed black). The polynomial rate is compatible within the range $p = 3 \pm 5\%$ in the interval $232 \lesssim U/M \lesssim 382$. The decay slightly is stronger in SdS. Lower panel: the centered $10M$ root-mean-square envelopes defined in (63). We see that the SdS waveform is slightly smaller than the Schwarzschild one, but the difference is small and consistent with the expected cosmological expansion.

numerically unresolved. Appendix B3 gives the transition criteria and their sensitivity to the decay-rate estimator.

V. EXTERIOR-SUPPORTED ARTIFICIAL COSMOLOGY

Uniform SdS alters the metric throughout the static patch, including the strong-field region. But Misner's proposal primarily concerns the regularization of the asymptotic domain. Therefore, it makes sense to confine the cosmological source to an exterior region while retaining the same asymptotic value

$$\Lambda = \frac{3}{L^2}. \quad (64)$$

We would like the cosmological source to vanish in a domain containing the event horizon and the photon sphere. Then, in an interior domain, the metric, boost, and scalar

potential are exactly Schwarzschild. However, the outer layer, cosmological boundary, and clock still affect the waveform. We perform comparisons with uniform SdS at fixed L to test whether this construction reduces the total waveform error. We first construct the source and its metric based on an idea by Dymnikova [83, 84] relying on the static spherical ansatz. We then specify the foliation and compare ringdown and tail signals.

A. Conserved cosmological tensor and exact metric

Restricting the cosmological term to an exterior region is not immediately trivial. The term Λ is called cosmological *constant* for a geometric reason. Simply replacing Λ by a radial profile $\lambda(r)$ in the vacuum equations violates the contracted Bianchi identity unless it is constant:

$$G_{ab} + \lambda g_{ab} = 0 \implies 0 = \nabla^a G_{ab} = -\nabla_b \lambda. \quad (65)$$

But we can replace the cosmological term by a conserved *tensor*,

$$G^a_b + \Lambda^a_b = 0, \quad \nabla_a \Lambda^a_b = 0. \quad (66)$$

In static spherical symmetry, radial variation is compatible with conservation if the source includes suitable anisotropic stresses. Following Dymnikova's anisotropic-vacuum framework [83, 84], we choose equal temporal and radial eigenvalues $\Lambda_{\parallel}$ and angular eigenvalues $\Lambda_{\perp}$:

$$\Lambda^a_b = \text{diag}(\Lambda_{\parallel}, \Lambda_{\parallel}, \Lambda_{\perp}, \Lambda_{\perp}). \quad (67)$$

For an arbitrary smooth profile $\lambda(r)$, set

$$\Lambda_{\parallel}(r) = \lambda + \frac{r}{3} \lambda', \quad (68)$$

$$\Lambda_{\perp}(r) = \lambda + r \lambda' + \frac{r^2}{6} \lambda''. \quad (69)$$

Equality of the temporal and radial eigenvalues cancels the metric-derivative terms in the radial conservation equation, leaving

$$\nabla_a \Lambda^a_r = \Lambda'_{\parallel} + \frac{2}{r} (\Lambda_{\parallel} - \Lambda_{\perp}) = 0. \quad (70)$$

The contributions proportional to λ' and λ'' cancel exactly for any smooth profile; symmetry makes the other components vanish. Where λ is constant, the tensor reduces to $\lambda \delta^a_b$. Where it varies, the derivative terms supply the additional stresses required by conservation. Thus $\lambda(r)$ parametrizes a conserved anisotropic source, allowing us to circumvent the obstacle against a variable vacuum term $\lambda(r)g_{ab}$.

We now choose the profile to give a Schwarzschild interior and an SdS exterior. Set $\lambda(r) = \Lambda \chi_L(r)$, with $\chi_L = 0$ below R_0 and $\chi_L = 1$ above R_1 . To join these regions smoothly, let $\eta(x) = e^{-1/x}$ for $x > 0$ and zero

otherwise, and define $S(x) = \eta(x)/[\eta(x) + \eta(1-x)]$. For $2M < R_0 < R_1 < r_c$, the switch is

$$\chi_L(r) = \begin{cases} 0, & r \leq R_0, \\ S(\zeta_L(r)), & R_0 < r < R_1, \\ 1, & r \geq R_1, \end{cases} \quad (71)$$

where ζ_L is a smooth monotone map with $\zeta_L(R_0) = 0$ and $\zeta_L(R_1) = 1$. We choose the transition radii so that both the transition layer and the outer exact-SdS region retain nonzero widths in the compactified radial coordinate as $L \rightarrow \infty$. Appendix C gives the explicit profile.

The Einstein equations with this source admit the metric

$$ds^2 = -f_\chi(r)dt^2 + \frac{dr^2}{f_\chi(r)} + r^2 d\omega^2, \quad (72)$$

with

$$\begin{aligned} f_\chi(r) &= 1 - \frac{2M}{r} - \frac{r^2}{3}\lambda(r) \\ &= 1 - \frac{2M}{r} - \frac{r^2}{L^2}\chi_L(r). \end{aligned} \quad (73)$$

The integration constant M is the same mass parameter in the interior and exterior. For $M > 0$, the metric is exactly Schwarzschild below R_0 and SdS above R_1 . Between them, the derivative terms in the source support a smooth, nonvacuum transition layer.

The enclosed Misner–Sharp mass is

$$f_\chi = 1 - \frac{2m_\chi}{r}, \quad m_\chi(r) = M + \frac{r^3\lambda(r)}{6}. \quad (74)$$

It includes the cosmological source as well as the central mass parameter M , which remains unchanged.

Interpreting the source as a stress tensor gives $T^a_b = -(8\pi)^{-1}\Lambda^a_b = \text{diag}(-\varepsilon, p_r, p_\perp, p_\perp)$, with $\varepsilon = \Lambda_\parallel/(8\pi)$, $p_r = -\varepsilon$, and $p_\perp = -\Lambda_\perp/(8\pi)$. The density $\varepsilon = (\lambda + r\lambda'/3)/(8\pi)$ obeys $m'_\chi = 4\pi r^2\varepsilon$. The energy-condition combinations are

$$\varepsilon + p_r = 0, \quad \varepsilon + p_\perp = -\frac{r}{16\pi}\Lambda'_\parallel. \quad (75)$$

The radial null energy condition is satisfied. Since $\int_{R_0}^{R_1} \Lambda'_\parallel dr = \Lambda > 0$, the tangential condition is violated somewhere in the layer. The regulator therefore cannot be interpreted as ordinary matter satisfying the null energy condition.

Extending the construction to nonlinear Einstein evolution would require an evolution law for the source, beyond background conservation. A metric perturbation changes the connection, so source perturbations must also satisfy

$$\delta(\nabla_a \Lambda^a_b) = 0. \quad (76)$$

The background tensor therefore cannot, in general, be held fixed during nonlinear evolution. Such an extension needs a covariant conserved regulator.

The Ricci scalar quantifies the curvature introduced by the source:

$$R_\chi = 4\lambda + \frac{8}{3}r\lambda' + \frac{1}{3}r^2\lambda''. \quad (77)$$

It vanishes in the Schwarzschild interior and equals 4Λ in the SdS exterior. The Misner–Sharp mass accumulated across the smooth layer is

$$\frac{\Delta m_{\text{tr}}}{M} := \frac{m_\chi(R_1) - m_\chi(R_0)}{M} = \frac{1}{2} \frac{(R_1/M)^3}{(L/M)^2}. \quad (78)$$

Appendix C gives the large- L scaling of the curvature and accumulated mass. At $L/M = 640$, the layer ends at $R_1 \simeq 440.4M$ and contains $\Delta m_{\text{tr}} = 104.3M$. Uniform SdS has $m_{\text{SdS}}(r) = M + r^3/(2L^2)$, so both constructions have exactly the same enclosed mass at R_1 and beyond. The derivative contribution to the source density compensates for removing the cosmological source from the interior, while retaining the same exterior mass parameter. The outer geometry can differ significantly from Schwarzschild: $|f_\chi(R_1) - f_{\text{Schw}}(R_1)| = (R_1/L)^2 \simeq 0.474$ in this example. This is compatible with the fixed-radius Schwarzschild limit; at finite L , neither construction need approximate Schwarzschild closely at radii comparable to L . The effect on extracted radiation must instead be assessed through waveform comparisons, as in the following subsections.

B. Bridge foliation, compactification, and clock

The exterior metric needs a bridge that crosses both horizons and recovers the Schwarzschild foliation as $L \rightarrow \infty$. For $M > 0$, we choose

$$B_\chi(r) = -1 + \frac{8M^2}{r^2} - \frac{8M^2}{r_c^2}\chi_L(r). \quad (79)$$

It has $B_\chi(2M) = 1$, $B_\chi(r_c) = -1$, and $-1 < B_\chi < 1$ between the horizons; beyond r_c , $B_\chi < -1$. The height function satisfies $h'_\chi = B_\chi/f_\chi$; the horizon crossings are regular, as verified in Appendix C. Since $0 \leq \chi_L \leq 1$ and $r_c/L \rightarrow 1$, the cosmological correction to the boost vanishes as $\mathcal{O}(M^2/L^2)$. Hence, at fixed r ,

$$B_\chi(r) = -1 + \frac{8M^2}{r^2} \quad (L \rightarrow \infty), \quad (80)$$

which is the minimal-gauge Schwarzschild boost. Likewise $r^2\chi_L/L^2 \rightarrow 0$, so the metric and bridge recover Schwarzschild at fixed radius even if χ_L has a nonzero limiting profile.

The compact radial coordinate maps the two horizons to fixed boundary points:

$$D_L = 1 - \frac{2M}{r_c}, \quad \rho_\chi(r) = \frac{1 - 2M/r}{D_L}. \quad (81)$$

TABLE II. Ringdown waveform errors relative to Schwarzschild for uniform and exterior-supported SdS, using minimally coupled ($\xi = 0$), $\ell = 2$ data on $15 \leq U/M \leq 45$. The relative error reduction is $100(1 - E_\chi/E_{\text{SdS}})\%$.

L/M	uniform SdS error (%)	exterior SdS error (%)	reduction (%)
320	3.18	2.77	12.8
640	1.55	1.42	8.6

It places $2M$ and r_c at $\rho_\chi = 0$ and 1 and goes to $1 - 2M/r$ at fixed radius. The conformal factor is

$$\Omega_{\chi,L} = \frac{2M}{r}, \quad \Omega_{\chi,L}(1) = \frac{2M}{r_c} > 0. \quad (82)$$

Its value at the outer boundary is positive at finite L and vanishes in the Schwarzschild limit. The coefficients multiplying spatial derivatives at the outer boundary approach their Schwarzschild values as $L \rightarrow \infty$ (Appendix C).

To compare waveforms on the same geometric clock, we keep the normalization $r_0 = 4M$ and define

$$q_\chi = \int_{r_0}^{r_c} \frac{1 + B_\chi}{f_\chi} dr, \quad U = \tau - q_\chi. \quad (83)$$

The integrand has a finite horizon limit. Comparing its large-radius behavior with the Schwarzschild expression gives

$$q_\chi = q_0 - \frac{8M^2}{r_c} + o\left(\frac{M^2}{r_c}\right) \Rightarrow q_0 = 4M \ln \frac{r_0}{r_0 - 2M}. \quad (84)$$

Therefore, the exterior family approaches the same Schwarzschild clock as the uniform construction.

C. Ringdown comparison

We test whether preserving the near-zone geometry improves the waveform at the outer boundary. We compare minimally coupled ($\xi = 0$), compact $\ell = 2$ data at $L/M = 320, 640$, using matched initial data and clocks for Schwarzschild, uniform SdS, and exterior-supported SdS. For either cosmological background, the relative waveform error against the same separately evolved Schwarzschild reference is

$$E_X(L; \mathcal{J}) = \frac{\|W_X - W_{\text{Schw}}\|_{\mathcal{J}}}{\|W_{\text{Schw}}\|_{\mathcal{J}}}. \quad (85)$$

Here $X \in \{\text{SdS}, \chi\}$. We compare waveforms at the outer boundary over the ringdown interval $15 \leq U/M \leq 45$.

Restricting the cosmological source to the exterior reduces the ringdown waveform error by 12.8% at $L/M = 320$ and 8.6% at $L/M = 640$ (Table II).

This fixed-mode test quantifies ringdown waveform agreement for the specified initial data. Determining the




FIG. 5. Matched dipole tails at $L/M = 640$, compared at $\mathcal{H}_c^+$ with Schwarzschild at $\mathcal{J}^+$. (a) Centered 10M RMS envelopes. (b) Local power indices from 40M fits; the gray band marks $p = 3 \pm 10\%$ and the gold shading the uniform-conformal Price-compatible interval.

global quasinormal spectrum requires a separate spectral analysis. Distant perturbations can strongly alter that spectrum [96] while leaving early ringdown nearly unchanged, as demonstrated for a black hole surrounded by a dynamical thin shell [97].

D. Exterior-supported tail comparison

Tail recovery tests the construction beyond the ringdown window. Matched $\ell = 1$ evolutions at $L/M = 640$ through $U/M = 1000$ compare uniform and exterior-supported SdS at $\mathcal{H}_c^+$ with Schwarzschild at $\mathcal{J}^+$, for both $\xi = 0$ and $1/6$. Here we require the local decay index to remain within 10% of $p = 3$ for at least 40M. Uniform conformal SdS meets this criterion; uniform minimal coupling does not. At fixed L , the field equation therefore matters for the intermediate polynomial decay rates.

At $L/M = 640$, the exterior-supported construction fails to recover the Schwarzschild tail decay rate for either coupling (Fig. 5). Both evolutions develop slowly decaying components and secondary lobes that persist under refinement, and neither meets the 40M criterion. This behavior is consistent with the sensitivity of tails to scattering in the outer geometry, which remains modified by the regulator. Appendix B 2 gives the intervals and convergence evidence.

VI. CONCLUSIONS

We tested Misner’s artificial cosmology as a regulator at the conformal boundary using scalar radiation from a compact source in a black-hole spacetime. The regulator introduces a small positive cosmological constant and a smooth foliation that reaches the cosmological horizon. The resulting SdS spacetime is asymptotically de Sitter, with a conformal boundary at finite radius. An important aspect of the construction is that the foliation must resolve the conformal boundary of the cosmological spacetime for it to have a numerically useful limit to the asymptotically flat case (see Fig. 1 and [78, 79]). Then, the limiting slices meet distinct cuts of future null infinity, and their conformal spatial metric remains smooth and nondegenerate. In this paper, we have used a minimal-gauge foliation that recovers the Schwarzschild slices in the limit of vanishing cosmological constant [44, 46, 78, 85].

Beyond testing Misner’s numerical regulator on wave propagation, these calculations provide a model for understanding the effects of a small positive cosmological constant on outgoing waveforms. The single-mode and localized-source comparisons show how Schwarzschild recovery improves with increasing L and provide benchmarks for choosing a cosmological scale. At sufficiently large L , we demonstrated that uniform SdS provides accurate ringdown waveforms and even gives the polynomial Price-law decay rate in intermediate intervals. The coupling comparison shows that tail recovery depends on the asymptotic field equation. This construction can also be used to numerically test the asymptotically flat approximation for isolated systems in a Universe with a small positive cosmological constant [72, 74].

We extend Misner’s artificial cosmology by restricting the cosmological modification to an exterior layer, based on Dymnikova’s conserved anisotropic source [83, 84]. This approach preserves the strong-field geometry and improves ringdown agreement relative to uniform SdS at fixed L . However, it does not improve tail recovery. The transition layer changes the scattering potential, and tails are sensitive to the outer geometry and source history [98, 99]. Extending this construction to nonlinear Einstein evolution is more difficult. Source conservation must hold as the metric evolves, requiring a covariant evolution law beyond the static spherical ansatz. The stresses in the transition layer also violate the null energy condition.

A natural next step is to compute gravitational waveforms from particle sources inspiraling into the central black hole. This would test cosmological effects on long, astrophysically relevant signals. A separate nonlinear Einstein study is needed to determine whether Misner’s artificial cosmology is a practical proposal for the treatment of the asymptotic boundary.

ACKNOWLEDGMENTS

A.Z. thanks Ted Jacobson, Rodrigo Panosso Macedo, and Alex Vañó-Viñuales for discussions. This work was inspired by Charles Misner’s investigations of artificial cosmology and by the memorial symposium held in his honor at the University of Maryland in 2023. This material is supported by NSF Grant No. 2309084. A.Z. used OpenAI Codex and G.A.K. used Anthropic Claude to assist with code development, numerical analysis, and manuscript review.

Appendix A: Numerical verification

This appendix details the numerical evidence supporting the waveform comparisons. Tail-specific criteria and refinement results are in Appendix B.

Independent implementations give consistent results for the sourced response and exterior ringdown. Finite-difference and spectral sourced waveforms on Schwarzschild and uniform SdS at $L/M = 80$ agree to 5.58×10^{-4} in the relative sphere-integrated norm. At $L/M = 320, 640$, the exterior ringdown waveforms from the spectral-element solver and an independent first-order calculation agree to 1.12×10^{-5} in the Schwarzschild-normalized L^2 norm on $15 \leq U/M \leq 45$. These differences are small compared with the reported finite- L effects; the exterior-supported improvement persists under spatial and temporal refinement.

1. Fixed-mode and localized-source convergence

The minimally coupled uniform fixed-mode sequence uses $(N, \Delta\tau/M) = (384, 0.005)$, $(512, 0.00375)$, and $(768, 0.0025)$, evolved to $\tau = 200M$ with boundary output every $0.03M$. The localized-source sequence uses $(N_r, \Delta\tau/M, \ell_{\max}) = (1024, 0.001, 42)$, $(1536, 1/1500, 46)$, and $(2048, 0.0005, 50)$, evolved to $\tau = 60M$ with output at every timestep. Each finite- L run has a separately evolved Schwarzschild control at the same resolution. Radial resolution, timestep, and, for the localized source, angular truncation are refined together.

We test convergence of waveform differences directly. For a residual $R_q = X_q - Y_q$ at refinement level q , the medium-to-fine change is

$$\delta_{mf} = \frac{\|R_{\text{medium}} - R_{\text{fine}}\|}{\|W_{\text{Schw, fine}}\|}, \quad (\text{A1})$$

using the fixed-mode or Parseval norm as appropriate. Richardson estimates use an effective refinement parameter $h_q = N_q^{-1}$ and an order p inferred from the two successive changes with their actual resolution ratios. This order describes the joint refinement. The estimated remaining fine-grid error is $\delta_{mf}/[(N_f/N_m)^p - 1]$; the larger of this estimate and the observed change sets the numerical scale used for the direct accuracy thresholds. These

TABLE III. Minimally coupled extrapolation checks. Each entry is the maximum over the cumulative intervals ending at $U/M = 40, 80, 160$. Fine-grid residuals, observed medium-to-fine residual changes, and propagated Richardson estimates are reported separately, in percent.

comparison	residual	change	estimate
$W_{\infty}^{(80)} - W_{\text{Schw}}$	0.0222	0.2666	0.0777
$W_{\infty}^{(160)} - W_{\text{Schw}}$	0.0265	0.1430	0.0413
$W_{\infty}^{(80)} - W_{\infty}^{(160)}$	0.0313	0.4092	0.1189

tests quantify refinement at fixed coefficient and matrix cutoffs. An independent cutoff check for the large- L tail is given in Appendix B1.

For the extrapolants, refinement changes in all three input waveforms propagate into the residual comparisons. Table III summarizes the largest values over the three cumulative intervals. The observed changes reach 0.409%, exceeding the central residuals near 0.03% and supporting the conservative 1% agreement stated in Sec. IV A.

The cumulative norms are dominated by the early signal. On the disjoint interval $40 \leq U/M \leq 80$, extrapolant residuals of 0.096–0.211% have refinement changes of 1.02–3.14%. Refinement also exceeds the residuals on $80 \leq U/M \leq 160$.

Localized-source refinement changes remain at the $10^{-4}\%$ level on the common simulated interval. For the Schwarzschild control, the sphere- and time-integrated relative differences from the fine solution are 3.44×10^{-7} and 7.66×10^{-8} at coarse and medium resolution. Truncating the fine solution at $\ell_{\text{max}} = 42$ and 46 gives omitted-mode norms of 2.29×10^{-9} and 1.26×10^{-10} , respectively. All four values use the fine solution’s Parseval norm for normalization.

2. Exterior ringdown

The minimally coupled quadrupole results in Table II use spectral-element calculations through $U = 80M$ with degrees 24, 32, 40 at $\Delta\tau = 0.05M$, and a degree-32 check at $0.025M$. All three geometries use the displacement data (44) and the same geometric clock. Reported errors use degree 40 and a common fine Schwarzschild reference.

On $15 \leq U/M \leq 45$, the largest successive spatial changes of the paired residuals are 3.05×10^{-9} and 6.73×10^{-11} ; timestep halving gives 5.79×10^{-8} . The largest individual waveform change under timestep halving is 1.96×10^{-6} . All are L^2 fractions normalized by the fine Schwarzschild waveform. Replacing the common reference in both errors by the half-step Schwarzschild solution changes $\Delta E = E_{\text{SdS}} - E_{\chi}$ by at most 2.44×10^{-8} . These changes are small compared with the measured improvements $\Delta E = 0.00408$ and 0.00134 at $L/M = 320$ and 640.

TABLE IV. Largest relative envelope changes on the $L/M = 3072$ Price-law interval, $232 \lesssim U/M \lesssim 382$. Values are fractions for SdS at $\mathcal{H}_c^+$ and Schwarzschild at $\mathcal{S}^+$. The timestep check uses $N = 2048$.

refinement	SdS	Schwarzschild
$N : 1536 \rightarrow 2048$	1.27×10^{-4}	3.58×10^{-4}
$N : 2048 \rightarrow 3072$	4.77×10^{-5}	2.20×10^{-5}
$\Delta\tau/M : 0.0025 \rightarrow 0.00125$	3.29×10^{-8}	3.36×10^{-8}

Appendix B: Tail decay and numerical accuracy

We describe how the tail decay rates are measured and checked for numerical accuracy.

1. Schwarzschild tail at large L

At $L/M = 3072$, we compare minimally coupled SdS at $\mathcal{H}_c^+$ with Schwarzschild at $\mathcal{S}^+$, using the same compact $\ell = 1$ velocity data and geometric clock. We use $N = 1536, 2048, 3072$ at $\Delta\tau = 0.0025M$ and halve the timestep at $N = 2048$.

We measure the local power index by fitting $\ln A$ against $\ln U$ over $40M$, using the $10M$ RMS envelope (63). Both indices must stay within 5% of $p = 3$ and satisfy $|p_{\text{SdS}} - p_{\text{Schw}}| \leq 0.15$ for a continuous $150M$ interval containing $U = 300M$. The two finest resolutions must also agree in envelope amplitude to within 1%. The reported interval ends at departure from this criterion: $232 \lesssim U/M \lesssim 382$. Table IV shows the changes under refinement.

Departure occurs near $U = 382M$ and varies by about $2M$ with spatial resolution. Halving the timestep leaves it unchanged. This spread is measured with the decay-rate estimator held fixed.

A further check at $N = 2048$ and $\Delta\tau = 0.0025M$ lowers the truncation thresholds for coefficients and matrices from $(10^{-6}, 10^{-12})$ to $(10^{-10}, 0)$. It extends to $U = 450M$ and also uses a different computing platform. The largest relative envelope change is 2.5×10^{-4} , exceeding the finest spatial-refinement change; the largest absolute index change is 3.5×10^{-4} . Departure moves about $0.8M$ earlier, leaving the Price-law conclusion unchanged.

The main runs extend to $U/M \simeq 1984$, or $\kappa_c U = 0.645$. To identify cosmological decay, we require the envelope to exceed ten times the numerical error estimated from refinement. The intermediate decay rate agrees with Schwarzschild to within 5%.

2. Uniform and exterior-supported tails

The $L/M = 640$ comparison in Fig. 5 uses the spectral-element method, with degrees 24, 32, 40 at $\Delta\tau = 0.1M$ and a degree-40 run at $0.05M$, through $U = 1000M$.

TABLE V. Longest intervals with $p_{\text{eff}} = 3 \pm 10\%$ that pass the numerical checks, for $\ell = 1$ and $L/M = 640$. The required duration is $40M$; the Schwarzschild interval ends with the usable data. The waveform changes δ_s and δ_t are relative L^2 norms.

case	$\Delta U/M$	δ_s	δ_t
Schwarzschild	≥ 834.6	4.59×10^{-11}	1.58×10^{-7}
uniform, $\xi = 0$	9.6	1.82×10^{-10}	2.18×10^{-7}
uniform, $\xi = 1/6$	141.5	1.82×10^{-10}	1.83×10^{-7}
exterior, $\xi = 0$	13.3	2.43×10^{-7}	3.06×10^{-8}
exterior, $\xi = 1/6$	8.5	3.52×10^{-6}	1.16×10^{-7}

Schwarzschild and uniform SdS use nine elements; the exterior cases use sixteen. The figure uses degree 40 and the smaller timestep, with the same Schwarzschild reference for both couplings.

Using the same envelope and fit as above, we require p_{eff} to remain within 10% of 3 for at least $40M$. We retain an index only where the envelope exceeds ten times the numerical error estimated from refinement and the index changes by less than 0.1 under both degree-32 to 40 refinement and timestep halving. We omit times within half a fit window of a waveform zero. Envelope widths $5M$, $10M$, and $20M$, combined with fit widths $30M$, $40M$, and $60M$, leave the outcome of this test unchanged.

Only uniform conformal SdS meets the $40M$ requirement among the finite- L cases, on $153 \lesssim U/M \lesssim 295$ (Table V). Both exterior cases fail the test. Spatial differences decrease from the degree-24–32 comparison to the degree-32–40 comparison in every case. The agreement concerns the decay rate alone.

Here δ_s compares degrees 32 and 40; δ_t measures the change on halving the timestep. Each is divided by the finer waveform norm and maximized over 150 – $300M$, 300 – $500M$, 500 – $750M$, and 750 – $950M$ for the same geometry and coupling.

For cosmological decay, we measure $\gamma_{\text{eff}} = -d \ln A/dU$. At large L , uniform SdS has the expected rate [75]

$$\frac{\gamma_{\ell,\xi}}{\kappa_c} = \ell + \frac{3}{2} - \frac{1}{2} \sqrt{9 - 48\xi} + \mathcal{O}(M/L), \quad (\text{B1})$$

giving dipole rates $\gamma_{1,0}/\kappa_c \simeq 1$ and $\gamma_{1,1/6}/\kappa_c \simeq 2$. Neither uniform waveform stays within 10% of its expected rate for a full exponential-fit interval $\Delta(\kappa_c U) = 0.25$ before the runs end at $\kappa_c U \simeq 1.56$. Cosmological decay is also unresolved for the exterior profiles.

3. Cosmological decay at finite radius

At $r = 8M$, we compare minimally coupled $\ell = 1$ signals at $L/M = 80, 160$ with Schwarzschild. Departure ends the last sustained agreement with Schwarzschild before cosmological entry; entry begins the first sustained

agreement with the expected SdS rate. We vary smoothing, envelope width, required duration, and tolerance over 54 settings, requiring both times to be found in at least half of them. Quoted medians and ranges use the settings that identify both times.

For $L/M = 80, 160$, the entry ranges are $\kappa_c U \in [2.56, 3.07]$ and $[2.68, 3.19]$; the departure ranges are $[1.15, 1.76]$ and $[0.71, 0.96]$. These ranges accompany the medians quoted in Sec. IV D. Both departures precede the settled Schwarzschild Price tail, which begins near $U = 220M$ under the 5% criterion. The observed transition thus connects a Schwarzschild-like transient to cosmological decay.

Appendix C: Exterior profile and boundary limits

The exterior profile keeps the transition layer and outer exact-SdS layer at nonvanishing compactified widths. We specify this profile and derive its geometric, boundary-coefficient, and clock limits.

1. Transition profile and large- L limits

Using (81), define the Chebyshev angle $\vartheta(r) = \arccos[2\rho_\chi(r) - 1]$ on $2M \leq r \leq r_c$. Chebyshev collocation points are uniformly spaced in ϑ . For $\beta = 9/10$, set

$$\begin{aligned} \hat{R}_1(L) &= \beta r_c, \\ \hat{\rho}_{\chi,1}(L) &= \frac{1 - 2M/\hat{R}_1}{D_L}, \\ \hat{\vartheta}_1(L) &= \arccos(2\hat{\rho}_{\chi,1} - 1). \end{aligned} \quad (\text{C1})$$

The transition edges obey

$$\begin{aligned} \frac{L_\star}{M} &= 160, \\ \vartheta_\star &= \hat{\vartheta}_1(L_\star) \simeq 0.07526, \\ \vartheta_1(L) &= \max\{\hat{\vartheta}_1(L), \vartheta_\star\}, \\ \vartheta_0(L) &= 2\vartheta_1(L), \\ \rho_{\chi,i}(L) &= \frac{1 + \cos \vartheta_i(L)}{2}, \\ R_i(L) &= \frac{2M}{1 - D_L \rho_{\chi,i}(L)} \quad (i = 0, 1). \end{aligned} \quad (\text{C2})$$

Thus $2M < R_0 < R_1 < r_c$, and the transition and outer layers have equal angular widths, $\vartheta_0 - \vartheta_1 = \vartheta_1$. For $L \leq L_\star$, $R_1 = \beta r_c$; the $L/M = 80, 160$ members retain the horizon-scaled profile. For $L > L_\star$, both widths equal $\vartheta_\star$, with fixed edge coordinates $\rho_{\chi,0}^\star = (1 + \cos 2\vartheta_\star)/2$ and $\rho_{\chi,1}^\star = (1 + \cos \vartheta_\star)/2$. The compactified widths are

$$\begin{aligned} \Delta\rho_{\text{tr}} &= \rho_{\chi,1}^\star - \rho_{\chi,0}^\star \simeq 0.004239, \\ \Delta\rho_{\text{layer}} &= 1 - \rho_{\chi,1}^\star \simeq 0.001416. \end{aligned} \quad (\text{C3})$$

The smooth switch (71) uses

$$\zeta_L(r) = \frac{\vartheta_0 - \vartheta(r)}{\vartheta_0 - \vartheta_1}. \quad (\text{C4})$$

At the two tested lengths above L_* , the radii are

L/M	R_0/M	R_1/M
320	168.2	260.5
640	228.1	440.4

(C5)

Their finite limits are

$$\begin{aligned} \frac{R_{0,\infty}}{M} &= \frac{2}{1 - \rho_{\chi,0}^*} \simeq 353.7, \\ \frac{R_{1,\infty}}{M} &= \frac{2}{1 - \rho_{\chi,1}^*} \simeq 1412.9, \\ \frac{R_0}{r_c} &\longrightarrow 0, \quad \frac{R_1}{r_c} \longrightarrow 0. \end{aligned} \quad (\text{C6})$$

The transition therefore approaches a finite, distant interval, while the exact-SdS layer extends from its outer edge to r_c . Since $R_0 > 6M$ throughout the numerical sequence, the strong-field region remains exactly Schwarzschild. Equation (78) gives

$$\frac{\Delta m_{\text{tr}}}{M} \sim \frac{1}{2} \left(\frac{R_{1,\infty}}{M} \right)^3 \left(\frac{M}{L} \right)^2 \longrightarrow 0. \quad (\text{C7})$$

For an areal width $\Delta = R_1 - R_0$, the derivative scales are

$$\lambda' = \mathcal{O}\left(\frac{\Lambda}{\Delta}\right), \quad \lambda'' = \mathcal{O}\left(\frac{\Lambda}{\Delta^2}\right). \quad (\text{C8})$$

Narrow layers can generate large curvature and sharp scattering features. Here $\Delta \rightarrow R_{1,\infty} - R_{0,\infty} \simeq 1059.2M$, so the limiting profile has $\chi'_L = \mathcal{O}(M^{-1})$ and $\chi''_L = \mathcal{O}(M^{-2})$. For $\lambda = 3\chi_L/L^2$ at fixed M ,

$$\lambda' = \mathcal{O}(L^{-2}M^{-1}), \quad \lambda'' = \mathcal{O}(L^{-2}M^{-2}). \quad (\text{C9})$$

The source eigenvalues and Ricci curvature in the transition are therefore $\mathcal{O}(L^{-2})$, although the lapse change near r_c remains order unity.

For conformal coupling, the potential (28) becomes

$$V_{\ell,1/6}^\chi = f_\chi \left[\frac{\ell(\ell+1)}{r^2} + \frac{2M}{r^3} + \frac{r\lambda'}{9} + \frac{r^2\lambda''}{18} \right]. \quad (\text{C10})$$

It equals the Schwarzschild potential below R_0 and the corresponding SdS potential above R_1 .

2. Boundary coefficients and clock asymptotics

The coefficients follow from Eqs. (31) and (32) with $G_\chi = 2M/(D_L r^2)$; removable horizon limits are assigned analytically using (39). The bridge crosses both horizons regularly:

$$\begin{aligned} \lim_{r \rightarrow 2M^+} \frac{1 - B_\chi^2}{f_\chi} &= 8, \\ \lim_{r \rightarrow r_c^-} \frac{1 - B_\chi^2}{f_\chi} &= \frac{16M^2}{r_c(r_c - 3M)} > 0. \end{aligned} \quad (\text{C11})$$

In the exact-SdS layer, $D_L = r_c^2/L^2 = 1 - 2M/r_c$, giving

$$\begin{aligned} A_\chi(1) &= \frac{r_c - 3M}{8M(r_c - 2M)}, \quad c_{+, \chi}(1) = 2A_\chi(1), \\ P_{\ell, \xi}^\chi(1) &= \frac{D_L}{2M} \left[\ell(\ell+1) - 2 + \frac{6M}{r_c} + 12\xi D_L \right]. \end{aligned} \quad (\text{C12})$$

Their flat limits are

$$\begin{aligned} A_\chi(1) &\longrightarrow \frac{1}{8M}, \quad c_{+, \chi}(1) \longrightarrow \frac{1}{4M}, \\ P_{\ell,0}^\chi(1) &\longrightarrow \frac{\ell(\ell+1) - 2}{2M}, \quad P_{\ell,1/6}^\chi(1) \longrightarrow \frac{\ell(\ell+1)}{2M}. \end{aligned} \quad (\text{C13})$$

For minimal coupling, convergence of the lower-order potential near the outer boundary remains nonuniform. With $\delta = 2M/r_c$, the potential in the outer layer contains

$$-\frac{D_L r^2}{ML^2} = -\frac{D_L^2}{M} \left[\frac{\delta}{\delta + D_L(1 - \rho_\chi)} \right]^2. \quad (\text{C14})$$

Its boundary amplitude is order M^{-1} , while its compactified width shrinks as M/L . Fixed geometric widths thus do not ensure accuracy at arbitrarily large L and fixed resolution. Conformal coupling cancels this term; the transition terms still require resolution checks.

The clock integrand has the finite horizon limit

$$\lim_{r \rightarrow r_c} \frac{1 + B_\chi}{f_\chi} = \frac{B'_\chi(r_c)}{f'_\chi(r_c)} = \frac{8M^2}{r_c(r_c - 3M)}. \quad (\text{C15})$$

For its flat limit, choose a fixed $R > R_{1,\infty}$. At sufficiently large L , $[R, r_c]$ is exactly SdS, while on $[r_0, R]$ the integrand in (83) differs from its Schwarzschild value by $\mathcal{O}(L^{-2})$. In the outer layer, a cancellation-free form is

$$\frac{1 + B_\chi}{f_\chi} = \frac{8M^2(r_c + r)}{r^2 r_c^2 [(r_c + r)/L^2 - 2M/(r r_c)]}. \quad (\text{C16})$$

Its integrated difference from the Schwarzschild integrand is $\mathcal{O}[M^3 \log(r_c/R)/r_c^2]$, while the omitted Schwarzschild tail is $8M^2/r_c + \mathcal{O}(M^3/r_c^2)$. Together these give $q_\chi = q_0 - 8M^2/r_c + o(M^2/r_c)$, with the same normalization $q_0 = 4M \ln[r_0/(r_0 - 2M)]$ as in (84).

-
- [1] C. W. Misner, Hyperboloidal slices and artificial cosmology for numerical relativity (2004), arXiv:gr-qc/0409073.
- [2] C. W. Misner, Over the rainbow: Numerical relativity beyond scri+ (2005), arXiv:gr-qc/0512167.
- [3] C. W. Misner, J. R. van Meter, and D. R. Fiske, Excising das All: Evolving Maxwell waves beyond scri, *Phys. Rev. D* **74**, 064003 (2006), arXiv:gr-qc/0603034.
- [4] N. Aghanim *et al.* (Planck), Planck 2018 results. VI. cosmological parameters, *Astron. Astrophys.* **641**, A6 (2020), arXiv:1807.06209 [astro-ph.CO].
- [5] J. Winicour, Characteristic evolution and matching, *Living Rev. Relativ.* **15**, 2 (2012), arXiv:0810.1903 [gr-qc].
- [6] S. Ma, M. A. Scheel, J. Moxon, K. C. Nelli, N. Deppe, L. E. Kidder, W. Throwe, and N. L. Vu, Merging black holes with Cauchy-characteristic matching: Computation of late-time tails, *Phys. Rev. D* **112**, 024003 (2025), arXiv:2412.06906 [gr-qc].
- [7] H. Friedrich, Cauchy problems for the conformal vacuum field equations in general relativity, *Commun. Math. Phys.* **91**, 445 (1983).
- [8] J. Frauendiener, Numerical treatment of the hyperboloidal initial value problem for the vacuum Einstein equations. II. The evolution equations, *Phys. Rev. D* **58**, 064003 (1998), arXiv:gr-qc/9712052.
- [9] P. Hübner, A scheme to numerically evolve data for the conformal Einstein equation, *Class. Quant. Grav.* **16**, 2823 (1999), arXiv:gr-qc/9903088.
- [10] P. Hübner, From now to timelike infinity on a finite grid, *Class. Quant. Grav.* **18**, 1871 (2001), arXiv:gr-qc/0010069.
- [11] J. Frauendiener and M. Hein, Numerical evolution of axisymmetric, isolated systems in general relativity, *Phys. Rev. D* **66**, 124004 (2002), arXiv:gr-qc/0207094.
- [12] S. Husa, Problems and successes in the numerical approach to the conformal field equations, in *The Conformal Structure of Space-Time: Geometry, Analysis, Numerics*, Lecture Notes in Physics, Vol. 604 (Springer, 2002) pp. 239–260, arXiv:gr-qc/0204043.
- [13] H. Friedrich, Einstein equations and conformal structure: Existence of anti-de Sitter-type space-times, *J. Geom. Phys.* **17**, 125 (1995).
- [14] H. Friedrich, Conformal geodesics on vacuum space-times, *Commun. Math. Phys.* **235**, 513 (2003), arXiv:gr-qc/0201006.
- [15] A. Zenginoğlu, Numerical calculations near spatial infinity, in *Journal of Physics: Conference Series*, Vol. 66 (2007) p. 012027.
- [16] A. Zenginoğlu, A conformal approach to numerical calculations of asymptotically flat spacetimes, arXiv preprint arXiv:0711.0873 (2007).
- [17] F. Beyer, J. Frauendiener, C. Stevens, and B. Whale, The numerical initial boundary value problem for the generalized conformal field equations, *Phys. Rev. D* **96**, 084020 (2017), arXiv:1706.01416 [gr-qc].
- [18] J. Frauendiener and C. Stevens, The non-linear perturbation of a black hole by gravitational waves. I. The Bondi-Sachs mass loss, *Class. Quant. Grav.* **38**, 194002 (2021), arXiv:2105.09515 [gr-qc].
- [19] J. Frauendiener, C. Stevens, and S. Thwala, Fully non-linear gravitational wave simulations from past to future null infinity, *Phys. Rev. Lett.* **134**, 161401 (2025), arXiv:2504.02188 [gr-qc].
- [20] B. Camden, J. Frauendiener, J. Galinski, K. Pillay, C. Stevens, and S. Thwala, A numerical framework for studying asymptotic quantities, *Gen. Relativ. Gravit.* **57**, 64 (2025), arXiv:2503.17631 [gr-qc].
- [21] J. Frauendiener and C. Stevens, The non-linear perturbation of a black hole by gravitational waves. II. Quasinormal modes and the compactification problem, *Class. Quant. Grav.* **40**, 125006 (2023), arXiv:2211.13276 [gr-qc].
- [22] A. Zenginoğlu, Hyperboloidal foliations and scri-fixing, *Class. Quant. Grav.* **25**, 145002 (2008), arXiv:0712.4333 [gr-qc].
- [23] A. Zenginoğlu, A hyperboloidal study of tail decay rates for scalar and Yang-Mills fields, *Class. Quant. Grav.* **25**, 175013 (2008), arXiv:0803.2018 [gr-qc].
- [24] A. Zenginoğlu, D. Núñez, and S. Husa, Gravitational perturbations of Schwarzschild spacetime at null infinity and the hyperboloidal initial value problem, *Class. Quant. Grav.* **26**, 035009 (2009), arXiv:0810.1929 [gr-qc].
- [25] A. Zenginoğlu and M. Tiglio, Spacelike matching to null infinity, *Phys. Rev. D* **80**, 024044 (2009), arXiv:0906.3342 [gr-qc].
- [26] I. Rácz and G. Z. Tóth, Numerical investigation of the late-time Kerr tails, *Class. Quant. Grav.* **28**, 195003 (2011), arXiv:1104.4199 [gr-qc].
- [27] S. Bernuzzi, A. Nagar, and A. Zenginoğlu, Binary black hole coalescence in the large-mass-ratio limit: The hyperboloidal layer method and waveforms at null infinity, *Phys. Rev. D* **84**, 084026 (2011), arXiv:1107.5402 [gr-qc].
- [28] A. Zenginoğlu and G. Khanna, Null infinity waveforms from extreme-mass-ratio inspirals in Kerr spacetime, *Phys. Rev. X* **1**, 021017 (2011), arXiv:1108.1816 [gr-qc].
- [29] E. Harms, S. Bernuzzi, and B. Brügmann, Numerical solution of the 2+1 Teukolsky equation on a hyperboloidal and horizon penetrating foliation of Kerr and application to late-time decays, *Class. Quant. Grav.* **30**, 115013 (2013), arXiv:1301.1591 [gr-qc].
- [30] R. Panosso Macedo and M. Ansorg, Axisymmetric fully spectral code for hyperbolic equations, *J. Comput. Phys.* **276**, 357 (2014), arXiv:1402.7343 [physics.comp-ph].
- [31] E. Harms, S. Bernuzzi, A. Nagar, and A. Zenginoğlu, A new gravitational wave generation algorithm for particle perturbations of the Kerr spacetime, *Class. Quant. Grav.* **31**, 245004 (2014), arXiv:1406.5983 [gr-qc].
- [32] K. Csukás, I. Rácz, and G. Z. Tóth, Numerical investigation of the dynamics of linear spin s fields on a Kerr background: Late-time tails of spin fields, *Phys. Rev. D* **100**, 104025 (2019), arXiv:1905.09082 [gr-qc].
- [33] L. J. Gomes Da Silva, R. Panosso Macedo, J. E. Thompson, J. A. Valiente Kroon, L. Durkan, and O. Long, Hyperboloidal discontinuous time-symmetric numerical algorithm with higher order jumps for gravitational self-force computations in the time domain (2023), arXiv:2306.13153 [gr-qc].
- [34] A. Vaswani, L. Barack, O. Long, and R. Panosso Macedo, Time-domain framework for the Teukolsky equation with a particle source using comoving hyperboloidal coordinates (2026), arXiv:2606.26339 [gr-qc].
- [35] J. L. Ripley, N. Loutrel, E. Giorgi, and F. Pretorius, Numerical computation of second order vacuum pertur-

- bations of Kerr black holes, *Phys. Rev. D* **103**, 104018 (2021), arXiv:2010.00162 [gr-qc].
- [36] H. Zhu, J. L. Ripley, A. Cárdenas-Avendaño, and F. Pretorius, Challenges in quasinormal mode extraction: Perspectives from numerical solutions to the Teukolsky equation, *Phys. Rev. D* **109**, 044010 (2024), arXiv:2309.13204 [gr-qc].
- [37] A. Zenginoğlu and L. E. Kidder, Hyperboloidal evolution of test fields in three spatial dimensions, *Phys. Rev. D* **81**, 124010 (2010), arXiv:1004.0760 [gr-qc].
- [38] A. Zenginoğlu and C. R. Galley, Caustic echoes from a Schwarzschild black hole, *Phys. Rev. D* **86**, 064030 (2012), arXiv:1206.1109 [gr-qc].
- [39] H. Yang, A. Zimmerman, A. Zenginoğlu, F. Zhang, E. Berti, and Y. Chen, Quasinormal modes of nearly extremal kerr spacetimes: spectrum bifurcation and power-law ringdown, *Physical Review D—Particles, Fields, Gravitation, and Cosmology* **88**, 044047 (2013).
- [40] C. Peterson, S. Gautam, I. Rainho, A. Vañó-Viñuales, and D. Hilditch, 3D evolution of a semilinear wave model for the Einstein field equations on compactified hyperboloidal slices, *Phys. Rev. D* **108**, 024067 (2023), arXiv:2303.16190 [gr-qc].
- [41] O. Rinne, A hyperboloidal method for numerical simulations of multidimensional nonlinear wave equations: nonlinear tails, *Nonlinearity* **38**, 105026 (2025), arXiv:2507.00674 [math.NA].
- [42] A. Zenginoğlu, S. Bernuzzi, and A. Nützi, Semilinear wave equations in homothetic hyperboloidal coordinates and tail decay (2026), arXiv:2608.08863 [gr-qc].
- [43] A. Zenginoğlu, A Geometric framework for black hole perturbations, *Phys. Rev. D* **83**, 127502 (2011), arXiv:1102.2451 [gr-qc].
- [44] M. Ansorg and R. Panosso Macedo, Spectral decomposition of black-hole perturbations on hyperboloidal slices, *Phys. Rev. D* **93**, 124016 (2016), arXiv:1604.02261 [gr-qc].
- [45] R. Panosso Macedo, J. L. Jaramillo, and M. Ansorg, Hyperboloidal slicing approach to quasi-normal mode expansions: The Reissner–Nordström case, *Phys. Rev. D* **98**, 124005 (2018), arXiv:1809.02837 [gr-qc].
- [46] R. Panosso Macedo, Hyperboloidal framework for the Kerr spacetime, *Class. Quant. Grav.* **37**, 065019 (2020), arXiv:1910.13452 [gr-qc].
- [47] R. Panosso Macedo and A. Zenginoğlu, Hyperboloidal approach to quasinormal modes, *Front. in Phys.* **12**, 1497601 (2024), arXiv:2409.11478 [gr-qc].
- [48] J. Besson and J. L. Jaramillo, Quasi-normal mode expansions of black hole perturbations: a hyperboloidal Keldysh’s approach, *Gen. Rel. Grav.* **57**, 110 (2025), arXiv:2412.02793 [gr-qc].
- [49] J. L. Jaramillo, R. Panosso Macedo, and L. Al Sheikh, Pseudospectrum and black hole quasinormal mode instability, *Phys. Rev. X* **11**, 031003 (2021), arXiv:2004.06434 [gr-qc].
- [50] E. Gasperín and J. L. Jaramillo, Energy scales and black hole pseudospectra: the structural role of the scalar product, *Class. Quant. Grav.* **39**, 115010 (2022), arXiv:2107.12865 [gr-qc].
- [51] K. Destounis, V. Boyanov, and R. Panosso Macedo, Pseudospectrum of de Sitter black holes, *Phys. Rev. D* **109**, 044023 (2024), arXiv:2312.11630 [gr-qc].
- [52] R. Panosso Macedo, B. Leather, N. Warburton, B. Wardell, and A. Zenginoğlu, Hyperboloidal method for frequency-domain self-force calculations, *Phys. Rev. D* **105**, 104033 (2022), arXiv:2202.01794 [gr-qc].
- [53] R. Panosso Macedo, P. Bourg, A. Pound, and S. D. Upton, Multidomain spectral method for self-force calculations, *Phys. Rev. D* **110**, 084008 (2024), arXiv:2404.10083 [gr-qc].
- [54] B. Leather, Gravitational self-force with hyperboloidal slicing and spectral methods, *Gen. Rel. Grav.* **57**, 112 (2025), arXiv:2411.14976 [gr-qc].
- [55] B. Leather, A. Spiers, A. Pound, S. D. Upton, B. Wardell, L. Durkan, and N. Warburton, Second-order Teukolsky calculations for nonspinning, quasicircular binaries (2026), arXiv:2609.08995 [gr-qc].
- [56] J. Frauendiener, Conformal infinity, *Living Rev. Rel.* **3**, 4 (2000).
- [57] A. Zenginoğlu, Hyperboloidal evolution with the Einstein equations, *Class. Quant. Grav.* **25**, 195025 (2008), arXiv:0808.0810 [gr-qc].
- [58] A. Vañó-Viñuales, S. Husa, and D. Hilditch, Spherical symmetry as a test case for unconstrained hyperboloidal evolution, *Class. Quant. Grav.* **32**, 175010 (2015), arXiv:1412.3827 [gr-qc].
- [59] A. Vañó-Viñuales and S. Husa, Spherical symmetry as a test case for unconstrained hyperboloidal evolution II: gauge conditions, *Class. Quant. Grav.* **35**, 045014 (2018), arXiv:1705.06298 [gr-qc].
- [60] A. Vañó-Viñuales and T. Valente, Height-function-based 4D reference metrics for hyperboloidal evolution, *Gen. Relativ. Gravit.* **56**, 135 (2024), arXiv:2408.08952 [gr-qc].
- [61] J. D. Álvares and A. Vañó-Viñuales, Free hyperboloidal evolution of the Einstein–Maxwell–Klein–Gordon system, *Class. Quant. Grav.* **42**, 155018 (2025), arXiv:2505.17176 [gr-qc].
- [62] D. Hilditch, Dual foliation formulations of general relativity (2015), arXiv:1509.02071 [gr-qc].
- [63] D. Hilditch, E. Harms, M. Bugner, H. Rüter, and B. Brügmann, The evolution of hyperboloidal data with the dual foliation formalism: Mathematical analysis and wave equation tests, *Class. Quant. Grav.* **35**, 055003 (2018), arXiv:1609.08949 [gr-qc].
- [64] M. Duarte, E. Gasperín, J. C. Feng, and D. Hilditch, Regularizing dual-frame generalized harmonic gauge at null infinity, *Class. Quant. Grav.* **40**, 025011 (2023), arXiv:2206.13661 [gr-qc].
- [65] C. Peterson, S. Gautam, A. Vañó-Viñuales, and D. Hilditch, Spherical evolution of the generalized harmonic gauge formulation of general relativity on compactified hyperboloidal slices, *Phys. Rev. D* **110**, 124033 (2024), arXiv:2409.02994 [gr-qc].
- [66] C. Peterson and D. Hilditch, Strong hyperboloidal compactification for the spherical dual-foliation-generalized harmonic gauge formulation of GR, *Phys. Rev. D* **112**, 024078 (2025), arXiv:2506.03078 [gr-qc].
- [67] V. Moncrief and O. Rinne, Regularity of the Einstein Equations at Future Null Infinity, *Class. Quant. Grav.* **26**, 125010 (2009), arXiv:0811.4109 [gr-qc].
- [68] O. Rinne, An Axisymmetric evolution code for the Einstein equations on hyperboloidal slices, *Class. Quant. Grav.* **27**, 035014 (2010), arXiv:0910.0139 [gr-qc].
- [69] O. Rinne and V. Moncrief, Hyperboloidal Einstein–matter evolution and tails for scalar and Yang–Mills fields, *Class. Quant. Grav.* **30**, 095009 (2013), arXiv:1301.6174 [gr-qc].

- [70] L. T. Buchman, H. P. Pfeiffer, and J. M. Bardeen, Black hole initial data on hyperboloidal slices, *Phys. Rev. D* **80**, 084024 (2009), arXiv:0907.3163 [gr-qc].
- [71] J. M. Bardeen, O. Sarbach, and L. T. Buchman, Tetrad formalism for numerical relativity on conformally compactified constant mean curvature hypersurfaces, *Phys. Rev. D* **83**, 104045 (2011), arXiv:1101.5479 [gr-qc].
- [72] A. Ashtekar, B. Bonga, and A. Kesavan, Asymptotics with a positive cosmological constant: I. Basic framework, *Class. Quant. Grav.* **32**, 025004 (2015), arXiv:1409.3816 [gr-qc].
- [73] A. Ashtekar, B. Bonga, and A. Kesavan, Asymptotics with a positive cosmological constant. II. Linear fields on de Sitter spacetime, *Phys. Rev. D* **92**, 044011 (2015), arXiv:1506.06152 [gr-qc].
- [74] A. Ashtekar, B. Bonga, and A. Kesavan, Asymptotics with a positive cosmological constant: III. The quadrupole formula, *Phys. Rev. D* **92**, 104032 (2015), arXiv:1510.05593 [gr-qc].
- [75] P. R. Brady, C. M. Chambers, W. G. Laarakkers, and E. Poisson, Radiative falloff in Schwarzschild–de Sitter spacetime, *Phys. Rev. D* **60**, 064003 (1999), arXiv:gr-qc/9902010.
- [76] R. Geroch, Limits of spacetimes, *Commun. Math. Phys.* **13**, 180 (1969).
- [77] M. Bugden and C. F. Paganini, The Λ to zero limit of spacetimes and its physical interpretation, *Class. Quant. Grav.* **36**, 045003 (2019), arXiv:1810.00436 [gr-qc].
- [78] Y. Zhou and R. Panosso Macedo, Limiting geometry and spectral instability in Schwarzschild–de Sitter spacetimes, *Phys. Rev. D* **112**, 084063 (2025), arXiv:2507.05370 [gr-qc].
- [79] A. Zenginoğlu, Bridging time across null horizons, *Gen. Rel. Grav.* **57**, 75 (2025), arXiv:2502.08581 [gr-qc].
- [80] R. H. Price, Nonspherical perturbations of relativistic gravitational collapse. I. Scalar and gravitational perturbations, *Phys. Rev. D* **5**, 2419 (1972).
- [81] P. R. Brady, C. M. Chambers, W. Krivan, and P. Laguna, Telling tails in the presence of a cosmological constant, *Phys. Rev. D* **55**, 7538 (1997), arXiv:gr-qc/9611056.
- [82] C. Molina, D. Giugno, E. Abdalla, and A. Saa, Field propagation in de Sitter black holes, *Phys. Rev. D* **69**, 104013 (2004), arXiv:gr-qc/0309079.
- [83] I. G. Dymnikova, The algebraic structure of a cosmological term in spherically symmetric solutions, *Phys. Lett. B* **472**, 33 (2000), arXiv:gr-qc/9912116 [gr-qc].
- [84] I. Dymnikova, Cosmological term as a source of mass, *Class. Quant. Grav.* **19**, 725 (2002), arXiv:gr-qc/0112052 [gr-qc].
- [85] R. Panosso Macedo, Hyperboloidal approach for static spherically symmetric spacetimes: a didactical introduction and applications in black-hole physics, *Phil. Trans. Roy. Soc. Lond. A* **382**, 20230046 (2024), arXiv:2307.15735 [gr-qc].
- [86] G. Mavrogianis, Quasilinear wave equations on Schwarzschild–de Sitter, *Commun. Part. Diff. Eq.* **49**, 38 (2024), arXiv:2111.09495 [gr-qc].
- [87] G. Mavrogianis, Morawetz estimates without relative degeneration and exponential decay on Schwarzschild–de Sitter spacetimes, *Ann. Henri Poincaré* **24**, 3113 (2023), arXiv:2111.09494 [gr-qc].
- [88] R. Gregory, D. Kastor, and J. Traschen, Evolving black holes in inflation, *Class. Quant. Grav.* **35**, 155008 (2018), arXiv:1804.03462 [hep-th].
- [89] P. R. Anderson and J. Traschen, Horizons and correlation functions in 2D Schwarzschild–de Sitter spacetime, *JHEP* **2022** (01), 192, arXiv:2012.08494 [hep-th].
- [90] P. R. Anderson, Z. P. Scofield, and J. Traschen, Linear growth of the two-point function for the Unruh state in $1+1$ dimensional black holes, in *Proceedings of the Sixteenth Marcel Grossmann Meeting* (World Scientific, 2023) pp. 1255–1264, arXiv:2205.00264 [gr-qc].
- [91] S. R. Dolan and A. C. Ottewill, Wave propagation and quasinormal mode excitation on Schwarzschild spacetime, *Phys. Rev. D* **84**, 104002 (2011), arXiv:1106.4318 [gr-qc].
- [92] M. Casals and B. C. Nolan, Global Hadamard form for the Green function in Schwarzschild spacetime, *Phys. Rev. D* **108**, 044033 (2023), arXiv:1606.03075 [gr-qc].
- [93] K. J. Burns, G. M. Vasil, J. S. Oishi, D. Lecoanet, and B. P. Brown, Dedalus: A flexible framework for numerical simulations with spectral methods, *Phys. Rev. Research* **2**, 023068 (2020), arXiv:1905.10388 [physics.comp-ph].
- [94] G. J. Gassner, A skew-symmetric discontinuous galerkin spectral element discretization and its relation to SBP-SAT finite difference methods, *SIAM J. Sci. Comput.* **35**, A1233 (2013).
- [95] E. Hairer and G. Wanner, Stiff differential equations solved by Radau methods, *J. Comput. Appl. Math.* **111**, 93 (1999).
- [96] M. H.-Y. Cheung, K. Destounis, R. P. Macedo, E. Berti, and V. Cardoso, Destabilizing the fundamental mode of black holes: The elephant and the flea, *Phys. Rev. Lett.* **128**, 111103 (2022), arXiv:2111.05415 [gr-qc].
- [97] A. Laeuger, C. Weller, D. Li, and Y. Chen, Ringdown of a black hole surrounded by a thin shell of matter, *Phys. Rev. D* **112**, 084042 (2025), arXiv:2506.00367 [gr-qc].
- [98] M. De Amicis, S. Albanesi, and G. Carullo, Inspiral-inherited ringdown tails, *Phys. Rev. D* **110**, 104005 (2024), arXiv:2406.17018 [gr-qc].
- [99] R. F. Rosato, M. De Amicis, and P. Pani, Singular structures and causality of the Schwarzschild Green’s function in the frequency domain (2026), arXiv:2603.20490 [gr-qc].