PREPRINT

Training clinical agents to ask, test, commit and defer

[Authors to be added]

[Affiliations to be added]

October 2026 · Code and the MIMIC-CDM-CR benchmark: github.com/GIND123/Clinical-Agent-Stanford

Abstract

Background. Interactive diagnostic agents for acute abdominal pain are evaluated in a closed world: they must name one of four diagnoses and cannot hand a case to a clinician. Abstention learned by reinforcement learning is expected to collapse into refusal, and theory recommends thresholding a calibrated confidence instead. Meanwhile, a randomised study has shown that physicians follow misdirecting AI advice, so what an agent hands over matters as much as when it does.

Methods. DEFER-Dx is a Qwen3-8B agent trained with multi-turn reinforcement learning to examine, order any of 22 priced tests, commit to a diagnosis with a stated probability, or defer to a clinician with a probabilistic differential. Each deferral is credited by a counterfactual escalation value: continuations branched from the state where the agent deferred, with deferral disabled, estimate the clinical value of continuing; the handoff is scored with a strictly proper rule. We evaluated on 480 held-out MIMIC-IV-Ext-CDM admissions, on an open-world extension with five time-critical diagnoses held out of training, and on MIMIC-CDM-CR, 32 cases rebuilt from recent published case reports, with two training seeds for DEFER-Dx and for an identically trained agent without deferral.

Results. DEFER-Dx answered 77.6% of in-set cases at 95.2% accuracy (95% CI 93.4–96.8) with 3.8 unflagged errors per 100 cases (zero-shot: 24.1). Against the agent without deferral thresholded at matched coverage, it was more accurate on answered cases in all four seed pairings (+1.1 to +5.3 points; pooled +3.2, 1.6–5.1). On 187 never-seen time-critical admissions both seeds escalated 87% and committed wrongly in 6.8% and 7.1%, against 11.4% for an identically trained deferral reward from prior work. A threshold sent complicated disease to the clinician less often than uncomplicated disease (16.7% vs 24.0%); DEFER-Dx did not. But what the handoff named depended on the training seed: when clinicians were modelled as adopting the handoff's top suggestion as often as physicians followed adversarial AI, one seed's handoffs lowered team accuracy (86.8% vs 91.5% with the threshold) and the other's steered towards wrong in-scope diagnoses on unfamiliar emergencies.

Conclusions. Crediting escalation against the counterfactual of continuing trains stable, well-targeted deferral that outperforms post-hoc thresholding and a published deferral reward. Whether escalation is safe depends on the content of the handoff, which should be trained and evaluated as carefully as the decision to escalate.

Introduction

A patient arrives in the emergency department with abdominal pain. An interactive diagnostic agent asks for the examination, orders a CT and names a diagnosis. Every such agent published on the MIMIC-IV clinical decision-making benchmark (MIMIC-CDM)[1] must name one of four conditions (appendicitis, cholecystitis, diverticulitis or pancreatitis), even when the patient has a perforated ulcer or mesenteric ischaemia, and even when its own evidence is split. None can hand the case back.

Three developments make this gap pressing. Regulators expect clinical decision support to leave the decision with a clinician who can independently review the basis of its recommendations rather than rely on them[2], and a randomised study of 225 physicians has just shown how fragile that safeguard is: conversational AI with a concealed objective raised harmful decisions from 2% to 49%, and clinical expertise did not protect[3]. The field's leaderboard shows the cost of optimising accuracy alone: the strongest reported MIMIC-CDM agent raises case-weighted accuracy to 93.4% while its accuracy on the minority condition, diverticulitis, falls below that of a random test-ordering policy[4]. And recent theory argues that the obvious fix, an "abstain" action trained by reinforcement learning, collapses into refusing everything, and that one should instead threshold a calibrated confidence after training[5].

We ask whether an agent can learn when to defer without collapsing, whether learned deferral beats thresholding the confidence of an identically trained agent, and what happens to the patient once the case reaches the clinician. Our

contributions:

1. **DEFER-Dx**, an agent trained with multi-turn group-relative policy optimisation over four actions: ask, order one of 22 individual tests, commit with a stated probability, or defer with a ranked probabilistic differential.
2. **The counterfactual escalation value (CEV).** At each state where the agent escalates, forced continuations branched from that exact state estimate the clinical value of not escalating. The escalation is credited with the difference, and the handoff is scored with a strictly proper rule. Prior deferral rewards price escalation by case-level difficulty[6–8] or by designated cases[9].
3. **Two open-world benchmarks.** MIMIC-CDM-OW, 713 admissions whose principal diagnosis is none of the four, built with MIMIC-CDM's own text pipeline, with five time-critical diagnosis groups held out of training; and MIMIC-CDM-CR, 32 cases rebuilt from open-access case reports published 2020–2025, which can be shared without a data use agreement.
4. **A head-to-head test of learned versus post-hoc deferral** at exactly matched coverage, with two training seeds per arm, and a model of the clinician who receives the handoff, calibrated to the randomised evidence on AI advice[3].

Results

An agent that asks, tests, commits or defers

Each episode is one emergency admission (Fig. 1). The agent sees the history at presentation and, turn by turn, asks for the examination, orders individual tests (each charged at its 2025 price), commits to one of the four conditions or OTHER with a stated probability, or defers to a clinician with a probabilistic differential. Observations are looked up from the record, never generated. DEFER-Dx was trained for 150 steps on 2,404 patient-disjoint training cases (CDM, OTHER and same-pipeline in-set controls) with the counterfactual escalation value (Fig. 2; Methods) and a PI-controlled cap of 30% on in-set deferral. A second training seed, an identically trained agent without DEFER (the GRPO control; two seeds), a case-level group-consensus deferral reward (the mechanism of prior work[6–8]) and zero-shot Qwen3-8B were evaluated on the same cases (Extended Data Fig. 1).





Fig. 1 | DEFER-Dx. One admission per episode; the agent asks, orders individual priced tests, commits with a stated probability or defers with a probabilistic differential. The label space is open: the four MIMIC-CDM conditions or OTHER.

Learned escalation without collapse

On the 480 held-out CDM admissions (val + test, three evaluation seeds), DEFER-Dx answered 77.6% (74.3–80.6) of cases at 95.2% (93.4–96.8) accuracy, with 3.8 (2.4–5.1) unflagged errors (wrong commitments without escalation) per 100 cases, against 24.1 for zero-shot Qwen3-8B with forced choice and 9.1 with deferral offered in the prompt (paired −5.3, −7.5 to −3.3). Its stated probabilities were calibrated (ECE 0.042; Brier 0.047, the lowest of all systems), and at 70% coverage it was 96.5% accurate (Extended Data Fig. 3). Deferral did not collapse: the PI controller held in-set deferral near its cap without the limit cycle that plain dual ascent produced (Extended Data Fig. 9). The



$R_k = \alpha\,1[d_k{=}y] - \lambda\,(p_k - 1[d_k{=}y])^2 - \kappa\,C[y, d_k] - c \cdot \text{cost(tests after } s)$; $K = 3$. In a forced continuation, a **DEFER** is executed as **COMMIT** of its differential's top; tests remain allowed. V_τ: reward of a calibrated decision that succeeds with probability τ (1.2 on OTHER cases). S: normalised multi-class Brier score of the handed-over differential q (strictly proper). $\mu = 0.1$: handoff cost. ν: coverage multiplier (Fig. S1).

Fig. 2 | The counterfactual escalation value. When an episode ends in DEFER at state s, the environment is replayed to s and K = 3 forced continuations are sampled with escalation disabled (a DEFER executes as COMMIT of the differential's top choice; further tests remain allowed). Their mean clinical return V̂(s) values not escalating; the DEFER turn's advantage is E − V̂(s), where E values the escalation itself, including a strictly proper score of the handed-over differential.

second training seed answered 73.2% (69.7–76.5) at 95.3% (93.2–97.1) with 3.5 unflagged errors per 100. Against a confidence threshold on zero-shot Qwen3-8B at matched coverage, DEFER-Dx was 12.4 points (9.4–15.7) more accurate on answered cases, and 11.1 (7.0–15.3) with three-sample voting for both.

Learned versus post-hoc deferral across training seeds

The fair comparison for learned deferral is a threshold on an identically trained agent[5]. The GRPO control was trained exactly as DEFER-Dx but without DEFER, and its stated probability was thresholded at DEFER-Dx's coverage, cross-fitted between val and test. Stated probabilities are coarse (a single value covers 14–17% of the control's cases), so commitments tied at the cut were kept with a probability fitted to reach the target coverage (Methods); without this, a plain cut overshot the target and overstated the advantage (+3.1 instead of +2.3 for the first seed).

DEFER-Dx was more accurate on the cases it answered in all four pairings of two DEFER-Dx seeds and two control seeds (Fig. 3): +2.3 (0.1–4.6), +5.3 (3.0–7.7), +1.1 (−0.8 to 3.3) and +4.1 (1.9–6.5) points, three of four intervals excluding zero. Pooled over the pairings, the difference was +3.2 points (1.6–5.1), with 2.4 fewer unflagged errors (1.1–3.9) and 2.1 fewer confident errors (1.0–3.3) per 100 cases and a lower area under the risk–coverage curve (−0.014, −0.031 to −0.005); DEFER-Dx's calibration error was higher (ECE +0.036, 0.014–0.050). The margin depended mainly on the control's seed: identically trained controls differed in how well their stated probabilities ranked their own errors (AURC 0.044 and 0.074), whereas DEFER-Dx's accuracy on answered cases was the same in both seeds (95.2% and 95.3%).




Fig. 3 | Learned versus post-hoc deferral across training seeds. Each row compares one DEFER-Dx training seed with one seed of the identically trained GRPO control, thresholded at that DEFER-Dx seed's coverage (paired, same 480 cases, matched coverage). The last row pools the four pairs: cases are resampled with the training runs held fixed, so its interval reflects case sampling; the rows show the variation between runs. Positive values favour DEFER-Dx; filled markers have a 95% interval that excludes 0.

Unfamiliar, time-critical presentations

On 187 admissions from five time-critical diagnosis groups never seen in training (mesenteric ischaemia, perforated or bleeding ulcer, ruptured aortic aneurysm, diabetic ketoacidosis, ectopic pregnancy), both DEFER-Dx seeds escalated 87.0% of cases and committed to one of the four conditions in 6.8% (4.3–10.0) and 7.1% (4.3–10.3) (Fig. 4). Against the identically trained group-consensus reward of prior work (11.4%), the differences were −4.6 (−8.0 to −1.4) and −4.3 (−7.7 to −1.1), and on 206 OTHER admissions from groups seen in training −7.0 (−9.9 to −4.4) and −6.6 (−9.9 to −3.6), at the same accuracy on answered in-set cases (+1.1 and +1.2, intervals including zero). Without a threshold the control committed wrongly in 15.0% (paired −8.2, −12.1 to −4.6). The thresholded control committed wrongly about as rarely as DEFER-Dx (7.5%), but in a different way: it escalated 21.0% of these cases and committed to “none of the four” in 64.3%. On 258 same-pipeline in-set controls DEFER-Dx named OTHER for 0.3%, so its open-world behaviour is not a source shortcut.




Fig. 4 | Open world. Outcomes on OTHER admissions from (a) diagnosis groups seen in training and (b) five time-critical groups never seen in training: wrong commitment to one of the four conditions, deferral, and naming OTHER. 95% case-level bootstrap intervals.

Published case reports

[[MIMIC-CDM-CR results: out-of-scope wrong commitments and escalation per system; held-out vs new groups; in-scope accuracy; complicated vs uncomplicated escalation; Fig. 5.]]

Complicated disease

DEFER-Dx names a diagnosis; it does not stage it, and a surgeon’s management depends on the stage. We labelled each in-set evaluation case as complicated or not from structured data only (admission ICD codes and admission



Fig. 5 | MIMIC-CDM-CR: behaviour on 32 published case reports. Majority outcome over three evaluation seeds for each case (rows) and system (columns): C, correct in-scope diagnosis; E, escalated (DEFER, or COMMIT of OTHER, which names no diagnosis to act on); W, wrong in-scope diagnosis; M, no majority. Rows are grouped by the five held-out time-critical groups, groups in no MIMIC cohort, a group seen in training, and in-scope cases with and without complications by the guideline grade the report supports. Cases and citations: benchmarks/casereports/.

laboratory values), following the guideline used for each condition: WSES and AAST grading for appendicitis[10], WSES staging for diverticulitis[11], the revised Atlanta classification for pancreatitis[12] and the Tokyo 2018 grades for cholecystitis[13]. 114 of 480 cases were complicated (appendicitis 48 of 191, cholecystitis 21 of 130, diverticulitis 21 of 51, pancreatitis 24 of 108).

A confidence threshold escalated complicated disease less often than uncomplicated disease (16.7% vs 24.0%; −7.4, −14.1 to −0.5): complicated presentations produce confident, often correct commitments. DEFER-Dx did not (23.7% vs 22.0%; +1.6, −6.1 to 9.8; second seed 28.1% vs 26.4%) (Extended Data Fig. 6). On the same complicated cases, at matched overall coverage, DEFER-Dx escalated more than the thresholded control in all four seed pairings (+4.4 to +7.0 points; +7.0, 0.6 to 13.7 for the first seeds) and more than the consensus reward (+7.6, 2.9 to 12.9; +12.0, 6.4 to 18.1), with fewer unflagged errors on complicated cases in all four control pairings (−1.8 to −5.0 per 100). No system, however, escalated complicated disease preferentially: none is severity-aware.

What the handoff says, and the clinician who reads it

A handoff is AI advice. Barrit and colleagues randomised 225 physicians to AI that advocated harmful options or correct ones and found 46.9 more harmful decisions per 100 targeted decision points with the misdirecting AI, with almost no detection[3]; incorrect AI suggestions have similarly lowered radiologists' accuracy from 79.7% to 19.8%[14]. We therefore examined what DEFER-Dx hands over and modelled the clinician who receives it (Methods): with probability f the clinician adopts the handoff's top suggestion, otherwise decides independently with the accuracy of the MIMIC-CDM physicians (0.89)[1]. A commitment to OTHER names no diagnosis to act on and was treated as a handoff suggesting OTHER.

The handoff's content depended on the training seed. 68.4% of the first seed's in-set handoffs ranked OTHER first (81% of its in-set handoffs had a wrong top suggestion), against 4.2% for the second seed (32%) and 8.6% for the consensus reward. On never-seen time-critical cases the pattern reversed: the first seed's handoffs had an in-scope top suggestion in 21%, the second seed's in 84%. With an independent clinician (f = 0) every system's team accuracy on in-set cases was about 94% (Fig. 6). At the susceptibility observed under adversarial advocacy (f = 0.47), the first seed's team accuracy fell to 86.8%, against 91.5% for the thresholded control (−4.7, −6.6 to −2.8), whereas the second seed's held (91.4%); on never-seen time-critical cases the second seed's handoffs led to wrong in-scope diagnoses in 37.0%, against 11.0% for the thresholded control and 25.9% for the consensus reward. A learning-to-defer rule on the control's confidence (defer whenever the stated probability is below the clinician's accuracy[15,16]) was the safest on these terms (8.4%), at almost twice the clinician workload (43% of in-set cases, against 23%).

What drives the escalation credit

The estimator. We re-sampled K = 16 forced continuations from 100 in-set and 60 OTHER deferral states of each final policy (Extended Data Fig. 7). The K = 3 estimate used in training gave the credit the same sign as K = 16 in 88% and 92% of in-set states (K = 1: 83% and 88%; K = 8: 95% and 96%). Had it not been able to defer, the first seed's policy would have been right in only 32% of its in-set deferral states, the second seed's in 67%: the first seed's deferrals were better targeted, and 60% of them were credited against 37% of the second seed's.

Test cost. Removing the cost of tests ordered after the deferral changed the sign of the credit in 0% and 1% of states, and forced continuations ordered a test in only 23% and 18% of cases. The early deferral (0.84 tests per case) reflects




Fig. 6 | The clinician who receives the handoff. Team outcomes as a function of the probability f that the clinician adopts the handoff's top suggestion; otherwise the clinician decides independently with accuracy 0.89 (MIMIC-CDM physicians[1]). a, Team accuracy on in-set cases. b, Wrong in-scope diagnoses caused by the AI on never-seen time-critical cases (its own wrong commitments plus clinicians who adopt an in-scope suggestion). The vertical line marks the adversarial-advocacy effect measured by Barrit et al.[3]. A COMMIT of OTHER counts as a handoff suggesting OTHER.

the agent's inability to resolve these cases, not the price of investigating. [[No-cost ablation: DEFER-Dx trained without the test-cost term, against DEFER-Dx at step 100.]]

The clinician the credit assumes. The escalation value credits a deferral with the value of a clinician who decides correctly with probability $\tau = 0.85$. If that clinician were right in 70% instead, 46% rather than 60% of the first seed's in-set deferrals would be credited (30% rather than 37% for the second seed).

Ablations at 100 training steps (Extended Data Fig. 8). A constant deferral reward with the same OTHER value and controller matched DEFER-Dx on accuracy on answered cases (−0.4, −2.3 to 1.5), unflagged errors and wrong commitments on never-seen time-critical cases (10.2% vs 10.7%), but deferred less and rarely named OTHER in its handoffs (24.7% vs 89.0% of never-seen time-critical cases). Removing the handoff score ($\eta = 0$) left accuracy on answered cases unchanged (−1.1, −2.8 to 0.6) but made handoffs less informative (−6.5 points, 3.5 to 9.3, in favour of the score), reduced deferral (coverage 75.9% vs 68.1%) and increased false commitments on OTHER from groups seen in training (9.5% vs 6.3%); OTHER-first handoffs were more common without the score (55.1% vs 35.1%), so they come with the counterfactual credit itself. [[No-cap ablation ($\rho_max = 1$) and DEFER-Dx with the handoff score weighted 1.0.]]

Robustness and equity

With every sentence containing CDM's diagnosis mask removed, DEFER-Dx's accuracy on answered cases changed by −1.0 points (−2.0 to 0.0) and its coverage by +1.6 (−0.2 to 3.5). The pattern of recorded tests predicts the diagnosis in MIMIC-CDM (macro AUROC 0.879 from recorded/not-recorded flags alone), but DEFER-Dx received a "not performed" reply in only 12–17% of episodes, and its accuracy on answered cases was lower after one (89.7% vs 95.7%; second seed 88.5% vs 96.2%), as for every trained model: there is no sign that it profits from test availability. Escalation and unflagged errors did not differ systematically by sex, race, insurance or language; patients aged 80 or older were escalated more often (crude 28.8% and 36.9% for the two seeds), consistent with atypical presentations in older patients (Extended Data Table 2). On LA-CDM's test split, counting handoffs through their top choice, DEFER-Dx reached 76.4% (71.1–81.0) mean-class accuracy and its second seed 85.2% (80.8–89.3), against 82.7% for the control, 86.3% for the consensus reward and 86.9% for zero-shot Qwen3-8B restricted to the four labels (LA-CDM reports 81.3% in its own closed-world environment[17]; these are context, not head-to-head comparisons).

Discussion

Learned deferral did not collapse. Crediting each escalation against what continuing from the same state would have achieved produced an agent that answered three-quarters of in-set cases at 95% accuracy with fewer than four unflagged errors per 100, escalated 87% of dangerous presentations it had never seen, and did both in two independent training runs. It outperformed the deferral reward of prior work on unfamiliar emergencies in both runs,

and outperformed a threshold on an identically trained agent in all four seed pairings; how much it outperformed the threshold depended mainly on the threshold, because identically trained agents differed in how well their confidence ranked their own errors. Theory is right that a proper-scored confidence makes thresholding a strong baseline[5]; in our runs a learned operating point was the more reliable of the two.

The more important finding concerns the handoff. A deferral agent is evaluated as if the clinician who receives the case were independent of it. Under that assumption every system here reached the same team accuracy of about 94%. Physicians are not independent of AI advice[3,14,18], and once the clinician can be steered, the content of the handoff dominates: one DEFER-Dx run labelled uncertain in-set cases “none of the four”, the other escalated unfamiliar emergencies with an in-scope suggestion, and either pattern would have cost patients under the susceptibility measured in physicians. Both runs were trained with the same strictly proper handoff score; at $\eta = 0.3$ it improved handoff content (the no-handoff ablation) but did not pin down which kind of differential the agent learned. A threshold, by contrast, hands over its own withheld guess, and a learning-to-defer rule hands over fewer guesses but doubles the clinician’s workload. Regulatory guidance asks that clinicians be able to review the basis of a recommendation independently[2]; our results suggest that agents should be trained and evaluated against a clinician who might not.

Complicated disease is where a surgeon’s decision matters most. A threshold sent it to the clinician less often than uncomplicated disease, because complicated presentations look certain; DEFER-Dx did not, but neither did it seek complicated disease out. Staging is a capability none of the agents has[19]: a clinically useful agent would name the condition, grade it by the relevant guideline[10–13] and escalate by grade.

Implications. Learned deferral should be compared with a threshold on an identically trained model, at exactly matched coverage, with more than one training run of each, since both the coarse confidences of language models and seed-to-seed variation make unmatched or single-run comparisons easy to misread. Agents that defer should be evaluated in the open world, on external cases, and with a model of the clinician who receives the case; the handoff should be trained with stronger pressure towards the right content and assessed by clinicians.

Limitations. The data come from one centre and one organ system (abdominal pain at BIDMC). Labels for OTHER cases are principal ICD codes, severity labels are proxies from codes and admission laboratory values, and the severity matrix and some test prices are proxies (Methods). The environment reveals only what was recorded, so test availability is informative, and it is static: a patient’s condition does not evolve while tests are pending, and the agent cannot treat and reassess, which real emergency care does. The clinician model is a sensitivity analysis calibrated to an adversarial study, not a measurement with these handoffs; clinician ratings of the handoffs (a blinded rating sheet is in the repository) and a user study are needed. Two training seeds per arm show seed-to-seed variation but cannot estimate it, and the consensus reward has one seed. MIMIC-CDM-CR has 32 cases, too few for precise rates; it is an external check of behaviour, not a validation study. No model here should be used for patient care.

Methods

Data

MIMIC-IV-Ext-CDM 1.1[1] has 2,400 emergency admissions with acute abdominal pain (957 appendicitis, 648 cholecystitis, 538 pancreatitis, 257 diverticulitis). We kept LA-CDM’s 80/10/10 split[17], so its numbers are comparable on its test set. When a test repeats within an admission, the earliest result is used. **MIMIC-CDM-OW** (this work; Extended Data Fig. 1) selects MIMIC-IV 2.2 admissions with an abdominal chief complaint whose principal diagnosis falls in one of nine non-CDM groups and that carry no CDM-condition code at any position. Each case is built with MIMIC-CDM’s own text pipeline (history and examination windows, radiology section filter, masking of the case’s own diagnosis, inclusion rule, earliest-value deduplication under a total row order); rebuilds are byte-identical. It yields 713 OTHER cases and 648 same-pipeline in-set controls, which remove the shortcut “built by the open-world pipeline, so OTHER”. Cohorts are patient-disjoint: 2,404 training cases (1,760 CDM, 279 OTHER, 365 controls), 212 development cases, 480 CDM evaluation cases (val + test), 206 OTHER cases from four groups seen in training, 187 from five time-critical groups held out of training entirely, and 258 in-set controls (Extended Data Table 1). Groups with fewer than 10 cases are reported only pooled.

Environment

The hidden state is the diagnosis y {appendicitis, cholecystitis, diverticulitis, pancreatitis, OTHER}. At reset the agent sees the history at presentation. Actions: ASK(physical examination), free; TEST(x) for 22 individual tests

(blood count, metabolic panels, liver tests, lipase, amylase, CRP, lactate, coagulation, triglycerides, pregnancy test, urinalysis, blood and urine cultures, CT, ultrasound, MRI, abdominal and chest radiographs, HIDA), each charged at its 2025 BIDMC standard charge where published (placeholders otherwise); COMMIT(d, p); and DEFER(q, reason) with a probabilistic differential q. At most eight investigations and two malformed turns are allowed.

Rewards and the counterfactual escalation value

A commitment earns $R = \alpha \cdot 1[d = y] - \lambda(p - 1[d = y])^2 - \kappa \cdot C[y, d] - c \cdot \Sigma\text{cost}$, with $\alpha = \lambda = 1$ (the second term is the Brier score), $\kappa = 0.5$ and C a 5 × 5 asymmetric time-to-harm severity matrix; c makes ordering every test cost one correct diagnosis. For an episode that ends with DEFER at state s, the environment is rebuilt by deterministic replay and K = 3 continuations are sampled from s with escalation disabled (a DEFER is executed as COMMIT of the differential's top choice at its stated probability; further tests remain allowed). Their mean return $\hat{V}(s)$, charging only tests ordered after s, values not escalating. The escalation is valued $E = V_\tau + \eta \cdot S(q, y) - \mu - \nu$, where V_τ is the expected commitment reward of a calibrated decision that succeeds with probability τ on in-set cases (1.2 on OTHER cases), S is the normalised multi-class Brier score of the handed-over differential (strictly proper; $\eta = 0.3$), $\mu = 0.1$ the handoff cost and ν the coverage multiplier. The DEFER turn's advantage is $E - \hat{V}(s)$; earlier turns keep the group-relative advantage. At most 32 deferrals per step are branched; others get no update on their DEFER turn. τ is annealed from 0.95 to 0.85 over the first 100 steps.

Training

Qwen3-8B in thinking mode with LoRA (rank 16, $\alpha = 32$) on every linear layer, trained with multi-turn GRPO on batches of 12 cases × 8 episodes at temperature 1.0 for 150 steps: advantages centred without standard-deviation normalisation, token-level clipped surrogate (ε 0.2/0.28) with truncated importance weights against the sampler's log-probabilities, AdamW, no KL term. Generation (vLLM) and training (PEFT) share one 48 GB GPU. In-set deferral is held at or below 30% by a PI controller on ν ($K_p = 1.5$, $K_i = 0.2$, $\nu_{max} = 2$); plain dual ascent wound up and limit-cycled (Extended Data Fig. 9). The control arm is identical without DEFER; the comparator arm replaces the escalation value with the case-level consensus reward $\gamma(\tau - \hat{p}) - \mu$ ($\gamma = 2$), where $\hat{p}$ is the share of the group's correct commitments (identical to TIAR at $\tau = 0.5$[6]). Second seeds change only the seed (case order, sampling, LoRA initialisation).

Comparators and post-hoc thresholds

Each forced-choice system was thresholded on its stated probability at DEFER-Dx's coverage, fitted on val and applied to test and vice versa. Because stated probabilities are coarse, commitments tied at the cut are kept with probability $k = (\text{target} \times N - n_{above})/n_{tied}$, decided by a fixed hash of the case and evaluation seed; we report one pre-specified draw and the range of the point difference over 20 draws (RESULTS.md §5b). Also evaluated: SGR (5% selective risk, $\delta = 0.05$)[22], self-consistency over three samples, and a learning-to-defer rule that defers whenever the stated probability is below the clinician's accuracy, the Bayes-optimal rejector for a calibrated model and an expert of known accuracy[15,16,21].

Statistics

Every number is given with a 95% case-level bootstrap interval (2,000 resamples, all evaluation seeds of a case resampled together); comparisons are paired bootstraps on the same cases with two-sided p-values. Pooled seed comparisons pool the four pairings with cases resampled and training runs held fixed. Holm-adjusted p-values over all paired tests in the report (one family; two-sided Wald p from the bootstrap standard error) are listed in RESULTS.md §5; the primary comparisons are DEFER-Dx against the thresholded control (pooled) and against the consensus reward on never-seen time-critical cases, and the others are exploratory.

Severity labels

Complicated disease was defined from the admission's ICD-9/10 codes and the first admission laboratory values in the case record (no note text): appendicitis with gangrene, perforation, abscess or generalised peritonitis; diverticuli-

tis with perforation, abscess, peritonitis, fistula or obstruction; pancreatitis with necrosis or pseudocyst, organ-failure codes or creatinine > 1.9 mg/dL; cholecystitis with Tokyo 2018 grade II (WBC > 18 × 10^9/L; perforation, fistula, pericholecystic or liver abscess, peritonitis) or grade III (organ-failure codes; creatinine > 2.0 mg/dL, INR > 1.5, platelets < 100 × 10^9/L) (`scripts/severity_analysis.py`).

The clinician model

For an escalated case (DEFER, or COMMIT of OTHER) the clinician adopts the handoff's top suggestion with probability f and otherwise decides correctly with probability a_c = 0.89, the mean accuracy of the four physicians in the MIMIC-CDM reader study[1]. Episodes without a usable answer carry no suggestion. On OTHER cases we count only AI-caused wrong in-scope diagnoses (the AI's own wrong commitments, and clinicians adopting an in-scope suggestion), because the only measured physician accuracy is for the four-way task. f ranged from 0 to 0.5; 0.47 is the adversarial-minus-aligned difference in harmful decisions per targeted decision point in Barrit et al.[3] (`scripts/team_analysis.py`).

Estimator audit

From the deferral states recorded in each final policy's held-out evaluation (100 in-set, 60 OTHER, sampled at random), K = 16 forced continuations were sampled with the trainer's own branching function and training-time sampling, and each continuation's return was stored by component; ν is the run's final value (`scripts/cev_audit.py`, `scripts/cev_audit_report.py`).

MIMIC-CDM-CR

Europe PMC was searched for open-access case reports with full text published 2020 or later, by target diagnosis; reports with adult patients presenting with abdominal pain were kept, and reports with an ambiguous principal diagnosis, in-hospital post-operative presentations or children were excluded. Each case section was paraphrased into the case format (history with past history, medications and social history; examination; laboratory values at presentation in MIMIC-IV units; imaging findings and impression); tests outside the catalogue were left out and noted, and nothing was added. 19 cases are out of scope (7 from the held-out time-critical groups, 11 from groups in no MIMIC cohort, 1 bowel obstruction) and 13 in scope with their guideline grade (9 complicated). The cases, citations and licences are in `benchmarks/casereports/`; the texts are released under CC BY 4.0 with attribution to the source reports.

Ethics

MIMIC-IV is de-identified and was used under the PhysioNet credentialed data use agreement. No MIMIC text was sent to any third-party service: all models ran on local weights. Published case reports were used under their open-access licences; MIMIC-CDM-CR contains no patient data beyond what the reports published.

Data and code availability


Code, configurations, aggregate results, figures and MIMIC-CDM-CR: github.com/GIND123/Clinical-Agent-Stanford. MIMIC-IV, MIMIC-IV-Note and MIMIC-IV-Ext-CDM are available to credentialed users from PhysioNet; MIMIC-CDM-OW is rebuilt deterministically from them by the released code. Model weights trained on MIMIC will be released only through PhysioNet's credentialed channel.

References


1. Hager P, Jungmann F, Holland R, et al. Evaluation and mitigation of the limitations of large language models in clinical decision-making. *Nat Med* 2024;30:2613–2622. doi:10.1038/s41591-024-03097-1
2. US Food and Drug Administration. Clinical Decision Support Software: guidance for industry and FDA staff. Revised January 2026.

3. Barrit S, Salvagno M, Corriero A, et al. Subversion of clinical judgment by conversational artificial intelligence. *medRxiv* 2026. doi:10.64898/2026.10.05.26364541
4. Shen et al. LDTL. arXiv:2604.05116, 2026.
5. Che et al. Abstention as an action can kill both the reward gradient and the KL anchor. arXiv:2608.00301, 2026.
6. Pan et al. TIAR. arXiv:2605.25850, 2026.
7. Gao et al. KARL. arXiv:2604.22779, 2026.
8. Zhang et al. AWA-RL. arXiv:2607.10738, 2026.
9. Zhan et al. TrustMed-RL. arXiv:2610.04387, 2026.
10. Di Saverio S, Podda M, De Simone B, et al. Diagnosis and treatment of acute appendicitis: 2020 update of the WSES Jerusalem guidelines. *World J Emerg Surg* 2020;15:27. doi:10.1186/s13017-020-00306-3
11. Sartelli M, Weber DG, Kluger Y, et al. 2020 update of the WSES guidelines for the management of acute colonic diverticulitis in the emergency setting. *World J Emerg Surg* 2020;15:32. doi:10.1186/s13017-020-00313-4
12. Banks PA, Bollen TL, Dervenis C, et al. Classification of acute pancreatitis—2012: revision of the Atlanta classification and definitions by international consensus. *Gut* 2013;62:102–111. doi:10.1136/gutjnl-2012-302779
13. Yokoe M, Hata J, Takada T, et al. Tokyo Guidelines 2018: diagnostic criteria and severity grading of acute cholecystitis. *J Hepatobiliary Pancreat Sci* 2018;25:41–54. doi:10.1002/jhbp.515
14. Dratsch T, Chen X, Rezazade Mehrizi M, et al. Automation bias in mammography: the impact of artificial intelligence BI-RADS suggestions on reader performance. *Radiology* 2023;307:e222176. doi:10.1148/radiol.222176
15. Mozannar H, Sontag D. Consistent estimators for learning to defer to an expert. *ICML* 2020.
16. Dvijotham KD, Winkens J, Barsbey M, et al. Enhancing the reliability and accuracy of AI-enabled diagnosis via complementarity-driven deferral to clinicians. *Nat Med* 2023;29:1814–1820. doi:10.1038/s41591-023-02437-x
17. Bani-Harouni et al. LA-CDM. *ICLR* 2026. arXiv:2506.13474.
18. Goddard K, Roudsari A, Wyatt JC. Automation bias: a systematic review of frequency, effect mediators, and mitigators. *J Am Med Inform Assoc* 2012;19:121–127. doi:10.1136/amiajnl-2011-000089
19. Nori H, Daswani M, Kelly C, et al. Sequential diagnosis with language models. arXiv:2506.22405, 2025.
20. Johri S, Jeong J, Tran BA, et al. An evaluation framework for clinical use of large language models in patient interaction tasks. *Nat Med* 2025. doi:10.1038/s41591-024-03328-5
21. Madras D, Pitassi T, Zemel R. Predict responsibly: improving fairness and accuracy by learning to defer. *NeurIPS* 2018.
22. Geifman Y, El-Yaniv R. Selective classification for deep neural networks. *NeurIPS* 2017.


Extended Data




Extended Data Fig. 1 | Data and evaluation design. Two sources and patient-disjoint cohorts. MIMIC-CDM-OW, built with MIMIC-CDM's own text pipeline, supplies OTHER cases and same-pipeline in-set controls; the five time-critical OTHER groups are held out of training entirely.




Extended Data Fig. 2 | One training step. Multi-turn GRPO with the counterfactual escalation value: rollouts, forced continuations from deferral states, group-relative advantages for every turn and the escalation credit for DEFER turns, and the PI controller on the coverage multiplier.




Extended Data Fig. 3 | Risk–coverage on CDM val + test (480 cases, 3 seeds). a, Accuracy on answered in-set cases as the stated-probability threshold is lowered; each curve ends (filled marker) at the system's own coverage. Hollow markers: forced-choice systems with a post-hoc threshold cross-fitted at the reference coverage (vertical line), ties at the cut broken to reach it exactly. b, Area under the risk–coverage curve.




Extended Data Fig. 4 | Paired differences against each comparator on the same cases. Circles: the first DEFER-Dx seed; diamonds (first two panels): the second seed, against the control thresholded at that seed's own coverage. Positive values favour DEFER-Dx (sign flipped for lower-is-better metrics, ↓); filled points have an interval that excludes 0.




Extended Data Fig. 5 | Clinical safety on CDM val + test: unflagged errors, confident errors (wrong commitments stated with p > 0.8), severity-weighted error from the time-to-harm matrix, and tests ordered per case.




Extended Data Fig. 6 | Escalation by severity. a, Share of complicated (filled) and uncomplicated (hollow) in-set cases escalated; b, unflagged errors per 100 cases. Complicated disease defined from admission ICD codes and laboratory values by the guideline for each condition (Methods). 95% case-level bootstrap intervals.




Extended Data Fig. 7 | The escalation-value estimator. a, Share of in-set deferral states whose credit has the same sign with K forced continuations as with K = 16 (2,000 random subsets per state). b, Share of in-set deferrals credited when the escalation value assumes a clinician of the given accuracy; vertical lines: τ = 0.85 used in training, and 0.89, the MIMIC-CDM physicians.




Extended Data Fig. 8 | Ablations at 100 training steps: DEFER-Dx's step-100 checkpoint against the same training with a constant deferral reward, without the scored handoff (η = 0), without the test-cost term, without the coverage cap, and with the handoff score weighted 1.0. Two evaluation seeds; 95% intervals.




Extended Data Fig. 9 | Training dynamics and the coverage controller: in-set deferral against the 30% cap, the coverage multiplier ν, and development-set accuracy and coverage, for plain dual ascent (first run; ν wound up and limit-cycled) and the PI controller.



Extended Data Fig. 10 | Reliability of stated probabilities on committed CDM val + test cases (bins with at least 10 commitments).



Extended Data Fig. 11 | Accuracy at full coverage by condition, CDM val + test; a deferral counts through the top of its handed-over differential.



Extended Data Fig. 12 | Never-seen time-critical OTHER groups: wrong commitments and deferral by diagnosis group (groups with at least 10 cases; ruptured aneurysm and ectopic pregnancy pooled).

Extended Data Table 1 | Cohorts. RL training 2,404 (1,760 CDM, 279 OTHER, 365 controls); development 212; CDM evaluation 480 (val 240, test 240; 51 diverticulitis); OTHER from groups seen in training 206; OTHER from five time-critical groups held out of training 187; same-pipeline in-set controls 258; MIMIC-CDM-CR 32. All cohorts are patient-disjoint.

Extended Data Table 2 | Escalation equity by subgroup. Escalation, accuracy on answered cases and unflagged errors per 100 cases by sex, age group, race, insurance and language, crude and label-adjusted, with 95% intervals, for both DEFER-Dx seeds, the consensus reward, the control and prompted zero-shot deferral (docs/fairness/; cells with fewer than 10 cases suppressed).

Supplementary Data | MIMIC-CDM-CR. The 32 cases, their source reports (authors, journal, DOI, PMID, PMCID, licence) and the construction protocol (benchmarks/casereports/).