## 2.11 (a) F(A,B,C)=(A+B')C'+A'C

7	<u>ው</u>	ហ	4	W	Ŋ	₩	0	
<u>,                                    </u>	<u></u>	Ē	<u>_</u>	0	0	0	0	D
$\vdash$	<u>-</u> -	0						
<u>_</u>	0	-	o	<u> </u>	0	-	0	C
0	0	0	0	<b>j</b> —	۳	ш	ш	Ą
0	0	L	H	0	0	H	<b>J-4</b>	œ
0	<b> -</b>	0	<b>J</b>	0	<b>–</b>	0	<u></u>	ဂ
<b>-</b>	۳-	۲	<b>j</b>	0	0	<u> </u>	H	A+B
0	P	0	)	C	0	0	101	(A+B')C'
C	0	0	0	F		) H	0	A'C
C	) <b> </b>	. 0	, ,	F	۰	) <b> </b>	٠,	π

 $F = \sum_{m} (0, 1, 3, 4, 6)$ 

## (b) F(X,Y,Z)=(X+Y')(X'+Z)(Z+Y')

7	<b>O</b> ^	ຜ	4	U	N	-	0	
-	↦	<del> </del>	_	0	0	0	0	×
<b>-</b>	-	0	0	<u> </u>	<b>,_</b> _	0	0	≺
-	0	-	0	$\vdash$	0	<u> </u>	0	7
0	0	0	0	<u> </u>	<u> </u>	_	<b> </b>	×
0	0	۳	<u></u>	0	0	-	<b>J</b>	۲,
<u> </u>	<b>1</b> -	1	Þ	0	0	-	<b>j</b>	χ+Υ,
F	0	-	0	Н	Н	<b>—</b>	<b></b>	X + Z
H	. 0	<u> </u>	1	-	0	) J-	. р-	7+Y
þ	. C	, <sub> -</sub>	0	C	· C	H		П

 $F = \sum_{m} (0, 1, 5, 7)$ 

- <u>@</u>@  $F(P,Q,R) = \sum m(2,3,4,6,7)$   $F(A,B,C,D) = \sum m(0,4,5,6,8,11,12,13,14)$
- (e) F(W,X,Y,Z)=WZ'+(W'+X')YZ'+W'Z'X'

	15	14	13	12	11	10	<b>v</b>	œ	7	0	UI	4	W	N	<b>)_</b>	0		(
	<u>⊢</u>	<u></u>	1 1	1												0		
	<b> </b>	_														0		
•	10	0					_	0	_		_	_		_		0	Σ	1
	0	0	0	0	<b>-</b>	Ь	<b>-</b>	<b>)</b>	0	0	0	0	-	<b>}</b>	<b>-</b>	⊢	×	
	0	_	0	<b>)</b>	0	H	0	<b>ب</b>	0	-	0	-	0	<b>}</b>	0	_	7,	
-	0	0	0	0	<u> </u>	-	<u> </u>	<b>.</b> ,	- <b>-</b>	<u> </u>	-	_		·	-	, <u></u>	w'+x'	
•	0	0	0	C	0	- 1-	. 0	0	0	· -	. 0	· c	) C	) <b> -</b> -		0	(W'+X')YZ'	
	-			) <b> </b>	. 0	) H		) h		) (	) (	) C	0 0	0 0	) (	) C	WZ,	
•	c	0 0	) C	0 0	) (	0 0	) (	0 0	0 0	) C	0 0	) C	0 0	) F	- 0	<b>&gt;</b> -	W'Z'X'	2 10 10
•	-	)  -	٠ ر	) F	٠ ر	) <b>+</b>	٠ ر	) h	٠ (	) h	٠ (	0 0	0	) <b>+</b>	- 0	) <b>-</b>	π	

 $F = \sum m(0,2,6,8,10,12,14)$ 

(f)  $F(A,B,C)=\sum_{m}(0,1,2,3,4,5,6,7)$ (g) F(A,B,C)= no minterm

2.12 (a) F(A,B,C)=(A+B')+C'+A'C

	1	0	U	1 +	) ،	N	N	<b>-</b>	0	
۳			-	-	٠ (	)	0	0	0	P
-	١	-	0	· C	) }-	- 1		0	0	œ
<u></u>	• (	0	-	C	<b>}</b>	4 (	0	H	0	ဂ
0	(	0	0	0	<b>-</b>	• 1	-	-	H	A,
0	•	0	<b>-</b>	_	C	) (	0	<b>)—</b> •	<b>j</b>	B
0	١	_	0	<b>—</b>	C	)  -		0	H	ဂ
1	٠	<u>-</u>	<u>-</u>		0		0	<b>→</b>	1	A+8 '
0	C	>	0	0	,_		0	-	0	A C
-	1-	•	<b>—</b>	_	,_	1-		<u> </u>	ш	η

F= No maxterm

- (b) F(X,Y,Z)=(X+Y')(X'+Z)+ZY'
  See the truth table of problem 2.10(2)
  F= MM(2,3,4,6)
- (c) F(P,Q,R,S)=(P+Q')R'+P'S'R'+PQ 2'(S'+R'+Q')

F	Ü	4	13	12	11	10	9	00	7	0	G	4	U	N	_	0	
∄	-	-	بر .	_		_	<u>,                                    </u>	ь	0	0	0	0	0	0	0	0	10
1(2	_	<u>,                                    </u>	<u> </u>	<u>-</u>	0	0	0	0	<b>-</b>	<b> </b>	<u> </u>	_	0	0	0	0	o
M N	_	<u> </u>	0	0	$\vdash$	<u>-</u>	0	0	<b>-</b>	<u>,</u>	0	0	<u>-</u>	$\vdash$	0	0	20
	1	0	<b>,_</b>	0	<u>-</u>	0	$\vdash$	0	-	0	-	0	<b>-</b>	0	-	0	တ
5.6	0	0	0	0	0	0	0	0	1	<b>)</b> 4	-	Н	₩.	<b>}</b>	<u> </u>	<b>-</b>	Ď
.7.14	0	0	0	0	<u> </u>	<u> </u>	<u> </u>	<b></b>	0	0	0	0	<b>-</b>	<u> </u>	<u> </u>	_	Ď
14	0	0	<u></u>	<u></u>	0	0	-	<b> </b>	0	0	<u>-</u>	j	0	0	<b>)</b>	۳	χ
15)	0	<b>J</b>	0	-	0	Ľ	0	<b>—</b>	0	<u>_</u>	0	<del> </del>	0	<u></u> -	0	1	ທ໌
	0	0	1	1 1	0	0	<u>-</u>	1	0	0	0	0	0	0	<b>,</b>	1	(P+Q')R'
	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	ָס'ט'ק אינט'ק
															1		PO,
	0	0	0	0 1	<b>-</b> 1	-	۱ ـــ	<b>⊢</b> (	0 (	0 (	0	0	0 (	0 (	0	0	(S'+R'+
-	_	_	<u> </u>	—— 						_			-			1	בן ינ <u>ב</u>
	-	_					,	`	_ `	- '		•	_ `			1	- 4

U, J, O, /, 14, 13)

- (d) F(A,B,C,D)=TM(2,3,4,6,8,9,10,12,13) (e) F(A,B,C)=No maxterm (f) F(A,B,C)=TM(0,1,2,3,4,5,6,7)

- 2.13 (a) XY'+X'Z+Y'Z=X'Y+X'Z

  See the truth table of problem 2.4(1)

  \( \sum\_{m(1,3,4,5)} \) NOT EQUAL TO \( \sum\_{m(1,2,3)} \)

  ----> False
- (b) (B'+C)(B'+D)=B'+CD See the truth table of problem 2.4(2)  $\sum m(0,1,2,3,7) = \sum m(0,1,2,3,7)$  ---> True

32