ī

₫DØV

<u>8</u>V

<u>aa</u> ⊲




Figure 5-72 Solution to Example 5-26 illustrating the wraparound feature.

Simplify the following equation using the Karnaugh mapping procedure:

$$X = \underline{B}(CD + \underline{C}) + C\underline{D}(\underline{V} + \underline{B} + \underline{V}\underline{B})$$

$$(av + a + v)a + (c + a + va) = \lambda$$

$$X = \underline{B}CD + \underline{B}\underline{C} + \underline{V}\underline{D}(\underline{V}\underline{B} + VB)$$

$$X = \underline{B}CD + \underline{B}\underline{C} + \underline{V}\underline{D}(\underline{V}\underline{B} + VB)$$

formed:

term in the final equation. encirclement (4-variable)

Students often neglect to

Misconception Common

əfgniz ədt əbuləni

 $X = \underline{AB} + \underline{BC} + \underline{BD} + \underline{ABCD}$

unattached I in the bottom row can be combined within a group of four, form \overline{BC} . That leaves two 1's that are not combined with any others. The Figure 5-73. Another group of four can be encircled using wraparound to The group of four 1's can be encircled to form $\overline{A}\overline{B}$, as shown in

The last I is not adjacent to any other, so it must be encircled by it-

Solution: Before filling in the K-map, an SOP expression must be

EXVMBLE 5-27

EXYMPLE 5-28

<u>₽</u>Ω

as shown, to form BD.

Figure 5-73 Solution to Example 5-27.

 $\underline{g}V$

<u>₽</u>

<u>av</u>

self to form $ABC\overline{D}$. The final simplified equation is