

Dataset Description

ExoHabitAI: Exoplanet Archive - TESS Mission

1. Overview

This dataset contains comprehensive scientific observations of exoplanets and their host stars from NASA's TESS (Transiting Exoplanet Survey Satellite) mission. Each record represents a confirmed or candidate exoplanet with detailed measurements of planetary properties, stellar characteristics, orbital parameters, and detection metadata.

Dataset Specifications:

- **Format:** CSV (Comma-Separated Values)
 - **Records:** ~39,000 exoplanet observations
 - **Features:** ~289 columns
 - **Granularity:** One row = one exoplanet
 - **Source:** <https://exoplanetarchive.ipac.caltech.edu/>
-

2. Feature Categories

a) Planetary Identifiers

Column	Description	Example
pl_name	Official planet name	TOI-700 d
hostname	Host star name	TOI-700
tic_id	TESS catalog ID	150428135
pl_letter	Planet designation in system	d
gaia_id	Gaia catalog ID	Gaia DR2 123456789
disc_year	Discovery year	2020

disc_facility	Discovery facility	TESS
----------------------	--------------------	------

b) Planetary Physical Properties

Column	Description	Units	Usage
pl_rade	Planet radius	Earth radii (R_{\oplus})	Size classification
pl_bmasse	Planet mass	Earth masses (M_{\oplus})	Type determination
pl_dens	Bulk density	g/cm ³	Composition inference
pl_eqt	Equilibrium temperature	Kelvin	Habitability assessment
pl_insol	Stellar flux received	Earth flux	Energy input
pl_orbeccen	Orbital eccentricity	0.0 - 1.0	Orbit stability

Classification Guidelines:

- **Rocky planets:** $R < 1.5 R_{\oplus}$, density $> 4 \text{ g/cm}^3$
- **Super-Earths:** $1.5 < R < 2.5 R_{\oplus}$
- **Neptune-like:** $2.5 < R < 6 R_{\oplus}$, low density
- **Gas giants:** $R > 6 R_{\oplus}$ or $M > 50 M_{\oplus}$
- **Potentially habitable:** $250 \text{ K} < \text{Temperature} < 350 \text{ K}$, $0.3 < \text{Flux} < 1.5$

c) Orbital Parameters

Column	Description	Units	Significance
pl_orbper	Orbital period	Days	Orbit duration
pl_orbsmax	Semi-major axis	AU	Star-planet distance
pl_trandep	Transit depth	ppm	Detection signal strength
pl_trandur	Transit duration	Hours	Transit geometry
pl_orbincl	Orbital inclination	Degrees	Viewing angle
pl_imppar	Impact parameter	Stellar radii	Transit path

Key Insights:

- Transit detection requires inclination $\geq 85^\circ$

- Shorter periods = closer, hotter planets
- Semi-major axis determines received stellar flux

d) Host Star Properties

Column	Description	Units	Classification Use
<code>st_teff</code>	Effective temperature	Kelvin	Spectral type
<code>st_rad</code>	Stellar radius	Solar radii (R_{\odot})	Star size
<code>st_mass</code>	Stellar mass	Solar masses (M_{\odot})	Evolutionary stage
<code>st_lum</code>	Luminosity	Solar luminosity	Energy output
<code>st_met</code>	Metallicity [Fe/H]	Dex	Chemical composition
<code>st_logg</code>	Surface gravity	$\log_{10}(\text{cm/s}^2)$	Stellar density

Stellar Types:

- **M-dwarf:** Teff < 4000 K (most common exoplanet hosts)
- **K-type:** Teff 4000-5200 K
- **G-type (Sun-like):** Teff 5200-6000 K
- **F-type:** Teff 6000-7500 K

e) Detection Metadata

Column	Description	Values
<code>disposition</code>	Confirmation status	"Confirmed" / "Candidate"
<code>disposition_source</code>	Confirming authority	Institution/Survey
<code>pl_refname</code>	Reference publication	ADS bibcode
<code>rv_flag</code>	Radial velocity detection	0/1
<code>tran_flag</code>	Transit detection	0/1
<code>ttv_flag</code>	Transit timing variation	0/1

Quality Indicators:

- **Confirmed + Multiple methods:** Highest confidence
- **Confirmed + Single method:** Medium confidence

- **Candidate:** Awaiting confirmation
-

3. Data Quality

a) Missing Data Patterns

Common Missing Values:

- **Planet mass (~60%):** Often unavailable for transit-only detections
- **Density:** Missing when mass or radius unavailable
- **Eccentricity:** Difficult to measure for long-period planets
- **Some stellar parameters:** Incomplete for faint stars

b) Data Validation

Physical Consistency Checks:

- Density should be 0.1 - 30 g/cm³
 - Temperature should correlate with stellar flux
 - Orbital period should match semi-major axis (Kepler's laws)
 - Transit depth should match radius ratio
-

4. Interpretation Caveats

- **Mass-radius degeneracy:** Similar sizes can have different compositions
 - **Atmospheric assumptions:** Equilibrium temperature assumes no atmosphere
 - **Stellar activity:** Can introduce measurement noise
 - **Multi-planet interactions:** Complex gravitational dynamics in some systems
-

5. Quick Reference

a) Planet Size Classification

Category	Radius Range	Mass Range	Example
Earth-sized	0.8-1.25 R [⊕]	0.5-2 M [⊕]	Earth, Venus

Super-Earth	1.25-2 R [⊕]	2-10 M [⊕]	Kepler-452b
Mini-Neptune	2-4 R [⊕]	10-20 M [⊕]	GJ 1214b
Neptune-like	4-6 R [⊕]	10-50 M [⊕]	Neptune
Jupiter-like	>6 R [⊕]	>50 M [⊕]	Jupiter

b) Stellar Classification

Type	Temperature (K)	Color	Common Hosts
M	<3,900	Red	Proxima Centauri
K	3,900-5,200	Orange	Epsilon Eridani
G	5,200-6,000	Yellow	Sun
F	6,000-7,500	White	Procyon

c) Habitability Criteria

- **Temperature:** 250-350 K (liquid water range)
 - **Stellar flux:** 0.3-1.5 Earth flux
 - **Radius:** 0.8-1.5 R[⊕] (rocky planets)
 - **Orbital eccentricity:** <0.3 (stable conditions)
 - **Star type:** K or G preferred (stable, long-lived)
-

6. Glossary

AU: Astronomical Unit - Earth-Sun distance (~150 million km)

Equilibrium Temperature: Expected surface temperature from energy balance

Flux: Stellar energy received per unit area

Habitable Zone: Orbital region where liquid water could exist

Inclination: Orbital tilt relative to observer's line of sight

Radial Velocity: Star's motion due to planet's gravitational pull

Semi-major Axis: Average orbital distance

Transit: Planet passing in front of its star

Transit Depth: Brightness decrease during transit

Document Version: 1.0

Last Updated: January 12, 2025

Prepared by: Dheeraj M