

Chapter 10 Equality constrained minimization

Last update on 2022-05-26 21:39

Table of contents

Equality constrained minimization

Eliminating equality constraints

Newton's method with equality constraints

Infeasible start Newton method

Implementation

Equality constrained minimization

Eliminating equality constraints

Newton's method with equality constraints

Infeasible start Newton method

Implementation

Equality constrained minimization

equality constrained minimization problem

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{subject to} & Ax = b\end{array}$$

- ▶ f convex and twice continuously differentiable
- ▶ $A \in \mathbb{R}^{p \times n}$ with **rank** $A = p$
- ▶ assume optimal value p^* is finite and attained

optimality condition (review)

$$x^* \text{ is optimal} \quad \Longleftrightarrow \quad \begin{array}{l} x^* \in \text{dom } f, \quad Ax^* = b, \\ \text{there exists } \nu^* \text{ such that } \nabla f(x^*) + A^T \nu^* = 0 \end{array}$$

equality constrained quadratic minimization (with $P \in \mathbb{S}_+^n$)

$$\begin{array}{ll} \text{minimize} & (1/2)x^T Px + q^T x + r \\ \text{subject to} & Ax = b \end{array}$$

optimality condition

$$\begin{bmatrix} P & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} x^* \\ \nu^* \end{bmatrix} = \begin{bmatrix} -q \\ b \end{bmatrix}$$

- ▶ coefficient matrix is called KKT matrix
- ▶ KKT matrix is nonsingular if and only if

$$Ax = 0, \quad x \neq 0 \quad \implies \quad x^T Px > 0$$

- ▶ equivalent condition for nonsingularity

$$P + A^T A \succ 0$$

Equality constrained minimization

Eliminating equality constraints

Newton's method with equality constraints

Infeasible start Newton method

Implementation

Eliminating equality constraints

represent solutions of $\{x \mid Ax = b\}$ as

$$\{x \mid Ax = b\} = \{Fz + \hat{x} \mid z \in \mathbb{R}^{n-p}\}$$

- ▶ \hat{x} is any particular solution
- ▶ range of $F \in \mathbb{R}^{n \times (n-p)}$ is nullspace of A

reduced or eliminated problem

$$\text{minimize} \quad f(Fz + \hat{x})$$

- ▶ unconstrained problem with variable $z \in \mathbb{R}^{n-p}$
- ▶ from solution z^* , obtain x^* and ν^* as

$$x^* = Fz^* + \hat{x}, \quad \nu^* = -(AA^T)^{-1} A \nabla f(x^*)$$

example optimal allocation with resource constraint

$$\begin{array}{ll}\text{minimize} & f_1(x_1) + \cdots + f_n(x_n) \\ \text{subject to} & x_1 + \cdots + x_n = b\end{array}$$

eliminate $x_n = b - x_1 - \cdots - x_{n-1}$, namely, choose

$$\hat{x} = be_n, \quad F = \begin{bmatrix} I \\ -\mathbf{1}^T \end{bmatrix} \in \mathbb{R}^{n \times (n-1)}$$

reduced problem

$$\text{minimize} \quad f_1(x_1) + \cdots + f_{n-1}(x_{n-1}) + f_n(b - x_1 - \cdots - x_{n-1})$$

Equality constrained minimization

Eliminating equality constraints

Newton's method with equality constraints

Infeasible start Newton method

Implementation

Newton step

Newton step Δx_{nt} of f at feasible x is given by the solution v of

$$\begin{bmatrix} \nabla^2 f(x) & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} v \\ w \end{bmatrix} = \begin{bmatrix} -\nabla f(x) \\ 0 \end{bmatrix}$$

interpretations

- ▶ Δx_{nt} solves second order approximation (with variable v)

$$\begin{array}{ll} \text{minimize} & \hat{f}(x+v) = f(x) + \nabla f(x)^T v + (1/2)v^T \nabla^2 f(x)v \\ \text{subject to} & A(x+v) = b \end{array}$$

- ▶ Δx_{nt} equations follow from linearizing optimality conditions

$$\nabla f(x+v) + A^T w \approx \nabla f(x) + \nabla^2 f(x)v + A^T w = 0, \quad A(x+v) = b$$

$$\lambda(x) = (\Delta x_{\text{nt}}^T \nabla^2 f(x) \Delta x_{\text{nt}})^{1/2} = (-\nabla f(x)^T \Delta x_{\text{nt}})^{1/2}$$

interpretations

- ▶ gives an estimate of $f(x) - p^*$ using quadratic approximation \hat{f}

$$f(x) - \inf_{Ay=b} f(y) = (1/2)\lambda(x)^2$$

- ▶ directional derivative in Newton direction

$$\left. \frac{d}{dt} f(x + t\Delta x_{\text{nt}}) \right|_{t=0} = -\lambda(x)^2$$

- ▶ in general $\lambda(x) \neq (\nabla f(x)^T \nabla^2 f(x)^{-1} \nabla f(x))^{1/2}$

Newton's method with equality constraints

given starting point $x \in \text{dom } f$ with $Ax = b$, tolerance $\epsilon > 0$

repeat

1. Compute the Newton step and decrement $\Delta x_{\text{nt}}, \lambda(x)$
 2. *Stopping criterion.* **quit** if $\lambda^2/2 \leq \epsilon$
 3. *Line search.* Choose step size t by backtracking line search
 4. *Update.* $x := x + t\Delta x_{\text{nt}}$
-

- ▶ feasible descent method: $x^{(k)}$ feasible and $f(x^{(k+1)}) < f(x^{(k)})$
- ▶ affine invariant

Newton's method for reduced problem

$$\text{minimize } \tilde{f}(z) = f(Fz + \hat{x})$$

- ▶ $z \in \mathbb{R}^{n-p}$ are variables, \hat{x} satisfies $A\hat{x} = b$, range of F is the nullspace of A
- ▶ Newton's method for \tilde{f} starts at $z^{(0)}$, generates iterates $z^{(k)}$

relation to Newton's method with equality constraints

when starting at $x^{(0)} = Fz^{(0)} + \hat{x}$, iterates are

$$x^{(k)} = Fz^{(k)} + \hat{x}$$

hence no separate convergence analysis is needed

Equality constrained minimization

Eliminating equality constraints

Newton's method with equality constraints

Infeasible start Newton method

Implementation

Newton step at infeasible points

Newton step Δx_{nt} of f at infeasible x is given by the solution of

$$\begin{bmatrix} \nabla^2 f(x) & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} \Delta x_{\text{nt}} \\ w \end{bmatrix} = - \begin{bmatrix} \nabla f(x) \\ Ax - b \end{bmatrix}$$

interpretation

- ▶ Δx_{nt} equations follow from linearizing optimality conditions

$$\nabla f(x + v) + A^T w \approx \nabla f(x) + \nabla^2 f(x)v + A^T w = 0, \quad A(x + v) = b$$

primal-dual interpretation

- ▶ write optimality condition as $r(y) = 0$ where

$$y = (x, \nu), \quad r(y) = (\nabla f(x) + A^T \nu, Ax - b)$$

- ▶ linearizing $r(y) = 0$ gives

$$r(y + \Delta y) \approx r(y) + Dr(y)\Delta y = 0$$

which is equivalent to

$$\begin{bmatrix} \nabla^2 f(x) & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} \Delta x_{\text{nt}} \\ \Delta \nu_{\text{nt}} \end{bmatrix} = - \begin{bmatrix} \nabla f(x) + A^T \nu \\ Ax - b \end{bmatrix}$$

same as the above equation with $w = \nu + \Delta \nu_{\text{nt}}$

Infeasible start Newton method

given starting point $x \in \mathbf{dom} f$, ν , tolerance $\epsilon > 0$, $\alpha \in (0, 1/2)$, $\beta \in (0, 1)$

repeat

1. Compute primal and dual Newton steps $\Delta x_{\text{nt}}, \Delta \nu_{\text{nt}}$
2. *Backtracking line search on $\|r\|_2$.* $t := 1$.
while $\|r(x + t\Delta x_{\text{nt}}, \nu + t\Delta \nu_{\text{nt}})\|_2 > (1 - \alpha t)\|r(x, \nu)\|_2$, $t := \beta t$
3. *Update.* $x := x + t\Delta x_{\text{nt}}, \nu := \nu + t\Delta \nu_{\text{nt}}$

until $Ax = b$ and $\|r(x, \nu)\|_2 \leq \epsilon$

- ▶ not a descent method: $f(x^{(k+1)}) > f(x^{(k)})$ is possible
- ▶ directional derivative of $\|r(y)\|_2$ in direction $\Delta y = (\Delta x_{\text{nt}}, \Delta \nu_{\text{nt}})$ is

$$\left. \frac{d}{dt} \|r(y + t\Delta y)\|_2 \right|_{t=0} = -\|r(y)\|_2$$

Equality constrained minimization

Eliminating equality constraints

Newton's method with equality constraints

Infeasible start Newton method

Implementation

Solving KKT systems

$$\begin{bmatrix} H & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} v \\ w \end{bmatrix} = - \begin{bmatrix} g \\ h \end{bmatrix}$$

solution methods

- ▶ LDL^T factorization
- ▶ elimination with nonsingular H

$$AH^{-1}A^Tw = h - AH^{-1}g, \quad Hv = -(g + A^Tw)$$

- ▶ elimination with singular H first write as

$$\begin{bmatrix} H + A^TQA & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} v \\ w \end{bmatrix} = - \begin{bmatrix} g + A^TQh \\ h \end{bmatrix}$$

with $Q \succeq 0$ for which $H + A^TQA \succ 0$, then apply elimination

Equality constrained analytic centering

primal problem

$$\begin{array}{ll}\text{minimize} & -\sum_{i=1}^n \log x_i \\ \text{subject to} & Ax = b\end{array}$$


dual problem

$$\text{maximize} \quad -b^T \nu + \sum_{i=1}^n \log(A^T \nu)_i + n$$

three methods for an example with $A \in \mathbb{R}^{100 \times 500}$, different starting points


Newton method with equality constraints

requires $x^{(0)} \succ 0$ and $Ax^{(0)} = b$




Newton method applied to dual problem

requires $A^T \nu^{(0)} \succ 0$



infeasible start Newton method

requires $x^{(0)} \succ 0$



dominant steps of three methods

1. use block elimination to solve KKT system

$$\begin{bmatrix} \mathbf{diag}(x)^{-2} & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} \Delta x \\ w \end{bmatrix} = \begin{bmatrix} \mathbf{diag}(x)^{-1} \mathbf{1} \\ 0 \end{bmatrix}$$

reduces to solving

$$A \mathbf{diag}(x)^2 A^T w = b$$

2. solve Newton system

$$A \mathbf{diag}(A^T \nu)^{-2} A^T \Delta \nu = -b + A \mathbf{diag}(A^T \nu)^{-1} \mathbf{1}$$

3. use block elimination to solve KKT system

$$\begin{bmatrix} \mathbf{diag}(x)^{-2} & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta \nu \end{bmatrix} = \begin{bmatrix} \mathbf{diag}(x)^{-1} \mathbf{1} - A^T \nu \\ b - Ax \end{bmatrix}$$

reduces to solving

$$A \mathbf{diag}(x)^2 A^T w = 2Ax - b$$

comparison of complexity per iteration

- ▶ in each case, solve

$$ADA^T w = h$$

with D positive diagonal

- ▶ complexity per iteration of three methods is identical

Network flow optimization

$$\begin{array}{ll}\text{minimize} & \sum_{i=1}^n \phi_i(x_i) \\ \text{subject to} & Ax = b\end{array}$$

- ▶ directed (connected) graph with n arcs and $p + 1$ nodes
- ▶ x_i is flow through arc i
- ▶ ϕ_i is cost flow function for arc i (with $\phi_i''(x) > 0$)
- ▶ A is (reduced) node-arc incidence matrix
- ▶ $b \in \mathbb{R}^p$ is (reduced) source vector

KKT system

$$\begin{bmatrix} H & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} v \\ w \end{bmatrix} = - \begin{bmatrix} g \\ h \end{bmatrix}$$

- ▶ $H = \mathbf{diag}(\phi_1''(x_1), \dots, \phi_n''(x_n))$ with positive diagonal
- ▶ solve via elimination

$$AH^{-1}A^Tw = h - AH^{-1}g, \quad Hv = -(g + A^Tw)$$

sparsity pattern of coefficient matrix is given by graph connectivity

$$\begin{aligned} (AH^{-1}A^T)_{ij} \neq 0 & \iff (AA^T)_{ij} \neq 0 \\ & \iff \text{nodes } i \text{ and } j \text{ are connected by an arc} \end{aligned}$$

Analytic center of linear matrix inequality

$$\begin{array}{ll}\text{minimize} & -\log \det X \\ \text{subject to} & \mathbf{tr}(A_i X) = b_i, \quad i = 1, \dots, p\end{array}$$

where $X \in \mathbb{S}^n$ is the variable, $A_i \in \mathbb{S}^n$, $b_i \in \mathbb{R}$

optimality conditions

$$X^* \succ 0, \quad -(X^*)^{-1} + \sum_{j=1}^p \nu_j^* A_j = 0, \quad \mathbf{tr}(A_i X^*) = b_i, \quad i = 1, \dots, p$$

Newton equation at feasible X

$$X^{-1}\Delta X X^{-1} + \sum_{j=1}^p w_j A_j = X^{-1}, \quad \mathbf{tr}(A_i \Delta X) = 0, \quad i = 1, \dots, p$$

- ▶ follows from linear approximation

$$(X + \Delta X)^{-1} \approx X^{-1} - X^{-1}\Delta X X^{-1}$$

- ▶ $n(n+1)/2 + p$ variables in ΔX and w

solution by block elimination

- ▶ compute ΔX from first equation

$$\Delta X = X - \sum_{j=1}^p w_j X A_j X$$

- ▶ substitute ΔX in second equation

$$\sum_{j=1}^p \text{tr}(A_i X A_j X) w_j = b_i, \quad i = 1, \dots, p$$

a (dense) positive definite set of linear equations with variable $w \in \mathbb{R}^p$

flop count (dominant terms) using Cholesky factorization $X = LL^T$

- ▶ form p products $L^T A_j L$: $(3/2)pn^3$
- ▶ form $p(p+1)/2$ inner products $\text{tr}((L^T A_i L)(L^T A_j L))$: $(1/2)p^2 n^2$
- ▶ solve for w_j via Cholesky factorization: $(1/3)p^3$