

Chapter 5 Duality

Last update on 2022-04-27 08:09

Table of contents

Lagrange dual problem

Weak and strong duality

Geometric interpretation

Optimality conditions

Perturbation and sensitivity analysis

Examples

Generalized inequalities

Lagrange dual problem

Weak and strong duality

Geometric interpretation

Optimality conditions

Perturbation and sensitivity analysis

Examples

Generalized inequalities

Lagrangian

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p \end{array}$$

variable $x \in \mathbb{R}^n$, domain \mathcal{D} , optimal value p^*

Lagrangian $L: \mathbb{R}^n \times \mathbb{R}^m \times \mathbb{R}^p \rightarrow \mathbb{R}$ with $\text{dom } L = \mathcal{D} \times \mathbb{R}^m \times \mathbb{R}^p$

$$L(x, \lambda, \nu) = f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) + \sum_{i=1}^p \nu_i h_i(x)$$

- ▶ weighted sum of objective and constraint functions
- ▶ λ_i and ν_i are Lagrange multipliers

Lagrange dual function

Lagrange dual function $g: \mathbb{R}^m \times \mathbb{R}^p \rightarrow \mathbb{R}$

$$g(\lambda, \nu) = \inf_{x \in \mathcal{D}} L(x, \lambda, \nu)$$

g is concave, can be $-\infty$ for some values of λ and ν

lower bound property $g(\lambda, \nu) \leq p^*$ for any $\lambda \succeq 0$

proof for any feasible \bar{x} and $\lambda \succeq 0$

$$g(\lambda, \nu) = \inf_{x \in \mathcal{D}} L(x, \lambda, \nu) \leq L(\bar{x}, \lambda, \nu) \leq f_0(\bar{x})$$

minimizing over all feasible \bar{x} gives $g(\lambda, \nu) \leq p^*$

Least-norm solution of linear equations

$$\begin{array}{ll}\text{minimize} & x^T x \\ \text{subject to} & Ax = b\end{array}$$

- ▶ Lagrangian $L(x, \nu) = x^T x + \nu^T (Ax - b)$
- ▶ to minimize L over x , set gradient equal to zero

$$\nabla_x L(x, \nu) = 2x + A^T \nu = 0 \quad \implies \quad x = -(1/2)A^T \nu$$

- ▶ dual function (concave in ν)

$$g(\nu) = L\left(\left(-1/2\right)A^T \nu, \nu\right) = -(1/4)\nu^T A A^T \nu - b^T \nu$$

- ▶ lower bound property $p^* \geq -(1/4)\nu^T A A^T \nu - b^T \nu \quad \text{for all } \nu$

Standard form LP

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \succeq 0\end{array}$$

- ▶ Lagrangian ($\lambda \succeq 0$, affine in x)

$$L(x, \lambda, \nu) = c^T x + \nu^T (Ax - b) - \lambda^T x = -b^T \nu + (c + A^T \nu - \lambda)^T x$$

- ▶ dual function (linear on affine domain hence concave)

$$g(\lambda, \nu) = \inf_x L(x, \lambda, \nu) = \begin{cases} -b^T \nu & A^T \nu - \lambda + c = 0 \\ -\infty & \text{otherwise} \end{cases}$$

- ▶ lower bound property $p^* \geq -b^T \nu$ if $A^T \nu + c \succeq 0$

Equality constrained norm minimization

$$\begin{array}{ll}\text{minimize} & \|x\| \\ \text{subject to} & Ax = b\end{array}$$

- ▶ Lagrangian $L(x, \nu) = \|x\| - \nu^T(Ax - b) = \|x\| - \nu^T Ax + b^T \nu$
- ▶ dual function

$$g(\nu) = \inf_x L(x, \nu) = \begin{cases} b^T \nu & \|A^T \nu\|_* \leq 1 \\ -\infty & \text{otherwise} \end{cases}$$

where $\|v\|_* = \sup_{\|u\| \leq 1} u^T v$ is the dual norm (proof on next page)

- ▶ lower bound property $p^* \geq b^T \nu$ if $\|A^T \nu\|_* \leq 1$

proof

observe that

$$\inf_x (\|x\| - y^T x) = \begin{cases} 0 & \|y\|_* \leq 1 \\ -\infty & \text{otherwise} \end{cases}$$

- ▶ if $\|y\|_* \leq 1$, then $y^T x \leq \|x\| \|y\|_* \leq \|x\|$ for all x , with equality if $x = 0$
- ▶ if $\|y\|_* > 1$, choose $x = tu$ such that $\|u\| \leq 1$ and $y^T u > 1$, then

$$\lim_{t \rightarrow \infty} (\|x\| - y^T x) = t (\|u\| - \|y\|_*) = -\infty$$

Two-way partitioning problem

$$\begin{array}{ll}\text{minimize} & x^T W x \\ \text{subject to} & x_i^2 = 1, \quad i = 1, \dots, n\end{array}$$

- ▶ nonconvex problem, feasible set contains 2^n discrete points
- ▶ $W \in \mathbb{S}^n$, W_{ij} is cost of assigning i and j to the same set
- ▶ interpretation: find the most harmonious way to divide $\{1, \dots, n\}$ in two sets

► Lagrangian

$$L = x^T W x + \sum_{i=1}^n \nu_i (x_i^2 - 1)$$

► dual function

$$g(\nu) = \inf_x (x^T (W + \mathbf{diag}(\nu))x - \mathbf{1}^T \nu) = \begin{cases} -\mathbf{1}^T \nu & W + \mathbf{diag}(\nu) \succeq 0 \\ -\infty & \text{otherwise} \end{cases}$$

► lower bound property

$$p^* \geq -\mathbf{1}^T \nu \quad \text{if } W + \mathbf{diag}(\nu) \succeq 0$$

► example

$$\nu = -\lambda_{\min}(W)\mathbf{1} \quad \text{gives bound } p^* \geq n\lambda_{\min}(W)$$

Lagrange dual & conjugate function

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & Ax \preceq b \\ & Cx = d\end{array}$$

dual function

$$\begin{aligned}g(\lambda, \nu) &= \inf_{x \in \text{dom } f_0} \left(f_0(x) + (A^T \lambda + C^T \nu)^T x - b^T \lambda - d^T \nu \right) \\ &= -f_0^* (-A^T \lambda - C^T \nu) - b^T \lambda - d^T \nu\end{aligned}$$

- ▶ recall definition of conjugate $f^*(y) = \sup_{x \in \text{dom } f} (y^T x - f(x))$
- ▶ simplifies derivation of dual if conjugate of f_0 is known

Entropy maximization

$$\begin{array}{ll}\text{minimize} & f_0(x) = \sum_{i=1}^n x_i \log x_i \\ \text{subject to} & Ax \preceq b \\ & \mathbf{1}^T x = 1\end{array}$$

- conjugate of $f_0(x)$

$$f_0^*(y) = \sum_{i=1}^n e^{y_i - 1}$$

- dual function

$$g(\lambda, \nu) = - \sum_{i=1}^n e^{-a_i^T \lambda - \nu - 1} - b^T \lambda - \nu = -e^{-\nu-1} \sum_{i=1}^n e^{-a_i^T \lambda} - b^T \lambda - \nu$$

Lagrange dual problem

$$\begin{array}{ll}\text{maximize} & g(\lambda, \nu) \\ \text{subject to} & \lambda \succeq 0\end{array}$$

- ▶ finds best lower bound on p^* , obtained from Lagrange dual function
- ▶ convex optimization problem, optimal value denoted d^*
- ▶ λ and ν are dual feasible if $\lambda \succeq 0$ and $(\lambda, \nu) \in \mathbf{dom} g$
- ▶ often simplified by making implicit constraint $(\lambda, \nu) \in \mathbf{dom} g$ explicit
- ▶ original problem is called primal problem

Standard form LP

primal problem

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \succeq 0\end{array}$$

dual problem

$$\begin{array}{ll}\text{maximize} & g(\lambda, \nu) = \begin{cases} -b^T \nu & A^T \nu - \lambda + c = 0 \\ -\infty & \text{otherwise} \end{cases} \\ \text{subject to} & \lambda \succeq 0\end{array}$$

equivalent form

$$\begin{array}{ll}\text{maximize} & -b^T \nu \\ \text{subject to} & A^T \nu + c \succeq 0\end{array}$$

Inequality form LP

primal problem

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \preceq b\end{array}$$

dual problem

$$\begin{array}{ll}\text{maximize} & g(\lambda) = \begin{cases} -b^T \lambda & A^T \lambda + c = 0 \\ -\infty & \text{otherwise} \end{cases} \\ \text{subject to} & \lambda \succeq 0\end{array}$$

equivalent form

$$\begin{array}{ll}\text{maximize} & -b^T \lambda \\ \text{subject to} & A^T \lambda + c = 0 \\ & \lambda \succeq 0\end{array}$$

Lagrange dual problem

Weak and strong duality

Geometric interpretation

Optimality conditions

Perturbation and sensitivity analysis

Examples

Generalized inequalities

$$d^* \leq p^*$$

- ▶ always holds (regardless of convexity)
- ▶ can be used to find nontrivial lower bounds for difficult problem

example

solving SDP

$$\begin{array}{ll} \text{maximize} & -\mathbf{1}^T \nu \\ \text{subject to} & W + \mathbf{diag}(\nu) \succeq 0 \end{array}$$

gives a lower bound for two-way partitioning problem

$$d^* = p^*$$

- ▶ does not hold in general
- ▶ usually holds for convex problems

constraint qualifications

- ▶ conditions that guarantee strong duality for convex problems
- ▶ there exist many types, example below

Slater's constraint qualification

strong duality holds for convex problem

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b \end{array}$$

if it is strictly feasible, namely

$$\exists x \in \mathbf{int} \mathcal{D} \quad \text{such that} \quad f_i(x) < 0, \quad i = 1, \dots, m, \quad Ax = b$$

- ▶ also guarantees that the dual optimum is attained if $p^* > -\infty$
- ▶ can replace $\mathbf{int} \mathcal{D}$ with $\mathbf{relint} \mathcal{D}$ (interior relative to affine hull)
- ▶ linear inequalities do not need to hold with strict inequality
- ▶ strong duality holds for LP unless both primal and dual are infeasible (for LP, dual of dual is primal, Slater's condition and feasibility agree)

Quadratic program

primal problem (assume $P \in \mathbb{S}_{++}^n$)

$$\begin{array}{ll}\text{minimize} & x^T P x \\ \text{subject to} & Ax \preceq b\end{array}$$

dual function

$$g(\lambda) = \inf_x (x^T P x + \lambda^T (Ax - b)) = -(1/4)\lambda^T A P^{-1} A^T \lambda - b^T \lambda$$

dual problem

$$\begin{array}{ll}\text{maximize} & -(1/4)\lambda^T A P^{-1} A^T \lambda - b^T \lambda \\ \text{subject to} & \lambda \succeq 0\end{array}$$

- ▶ by Slater's condition $p^* = d^*$ holds if primal problem is feasible
- ▶ in fact $p^* = d^*$ always holds (dual of dual is primal, dual always satisfies Slater)

A nonconvex problem with strong duality

primal problem (nonconvex if $A \not\preceq 0$)

$$\begin{array}{ll}\text{minimize} & x^T A x + 2b^T x \\ \text{subject to} & x^T x \leq 1\end{array}$$

dual problem

$$\begin{array}{ll}\text{maximize} & -b^T(A + \lambda I)^\dagger b - \lambda \\ \text{subject to} & A + \lambda I \succeq 0 \\ & b \in \mathcal{R}(A + \lambda I)\end{array}$$

equivalent SDP

$$\begin{array}{ll}\text{maximize} & -t - \lambda \\ \text{subject to} & \begin{bmatrix} A + \lambda I & b \\ b^T & t \end{bmatrix} \succeq 0\end{array}$$

strong duality holds although primal problem is nonconvex (not easy to show)

Lagrange dual problem

Weak and strong duality

Geometric interpretation

Optimality conditions

Perturbation and sensitivity analysis



Examples

Generalized inequalities

Geometric interpretation

interpretation of dual function consider problem with one constraint $f_1(x) \leq 0$


$$g(\lambda) = \inf_{(u,t) \in \mathcal{G}} (t + \lambda u) \quad \text{where} \quad \mathcal{G} = \{(u,t) \mid u = f_1(x), t = f_0(x) \text{ for some } x \in \mathcal{D}\}$$



$\lambda u + t = g(\lambda)$ is (non-vertical) supporting hyperplane to \mathcal{G} meeting t -axis at $t = g(\lambda)$

epigraph variation same interpretation if \mathcal{G} is replaced by

$$\mathcal{A} = \{(u, t) \mid u \geq f_1(x), t \geq f_0(x) \text{ for some } x \in \mathcal{D}\}$$



proof of strong duality (under Slater's condition)

- ▶ holds if there is a non-vertical supporting hyperplane H to \mathcal{A} at $(0, p^*)$
- ▶ for convex problems, \mathcal{A} is convex, hence H exists
- ▶ Slater's condition guarantees the existence of $(\tilde{u}, \tilde{t}) \in \mathcal{A}$ with $\tilde{u} < 0$
- ▶ it follows that H cannot be vertical

Lagrange dual problem

Weak and strong duality

Geometric interpretation

Optimality conditions

Perturbation and sensitivity analysis

Examples

Generalized inequalities

Complementary slackness

assume x^* is primal optimal, (λ^*, ν^*) is dual optimal

$$\begin{aligned} g(\lambda^*, \nu^*) &= \inf_x \left(f_0(x) + \sum_{i=1}^m \lambda_i^* f_i(x) + \sum_{i=1}^p \nu_i^* h_i(x) \right) \\ &\leq f_0(x^*) + \sum_{i=1}^m \lambda_i^* f_i(x^*) + \sum_{i=1}^p \nu_i^* h_i(x^*) \leq f_0(x^*) \end{aligned}$$

assume strong duality holds, then both inequalities hold with equality

- ▶ x^* minimizes $L(x, \lambda^*, \nu^*)$
- ▶ $\lambda_i^* f_i(x^*) = 0$ for each $i = 1, \dots, m$, namely, for each pair of inequalities

$$\lambda_i^* \geq 0 \quad \text{and} \quad f_i(x^*) \leq 0$$

at least one of them achieves equality (complementary slackness)

assume f_0, f_1, \dots, f_m and h_1, \dots, h_p are all differentiable (hence with open domains)

Karush-Kuhn-Tucker conditions

1. primal constraints $f_i(x) \leq 0, i = 1, \dots, m; \quad h_i(x) = 0, i = 1, \dots, p$
2. dual constraints $\lambda \succeq 0$
3. complementary slackness $\lambda_i f_i(x) = 0, i = 1, \dots, m$
4. gradient of Lagrangian with respect to x vanishes

$$\nabla f_0(x) + \sum_{i=1}^m \lambda_i \nabla f_i(x) + \sum_{i=1}^p \nu_i \nabla h_i(x) = 0$$

necessity if strong duality holds

$$(\tilde{x}, \tilde{\lambda}, \tilde{\nu}) \text{ are optimal} \quad \implies \quad (\tilde{x}, \tilde{\lambda}, \tilde{\nu}) \text{ satisfy KKT}$$

sufficiency if primal problem is convex

$$(\tilde{x}, \tilde{\lambda}, \tilde{\nu}) \text{ satisfy KKT} \quad \implies \quad (\tilde{x}, \tilde{\lambda}, \tilde{\nu}) \text{ are optimal}$$

proof

- ▶ conditions 1 & 2 imply primal and dual feasibility
- ▶ condition 3 implies $f_0(\tilde{x}) = L(\tilde{x}, \tilde{\lambda}, \tilde{\nu})$
- ▶ condition 4 and convexity imply $g(\tilde{\lambda}, \tilde{\nu}) = L(\tilde{x}, \tilde{\lambda}, \tilde{\nu})$

necessity + sufficiency if Slater's condition holds for convex problem

$$\tilde{x} \text{ is optimal} \quad \iff \quad (\tilde{x}, \tilde{\lambda}, \tilde{\nu}) \text{ satisfy KKT for some } \tilde{\lambda} \text{ and } \tilde{\nu}$$

Example

assume $\alpha_i > 0$ for $i = 1, \dots, n$

$$\begin{array}{ll}\text{minimize} & -\sum_{i=1}^n \log(x_i + \alpha_i) \\ \text{subject to} & x \succeq 0 \\ & \mathbf{1}^T x = 1\end{array}$$

x is optimal $\iff x \succeq 0, \mathbf{1}^T x = 1$, there exists $\lambda \in \mathbb{R}^n$ and $\nu \in \mathbb{R}$ such that

$$\lambda \succeq 0, \quad \lambda_i x_i = 0, \quad \frac{1}{x_i + \alpha_i} + \lambda_i = \nu, \quad i = 1, \dots, n$$


- ▶ if $\nu \leq 1/\alpha_i$, then $\lambda_i = 0$ and $x_i = 1/\nu - \alpha_i$
- ▶ if $\nu \geq 1/\alpha_i$, then $\lambda_i = \nu - 1/\alpha_i$ and $x_i = 0$

determine ν from

$$\mathbf{1}^T x = \sum_{i=1}^n \max\{0, 1/\nu - \alpha_i\} = 1$$

water-filling algorithm

- ▶ left-hand side is a piecewise linear increasing function in $1/\nu$
- ▶ n patches, level of patch i is at height α_i
- ▶ flood area with unit amount of water, resulting level is $1/\nu^*$



Lagrange dual problem

Weak and strong duality

Geometric interpretation

Optimality conditions

Perturbation and sensitivity analysis

Examples

Generalized inequalities

Perturbed problem

perturbed primal problem

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq u_i, \quad i = 1, \dots, m \\ & h_i(x) = v_i, \quad i = 1, \dots, p \end{array}$$

perturbed dual problem

$$\begin{array}{ll} \text{maximize} & g(\lambda, \nu) - u^T \lambda - v^T \nu \\ \text{subject to} & \lambda \succeq 0 \end{array}$$

- ▶ u and v are parameters
- ▶ original primal & dual problems are recovered when $u = 0$ and $v = 0$
- ▶ $p^*(u, v)$ is optimal value as a function of u and v
- ▶ need to understand $p^*(u, v)$ from solution to unperturbed problem

Global sensitivity

assume for the unperturbed problem that

- ▶ strong duality holds (e.g. convex + Slater)
- ▶ λ^* and ν^* are dual optimal

then weak duality for the perturbed problem implies


$$\begin{aligned} p^*(u, v) &\geq g(\lambda^*, \nu^*) - u^T \lambda^* - v^T \nu^* \\ &= p^*(0, 0) - u^T \lambda^* - v^T \nu^* \end{aligned}$$

- ▶ λ_i^* large $\implies p^*$ increases greatly if $u_i < 0$ (tighten constraint)
- ▶ λ_i^* small $\implies p^*$ does not decrease much if $u_i > 0$ (loosen constraint)
- ▶ $\nu_i^* > 0$ large $\implies p^*$ increases greatly if $v_i < 0$
- ▶ $\nu_i^* < 0$ large $\implies p^*$ increases greatly if $v_i > 0$
- ▶ $\nu_i^* > 0$ small $\implies p^*$ does not decrease much if $v_i > 0$
- ▶ $\nu_i^* < 0$ small $\implies p^*$ does not decrease much if $v_i < 0$

Local sensitivity

assume in addition that $p^*(u, v)$ is differentiable at $(0, 0)$ then

$$\lambda_i^* = -\frac{\partial p^*}{\partial u_i}(0, 0), \quad \nu_i^* = -\frac{\partial p^*}{\partial v_i}(0, 0)$$



(above picture exhibits $p^*(u)$ for a problem with one inequality constraint)

Lagrange dual problem

Weak and strong duality

Geometric interpretation

Optimality conditions

Perturbation and sensitivity analysis

Examples

Generalized inequalities

principle

- ▶ equivalent formulations of a problem can lead to very different duals
- ▶ reformulation can be useful when dual is difficult to derive or uninteresting

common reformulations

- ▶ introduce new variables and equality constraints
- ▶ make explicit constraints implicit or vice-versa
- ▶ apply an increasing function to objective or constraint functions

Introducing new variables and equality constraints

unconstrained problem

primal problem

$$\text{minimize} \quad f_0(Ax + b)$$

dual problem

$$g = \inf_x f_0(Ax + b) = p^*$$

- ▶ no dual variable, hence dual function is constant
- ▶ strong duality holds, but dual is useless

reformulated primal problem

$$\begin{array}{ll}\text{minimize} & f_0(y) \\ \text{subject to} & Ax + b - y = 0\end{array}$$

dual of reformulated problem

$$\begin{array}{ll}\text{maximize} & b^T \nu - f_0^*(\nu) \\ \text{subject to} & A^T \nu = 0\end{array}$$

it follows from

$$g(\nu) = \inf_{x,y} (f_0(y) - \nu^T y + \nu^T Ax + b^T \nu) = \begin{cases} -f_0^*(\nu) + b^T \nu & A^T \nu = 0 \\ -\infty & \text{otherwise} \end{cases}$$

norm approximation problem

$$\text{minimize} \quad \|Ax - b\|$$

reformulated problem

$$\begin{array}{ll} \text{minimize} & \|y\| \\ \text{subject to} & y = Ax - b \end{array}$$

dual of the reformulated problem

$$\begin{array}{ll} \text{maximize} & b^T \nu \\ \text{subject to} & A^T \nu = 0 \\ & \|\nu\|_* \leq 1 \end{array}$$

LP with box constraints

primal problem

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & -1 \preceq x \preceq 1\end{array}$$

dual problem

$$\begin{array}{ll}\text{maximize} & -b^T \nu - \mathbf{1}^T \lambda_1 - \mathbf{1}^T \lambda_2 \\ \text{subject to} & c + A^T \nu + \lambda_1 - \lambda_2 = 0 \\ & \lambda_1 \succeq 0, \quad \lambda_2 \succeq 0\end{array}$$

reformulated primal problem

$$\begin{array}{ll}\text{minimize} & f_0(x) = \begin{cases} c^T x & -1 \preceq x \preceq 1 \\ \infty & \text{otherwise} \end{cases} \\ \text{subject to} & Ax = b\end{array}$$

dual function

$$\begin{aligned}g(\nu) &= \inf_{-1 \preceq x \preceq 1} (c^T x + \nu^T (Ax - b)) \\ &= -b^T \nu - \|A^T \nu + c\|_1\end{aligned}$$

dual of the reformulated problem

$$\text{maximize} \quad -b^T \nu - \|A^T \nu + c\|_1$$

Lagrange dual problem

Weak and strong duality

Geometric interpretation

Optimality conditions

Perturbation and sensitivity analysis

Examples

Generalized inequalities

Problems with generalized inequalities

primal problem (proper cone $K_i \subseteq \mathbb{R}^{k_i}$ for $i = 1, \dots, m$)

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \preceq_{K_i} 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p \end{array}$$

- ▶ Lagrange multiplier for $f_i(x) \preceq_{K_i} 0$ is vector $\lambda_i \in \mathbb{R}^{k_i}$, for $h_i(x) = 0$ scalar $\nu_i \in \mathbb{R}$
- ▶ Lagrangian $L: \mathbb{R}^n \times \mathbb{R}^{k_1} \times \dots \times \mathbb{R}^{k_m} \times \mathbb{R}^p \longrightarrow \mathbb{R}$

$$L(x, \lambda_1, \dots, \lambda_m, \nu) = f_0(x) + \sum_{i=1}^m \lambda_i^T f_i(x) + \sum_{i=1}^p \nu_i h_i(x)$$

- ▶ dual function $g: \mathbb{R}^{k_1} \times \dots \times \mathbb{R}^{k_m} \times \mathbb{R}^p \longrightarrow \mathbb{R}$

$$g(\lambda_1, \dots, \lambda_m, \nu) = \inf_{x \in \mathcal{D}} L(x, \lambda_1, \dots, \lambda_m, \nu)$$

lower bound property if $\lambda_i \succeq_{K_i^*} 0$, then $g(\lambda_1, \dots, \lambda_m, \nu) \leq p^*$

proof

$$\begin{aligned} f_0(\tilde{x}) &\geq f_0(\tilde{x}) + \sum_{i=1}^m \lambda_i^T f_i(\tilde{x}) + \sum_{i=1}^p \nu_i h_i(\tilde{x}) \\ &\geq \inf_{x \in \mathcal{D}} L(x, \lambda_1, \dots, \lambda_m, \nu) \\ &= g(\lambda_1, \dots, \lambda_m, \nu) \end{aligned}$$

holds for all feasible \tilde{x} , then minimize over all such \tilde{x} to conclude

dual problem

$$\begin{array}{ll}\text{maximize} & g(\lambda_1, \dots, \lambda_m, \nu) \\ \text{subject to} & \lambda_i \succeq_{K_i^*} 0, \quad i = 1, \dots, m\end{array}$$

weak duality (always holds)

$$p^* \geq d^*$$

strong duality (holds for convex problem with constraint qualification)

$$p^* = d^*$$

Slater's condition: primal problem is strictly feasible

Semidefinite program

primal SDP (assume $F_i, G \in \mathbb{S}^k$)

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & x_1 F_1 + \cdots + x_n F_n \preceq G\end{array}$$

Lagrange multiplier

$$Z \in \mathbb{S}^k$$

Lagrangian

$$L(x, Z) = c^T x + \mathbf{tr}(Z(x_1 F_1 + \cdots + x_n F_n - G))$$

dual function

$$g(Z) = \inf_x L(x, Z) = \begin{cases} -\operatorname{tr}(GZ) & \operatorname{tr}(F_i Z) + c_i = 0 \text{ for all } i = 1, \dots, n \\ -\infty & \text{otherwise} \end{cases}$$

dual SDP

$$\begin{array}{ll} \text{maximize} & -\operatorname{tr}(GZ) \\ \text{subject to} & Z \succeq 0 \\ & \operatorname{tr}(F_i Z) + c_i = 0, \quad i = 1, \dots, n \end{array}$$

strong duality

$p^* = d^*$ holds if primal SDP is strictly feasible ($\exists x$ such that $x_1 F_1 + \dots + x_n F_n \prec G$)