

Chapter 4 Convex optimization problems

Last update on 2022-03-23 10:04

Table of contents

Optimization problems

Convex optimization

Linear optimization

Quadratic optimization

Geometric programming

Generalized inequality constraints

Vector optimization

Optimization problems

Convex optimization

Linear optimization

Quadratic optimization

Geometric programming

Generalized inequality constraints

Vector optimization

Optimization problem in standard form

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

$x \in \mathbb{R}^n$	optimization variable
$f_0: \mathbb{R}^n \rightarrow \mathbb{R}$	objective function (cost function)
$f_i: \mathbb{R}^n \rightarrow \mathbb{R}$	inequality constraint functions
$h_i: \mathbb{R}^n \rightarrow \mathbb{R}$	equality constraint functions

$$p^* = \inf \left\{ f_0(x) \mid \begin{array}{l} f_i(x) \leq 0 \text{ for } 1 \leq i \leq m \\ h_i(x) = 0 \text{ for } 1 \leq i \leq p \end{array} \right\}$$

- ▶ $p^* = \infty$ if problem is infeasible (no x satisfies the constraints)
- ▶ $p^* = -\infty$ if problem is unbounded below

(Locally) optimal points

- ▶ x is **feasible** if $x \in \text{dom } f_0$ and it satisfies the constraints
- ▶ x is **optimal** if it is feasible and $f_0(x) = p^*$; set of optimal points X_{opt}
- ▶ x is **locally optimal** if there exists $R > 0$ such that x is optimal for

$$\begin{array}{ll}\text{minimize} & f_0(z) \\ \text{subject to} & f_i(z) \leq 0, \quad i = 1, \dots, m \\ & h_i(z) = 0, \quad i = 1, \dots, p \\ & \|z - x\|_2 \leq R\end{array}$$

Examples

$f_0(x) = 1/x$	$\text{dom } f_0 = \mathbb{R}_{++}$	$p^* = 0$	no optimal point
$f_0(x) = -\log x$	$\text{dom } f_0 = \mathbb{R}_{++}$	$p^* = -\infty$	no optimal point
$f_0(x) = x \log x$	$\text{dom } f_0 = \mathbb{R}_{++}$	$p^* = -1/e$	$x = 1/e$ is optimal
$f_0(x) = x^3 - 3x$		$p^* = -\infty$	$x = 1$ is locally optimal

($n = 1$, $m = p = 0$ in the above examples)

Implicit constraints

- ▶ **explicit constraints**

$$f_i(x) \leq 0 \text{ for } 1 \leq i \leq m \quad \text{and} \quad h_i(x) = 0 \text{ for } 1 \leq i \leq p$$

- ▶ **implicit constraints**

$$x \in \mathcal{D} = \left(\bigcap_{i=0}^m \text{dom } f_i \right) \cap \left(\bigcap_{i=1}^p \text{dom } h_i \right)$$

- ▶ \mathcal{D} is called the **domain** of the problem
- ▶ problem is **unconstrained** if it has no explicit constraints ($m = p = 0$)

example

$$\text{minimize} \quad f_0(x) = - \sum_{i=1}^k \log(b_i - a_i^T x)$$

is an unconstrained problem with implicit constraints $a_i^T x < b_i$ for $1 \leq i \leq k$

Feasibility problem

$$\begin{array}{ll}\text{find} & x \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

can be considered as an optimization problem

$$\begin{array}{ll}\text{minimize} & 0 \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

- ▶ $p^* = 0$ if constraints are feasible; any feasible x is optimal
- ▶ $p^* = \infty$ if constraints are infeasible

Optimization problems

Convex optimization

Linear optimization

Quadratic optimization

Geometric programming

Generalized inequality constraints

Vector optimization

Convex optimization problem in standard form

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & a_i^T x = b_i, \quad i = 1, \dots, p\end{array}$$

- ▶ f_0, f_1, \dots, f_m are convex
- ▶ equality constraints are affine, often written as $Ax = b$
- ▶ important property: feasible set of a convex problem is convex
- ▶ problem is **quasiconvex** if f_0 is quasiconvex (and f_1, \dots, f_m convex)

Example

$$\begin{array}{ll}\text{minimize} & f_0(x) = x_1^2 + x_2^2 \\ \text{subject to} & f_1(x) = x_1/(1 + x_2^2) \leq 0 \\ & h_1(x) = (x_1 + x_2)^2 = 0\end{array}$$

- ▶ f_0 is convex
- ▶ feasible set $\{(x_1, x_2) \mid x_1 = -x_2 \leq 0\}$ is convex
- ▶ not a convex problem: f_1 is not convex, h_1 is not affine
- ▶ equivalent (but not identical) to the convex problem

$$\begin{array}{ll}\text{minimize} & x_1^2 + x_2^2 \\ \text{subject to} & x_1 \leq 0 \\ & x_1 + x_2 = 0\end{array}$$

any locally optimal point of a convex optimization problem is globally optimal


proof

- ▶ suppose x is locally optimal, but there exists feasible y with $f_0(y) < f_0(x)$
- ▶ there exists $R > 0$ such that $f_0(z) \geq f_0(x)$ for all feasible z with $\|z - x\|_2 < R$
- ▶ consider $z = \theta y + (1 - \theta)x$ with $\theta = R/(2\|y - x\|_2)$, then $\|z - x\|_2 = R/2$
- ▶ $\|y - x\|_2 > R$ implies $0 < \theta < 1/2$, hence z is feasible by convexity of domain
- ▶ by convexity of objective $f_0(z) \leq \theta f_0(y) + (1 - \theta)f_0(x) < f_0(x)$, contradiction

Optimality criterion for differentiable objective

suppose f_0 is differentiable, then

$$x \text{ is optimal} \iff \begin{array}{l} x \text{ is feasible and} \\ \nabla f_0(x)^T (y - x) \geq 0 \text{ for all feasible } y \end{array}$$



either $\nabla f_0(x) = 0$ or it defines a supporting hyperplane to feasible set X at x

unconstrained problem

minimize $f_0(x)$

x is optimal $\iff x \in \text{dom } f_0, \nabla f_0(x) = 0$

equality constrained problem

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & Ax = b\end{array}$$

$$x \text{ is optimal} \quad \Longleftrightarrow \quad \begin{array}{l} x \in \text{dom } f_0, \quad Ax = b, \\ \text{there exists } \nu \text{ such that } \nabla f_0(x) + A^T \nu = 0 \end{array}$$

minimization over nonnegative orthant

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & x \succeq 0\end{array}$$

$$x \text{ is optimal} \iff x \in \mathbf{dom} f_0, \quad x \succeq 0, \quad \begin{cases} \nabla f_0(x)_i \geq 0, & \text{if } x_i = 0 \\ \nabla f_0(x)_i = 0, & \text{if } x_i > 0 \end{cases}$$

Equivalent convex problem

two problems are (informally) **equivalent** if the solution of one is readily obtained from the solution of the other, and vice-versa

some common transformations that preserve convexity

- ▶ eliminating equality constraints
- ▶ introducing equality constraints
- ▶ introducing slack variables for linear inequalities
- ▶ epigraph form
- ▶ minimizing over some variables

eliminating equality constraints

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

is equivalent to

$$\begin{array}{ll}\text{minimize} & f_0(Fz + x_0) \quad (\text{over } z) \\ \text{subject to} & f_i(Fz + x_0) \leq 0, \quad i = 1, \dots, m\end{array}$$

where F and x_0 are such that

$$Ax = b \iff x = Fz + x_0 \text{ for some } z$$

introducing equality constraints

$$\begin{array}{ll}\text{minimize} & f_0(A_0x + b_0) \\ \text{subject to} & f_i(A_ix + b_i) \leq 0, \quad i = 1, \dots, m\end{array}$$

is equivalent to

$$\begin{array}{lll}\text{minimize} & f_0(y_0) & (\text{over } x, y_i) \\ \text{subject to} & f_i(y_i) \leq 0, & i = 1, \dots, m \\ & y_i = A_ix + b_i, & i = 0, 1, \dots, m\end{array}$$

introducing slack variables for linear inequalities

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & a_i^T x \leq b_i, \quad i = 1, \dots, m\end{array}$$

is equivalent to

$$\begin{array}{lll}\text{minimize} & f_0(x) & (\text{over } x, z) \\ \text{subject to} & a_i^T x + s_i = b_i, & i = 1, \dots, m \\ & s_i \geq 0, & i = 1, \dots, m\end{array}$$

epigraph form

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

is equivalent to

$$\begin{array}{ll}\text{minimize} & t \quad (\text{over } x, t) \\ \text{subject to} & f_0(x) - t \leq 0 \\ & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

minimizing over some variables

$$\begin{array}{ll}\text{minimize} & f_0(x_1, x_2) \\ \text{subject to} & f_i(x_1) \leq 0, \quad i = 1, \dots, m\end{array}$$

is equivalent to

$$\begin{array}{ll}\text{minimize} & \tilde{f}_0(x_1) \\ \text{subject to} & f_i(x_1) \leq 0, \quad i = 1, \dots, m\end{array}$$

where


$$\tilde{f}_0(x_1) = \inf_{x_2} f_0(x_1, x_2)$$

Quasiconvex optimization

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

with $f_0: \mathbb{R}^n \rightarrow \mathbb{R}$ quasiconvex, f_1, \dots, f_m convex

locally optimal points may not be globally optimal



convex representation of sublevel sets of f_0

for quasiconvex f_0 there exists a family of functions ϕ_t such that

- ▶ $\phi_t(x)$ is convex in x for each fixed t
- ▶ t -sublevel set of f_0 is 0-sublevel set of ϕ_t , namely $f_0(x) \leq t \iff \phi_t(x) \leq 0$
- ▶ $\phi_t(x)$ is nonincreasing in t for each fixed x , namely $\phi_s(x) \leq \phi_t(x)$ if $s \geq t$

example

$$f_0(x) = \frac{p(x)}{q(x)}$$

with p convex, q concave, and $p(x) \geq 0$, $q(x) > 0$ on $\text{dom } f_0$

we can choose

$$\phi_t(x) = p(x) - tq(x)$$

- ▶ $\phi_t(x)$ convex in x for $t \geq 0$
- ▶ $p(x)/q(x) \leq t \iff \phi_t(x) \leq 0$

quasiconvex optimization via convex feasibility problems

$$\phi_t(x) \leq 0, \quad f_i(x) \leq 0, \quad i = 1, \dots, m, \quad Ax = b$$

- ▶ convex feasibility problem in x for each fixed t
- ▶ if feasible, then $t \geq p^*$; if infeasible, then $t \leq p^*$

bisection method

given $l \leq p^*, u \geq p^*, \text{ tolerance } \epsilon > 0$

repeat

1. $t := (l + u)/2$
2. solve the above convex feasibility problem
3. **if** feasible, $u := t$; **else** $l := t$

until $u - l \leq \epsilon$

requires exactly $\lceil \log_2((u - l)/\epsilon) \rceil$ iterations

Optimization problems

Convex optimization

Linear optimization

Quadratic optimization

Geometric programming


Generalized inequality constraints

Vector optimization

Linear program (LP)

$$\begin{array}{ll}\text{minimize} & c^T x + d \\ \text{subject to} & Gx \preceq h \\ & Ax = b\end{array}$$

- ▶ convex problem with affine objective and constraint functions
- ▶ feasible set is a polyhedron



Examples

diet problem: choose quantities x_1, \dots, x_n of n kinds of food

- ▶ one unit of food j costs c_j , contains amount a_{ij} of nutrient i
- ▶ healthy diet requires nutrient i in quantity at least b_i

to find cheapest healthy diet

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \succeq b \\ & x \succeq 0\end{array}$$

piecewise-linear minimization

$$\text{minimize} \quad \max \{a_i^T x + b_i \mid i = 1, \dots, m\}$$

equivalent to the LP

$$\begin{array}{ll} \text{minimize} & t \\ \text{subject to} & a_i^T x + b_i \leq t, \quad i = 1, \dots, m \end{array}$$


Chebyshev center of a polyhedron

Chebyshev center of

$$\mathcal{P} = \{x \mid a_i^T x \leq b_i, \quad i = 1, \dots, m\}$$

is center of largest inscribed ball

$$\mathcal{B} = \{x_c + u \mid \|u\|_2 \leq r\}$$



$a_i^T x \leq b_i$ for all $x \in \mathcal{B}$ if and only if

$$\sup\{a_i^T(x_c + u) \mid \|u\|_2 \leq r\} = a_i^T x_c + r\|a_i\|_2 \leq b_i$$

hence x_c and r can be determined by solving the LP

$$\begin{array}{ll} \text{maximize} & r \\ \text{subject to} & a_i^T x_c + r\|a_i\|_2 \leq b_i, \quad i = 1, \dots, m \end{array}$$

Linear-fractional program

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & Gx \preceq h \\ & Ax = b\end{array}$$

where

$$f_0(x) = \frac{c^T x + d}{e^T x + f}, \quad \text{dom } f_0(x) = \{x \mid e^T x + f > 0\}$$

- ▶ a quasiconvex optimization problem; can be solved by bisection
- ▶ also equivalent to the LP

$$\begin{array}{ll}\text{minimize} & c^T y + dz \\ \text{subject to} & Gy \preceq hz \\ & Ay = bz \\ & e^T y + fz = 1 \\ & z \geq 0\end{array}$$

Generalized linear-fractional program

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & Gx \preceq h \\ & Ax = b\end{array}$$

where

$$f_0(x) = \max \left\{ \frac{c_i^T x + d_i}{e_i^T x + f_i} \mid i = 1, \dots, r \right\}$$
$$\text{dom } f_0(x) = \left\{ x \mid e_i^T x + f_i > 0, \ i = 1, \dots, r \right\}$$

a quasiconvex optimization problem; can be solved by bisection

Example

Von Neumann model of a growing economy

$$\begin{array}{ll} \text{maximize} & \min \{x_i^+ / x_i \mid i = 1, \dots, n\} \quad (\text{over } x, x^+) \\ \text{subject to} & x^+ \succeq 0 \\ & Bx^+ \preceq Ax \end{array}$$

with domain $\{(x, x^+) \mid x \succ 0\}$

- ▶ $x, x^+ \in \mathbb{R}^n$: activity levels of n sectors, in current and next period
- ▶ $(Ax)_i, (Bx^+)_i$: produced resp. consumed amounts of good i
- ▶ x_i^+ / x_i : growth rate of sector i

allocate activity to maximize growth rate of slowest growing sector

Optimization problems

Convex optimization

Linear optimization

Quadratic optimization

Geometric programming


Generalized inequality constraints

Vector optimization

Quadratic program (QP)

$$\begin{array}{ll}\text{minimize} & (1/2)x^T Px + q^T x + r \\ \text{subject to} & Gx \preceq h \\ & Ax = b\end{array}$$

- ▶ $P \in \mathbb{S}_+^n$ thus objective is convex quadratic
- ▶ minimize a convex quadratic function over a polyhedron



least-squares

$$\text{minimize} \quad \|Ax - b\|_2^2$$

- ▶ analytical solution $x^* = A^\dagger b$ (where A^\dagger is pseudo-inverse)
- ▶ can add linear constraints such as $l \preceq x \preceq u$

linear program with random cost

$$\begin{array}{ll}\text{minimize} & \bar{c}^T x + \gamma x^T \Sigma x \\ \text{subject to} & Gx \preceq h \\ & Ax = b\end{array}$$

- ▶ c is random vector with mean \bar{c} and covariance Σ
- ▶ hence $c^T x$ is random variable with mean $\bar{c}^T x$ and variance $x^T \Sigma x$

$$\bar{c}^T x + \gamma x^T \Sigma x = \mathbb{E} \left(c^T x \right) + \gamma \mathbf{var} \left(c^T x \right)$$

- ▶ $\gamma > 0$ is risk-aversion parameter, controls the trade-off between expected cost and variance (risk)

Quadratically constrained quadratic program (QCQP)

$$\begin{array}{ll}\text{minimize} & (1/2)x^T P_0 x + q_0^T x + r_0 \\ \text{subject to} & (1/2)x^T P_i x + q_i^T x + r_i \leq 0, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

- ▶ $P_i \in \mathbb{S}_+^n$ thus objective and constraints are convex quadratic
- ▶ feasible region is intersection of m ellipsoids and an affine set if $P_1, \dots, P_m \in \mathbb{S}_{++}^n$

Second-order cone program (SOCP)

$$\begin{array}{ll}\text{minimize} & f^T x \\ \text{subject to} & \|A_i x + b_i\|_2 \leq c_i^T x + d_i, \quad i = 1, \dots, m \\ & Fx = G\end{array}$$

with $A_i \in \mathbb{R}^{n_i \times n}$ and $F \in \mathbb{R}^{p \times n}$

- ▶ inequalities are called second-order cone constraints since

$$(A_i x + b_i, c_i^T x + d_i) \in \text{second-order cone in } \mathbb{R}^{n_i+1}$$

- ▶ if $n_i = 0$, reduces to LP
- ▶ if $c_i = 0$, reduces to QCQP (with linear objective)

Robust linear program

parameters in optimization problems are often uncertain, e.g. in LP

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & a_i^T x \leq b_i, \quad i = 1, \dots, m\end{array}$$

there can be uncertainty in c, a_i, b_i

two common approaches to handle uncertainty (in a_i for simplicity)

- ▶ deterministic model: constraints must hold for all $a_i \in \mathcal{E}_i$

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & a_i^T x \leq b_i \text{ for all } a_i \in \mathcal{E}_i, \quad i = 1, \dots, m\end{array}$$

- ▶ stochastic model: a_i is random variable; constraints must hold with probability η

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & \text{prob}(a_i^T x \leq b_i) \geq \eta, \quad i = 1, \dots, m\end{array}$$

deterministic approach via SOCP

- ▶ choose ellipsoid as \mathcal{E}_i with $\bar{a}_i \in \mathbb{R}^n$ and $P_i \in \mathbb{R}^{n \times n}$

$$\mathcal{E}_i = \{\bar{a}_i + P_i u \mid \|u\|_2 \leq 1\}$$

- ▶ robust LP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & a_i^T x \leq b_i \text{ for all } a_i \in \mathcal{E}_i, \quad i = 1, \dots, m \end{array}$$

is equivalent to SOCP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & \bar{a}_i^T x + \|P_i^T x\|_2 \leq b_i, \quad i = 1, \dots, m \end{array}$$

which follows from

$$\sup_{\|u\|_2 \leq 1} (\bar{a}_i + P_i u)^T x = \bar{a}_i^T x + \|P_i^T x\|_2$$

stochastic approach via SOCP

- ▶ assume a_i is Gaussian with mean \bar{a}_i and covariance Σ_i , namely $a_i \sim \mathcal{N}(\bar{a}_i, \Sigma_i)$
- ▶ $a_i^T x$ is Gaussian with mean $\bar{a}_i^T x$ and variance $x^T \Sigma_i x$, hence

$$\text{prob}(a_i^T x \leq b_i) = \Phi \left(\frac{b_i - \bar{a}_i^T x}{\|\Sigma_i^{1/2} x\|_2} \right)$$

with $\Phi(x) = (1/\sqrt{2\pi}) \int_{-\infty}^x e^{-t^2/2} dt$ cumulative distribution function of $\mathcal{N}(0, 1)$

- ▶ robust LP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & \text{prob}(a_i^T x \leq b_i) \geq \eta, \quad i = 1, \dots, m \end{array}$$

with $\eta > 1/2$ is equivalent to SOCP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & \bar{a}_i^T x + \Phi^{-1}(\eta) \|\Sigma_i^{1/2} x\|_2 \leq b_i, \quad i = 1, \dots, m \end{array}$$

Optimization problems

Convex optimization

Linear optimization

Quadratic optimization

Geometric programming

Generalized inequality constraints

Vector optimization

► **monomial function**

$$f(x) = cx_1^{a_1} \cdots x_n^{a_n}, \quad \text{dom } f = \mathbb{R}_{++}^n$$

with $c > 0$ and $a_i \in \mathbb{R}$

► **posynomial function**

$$f(x) = \sum_{k=1}^K c_k x_1^{a_{1k}} \cdots x_n^{a_{nk}}, \quad \text{dom } f = \mathbb{R}_{++}^n$$

sum of monomials

change variables to $y_i = \log x_i$ and take logarithm

- ▶ monomial $f(x) = cx_1^{a_1} \cdots x_n^{a_n}$ transforms to

$$\log f(e^{y_1}, \dots, e^{y_n}) = a^T y + b, \quad (b = \log c)$$

- ▶ posynomial $f(x) = \sum_{k=1}^K c_k x_1^{a_{1k}} \cdots x_n^{a_{nk}}$ transforms to

$$\log f(e^{y_1}, \dots, e^{y_n}) = \log \left(\sum_{k=1}^K e^{a_k^T y + b_k} \right), \quad (b_k = \log c_k)$$

Geometric program (GP)

geometric program in standard form

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 1, \quad i = 1, \dots, m \\ & h_i(x) = 1, \quad i = 1, \dots, p \end{array}$$


with f_i posynomial, h_i monomial

geometric program in convex form

change variables to $y_i = \log x_i$ and take logarithm of objective and constraints

$$\begin{array}{ll} \text{minimize} & \log \left(\sum_{k=1}^K e^{a_{0k}^T y + b_{0k}} \right) \\ \text{subject to} & \log \left(\sum_{k=1}^K e^{a_{ik}^T y + b_{ik}} \right) \leq 0, \quad i = 1, \dots, m \\ & Gy + d = 0 \end{array}$$

Design of cantilever beam



- ▶ N segments with unit length, rectangular cross-sections of width w_i and height h_i
- ▶ given vertical force F applied at the right end

design problem

variables w_i, h_i for $i = 1, \dots, N$

minimize total weight

subject to upper & lower bounds on w_i and h_i

upper & lower bounds on aspect ratios h_i/w_i

upper bound on stress in each segment

upper bound on vertical deflection at the end of the beam

objective and constraint functions

- ▶ total weight $w_1 h_1 + \cdots + w_N h_N$ is posynomial
- ▶ aspect ratio h_i/w_i and inverse aspect ratio w_i/h_i are monomials
- ▶ maximum stress in segment i given by $6iF/(w_i h_i^2)$ is monomial
- ▶ vertical deflection y_i and slope v_i of central axis at the right end of segment i defined recursively as (constant E is Young's modulus)

$$v_i = 12 \left(i - \frac{1}{2} \right) \frac{F}{E w_i h_i^3} + v_{i+1}$$

$$y_i = 6 \left(i - \frac{1}{3} \right) \frac{F}{E w_i h_i^3} + v_{i+1} + y_{i+1}$$

for $i = N, N-1, \dots, 1$, with $v_{N+1} = y_{N+1} = 0$, are posynomial functions

formulation as GP

$$\begin{aligned} & \text{minimize} && w_1 h_1 + \cdots + w_N h_N \\ & \text{subject to} && w_{\max}^{-1} w_i \leq 1, \quad w_{\min} w_i^{-1} \leq 1, \quad i = 1, \dots, N \\ & && h_{\max}^{-1} h_i \leq 1, \quad h_{\min} h_i^{-1} \leq 1, \quad i = 1, \dots, N \\ & && S_{\max}^{-1} w_i^{-1} h_i \leq 1, \quad S_{\min} w_i h_i^{-1} \leq 1, \quad i = 1, \dots, N \\ & && 6iF \sigma_{\max}^{-1} w_i^{-1} h_i^{-2} \leq 1, \quad i = 1, \dots, N \\ & && y_{\max}^{-1} y_1 \leq 1 \end{aligned}$$

- ▶ first two lines of constraints equivalent to

$$w_{\min} \leq w_i \leq w_{\max} \quad \text{and} \quad h_{\min} \leq h_i \leq h_{\max}$$

- ▶ third line of constraints equivalent to

$$S_{\min} \leq h_i / w_i \leq S_{\max}$$

Optimization problems

Convex optimization

Linear optimization

Quadratic optimization

Geometric programming

Generalized inequality constraints

Vector optimization

Convex problem with generalized inequality constraints

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \preceq_{K_i} 0, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

- ▶ $f_0: \mathbb{R}^n \rightarrow \mathbb{R}$ is convex
- ▶ $f_i: \mathbb{R}^n \rightarrow \mathbb{R}^{k_i}$ is K_i -convex, where K_i is a proper cone
- ▶ same properties as standard convex problem
(convex feasible set, local optimum is global, etc)

Conic form problem (cone program)

special case of above with affine objective and constraints

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Fx + g \preceq_K 0 \\ & Ax = b\end{array}$$

extends linear programming ($K = \mathbb{R}_+^m$) to nonpolyhedral cones

Semidefinite program (SDP)

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & x_1 F_1 + \cdots + x_n F_n + G \preceq 0 \\ & Ax = b\end{array}$$

with $F_i, G \in \mathbb{S}^k$

- ▶ inequality constraint is called **linear matrix inequality** (LMI)
- ▶ includes problems with multiple LMI constraints:

$$x_1 F'_1 + \cdots + x_n F'_n + G' \preceq 0 \quad \text{and} \quad x_1 F''_1 + \cdots + x_n F''_n + G'' \preceq 0$$

is equivalent to single LMI

$$x_1 \begin{bmatrix} F'_1 & 0 \\ 0 & F''_1 \end{bmatrix} + \cdots + x_n \begin{bmatrix} F'_n & 0 \\ 0 & F''_n \end{bmatrix} + \begin{bmatrix} G' & 0 \\ 0 & G'' \end{bmatrix} \preceq 0$$

LP as equivalent SDP

LP

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \preceq b\end{array}$$

equivalent SDP

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & \mathbf{diag}(Ax - b) \preceq 0\end{array}$$

note different interpretation of generalized inequality

SOCP as equivalent SDP

SOCP

$$\begin{array}{ll}\text{minimize} & f^T x \\ \text{subject to} & \|A_i x + b_i\|_2 \leq c_i^T x + d_i, \quad i = 1, \dots, m\end{array}$$

equivalent SDP

$$\begin{array}{ll}\text{minimize} & f^T x \\ \text{subject to} & \begin{bmatrix} (c_i^T x + d_i)I & A_i x + b_i \\ (A_i x + b_i)^T & c_i^T x + d_i \end{bmatrix} \succeq 0, \quad i = 1, \dots, m\end{array}$$

Eigenvalue minimization

$$\text{minimize} \quad \lambda_{\max}(A(x))$$

where $A(x) = A_0 + x_1 A_1 + \cdots + x_n A_n$ with given $A_i \in \mathbb{S}^k$

equivalent SDP

$$\begin{array}{ll} \text{minimize} & t \\ \text{subject to} & A(x) \preceq tI \end{array}$$

- ▶ variables $x \in \mathbb{R}^n$ and $t \in \mathbb{R}$
- ▶ follows from

$$\lambda_{\max}(A) \leq t \quad \Longleftrightarrow \quad A \preceq tI$$

Matrix norm minimization

$$\text{minimize} \quad \|A(x)\|_2 = \left(\lambda_{\max} \left(A(x)^T A(x) \right) \right)^{1/2}$$

where $A(x) = A_0 + x_1 A_1 + \cdots + x_n A_n$ with given $A_i \in \mathbb{R}^{p \times q}$

equivalent SDP

$$\begin{array}{ll} \text{minimize} & t \\ \text{subject to} & \begin{bmatrix} tI & A(x) \\ A(x)^T & tI \end{bmatrix} \succeq 0 \end{array}$$

- ▶ variables $x \in \mathbb{R}^n$ and $t \in \mathbb{R}$
- ▶ follows from

$$\begin{aligned} \|A\|_2 \leq t & \iff A^T A \preceq t^2 I, \quad t \geq 0 \\ & \iff \begin{bmatrix} tI & A \\ A^T & tI \end{bmatrix} \succeq 0 \end{aligned}$$

Optimization problems

Convex optimization

Linear optimization

Quadratic optimization

Geometric programming

Generalized inequality constraints

Vector optimization

Vector optimization

general vector optimization problem

$$\begin{array}{ll} \text{minimize (with respect to } K) & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p \end{array}$$

vector objective $f_0: \mathbb{R}^n \rightarrow \mathbb{R}^q$ minimized with respect to proper cone $K \subseteq \mathbb{R}^q$

convex vector optimization problem

$$\begin{array}{ll} \text{minimize (with respect to } K) & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b \end{array}$$



where f_0 is K -convex and f_1, \dots, f_m are convex

Optimal and Pareto optimal points

set of achievable values

$$\mathcal{O} = \{f_0(x) \mid x \text{ feasible}\}$$

- ▶ feasible x is optimal if $f_0(x)$ is the minimum value of \mathcal{O} (optimal value)
- ▶ feasible x is Pareto optimal if $f_0(x)$ is a minimal value of \mathcal{O} (Pareto optimal value)



Multicriterion optimization

vector optimization problem with $K = \mathbb{R}_+^q$

$$f_0(x) = (F_1(x), \dots, F_q(x))$$

- ▶ q different objectives F_i , we want all of them to be small
- ▶ feasible x^* is optimal if

$$y \text{ feasible} \implies f_0(x^*) \preceq f_0(y)$$

if there exists an optimal point, the objectives are noncompeting


- ▶ feasible x^{po} is Pareto optimal if

$$y \text{ feasible, } f_0(y) \preceq f_0(x^{\text{po}}) \implies f_0(x^{\text{po}}) = f_0(y)$$

if multiple Pareto optimal values exist, there is a trade-off between the objectives

Regularized least-squares

minimize (with respect to \mathbb{R}_+^2) $(\|Ax - b\|_2^2, \|x\|_2^2)$





the optimal trade-off curve, shown darker, is formed by Pareto optimal points

Risk-return trade-off in portfolio optimization

$$\begin{array}{ll} \text{minimize (with respect to } \mathbb{R}_+^2) & \left(-\bar{p}^T x, x^T \Sigma x \right) \\ \text{subject to} & \mathbf{1}^T x = 1 \\ & x \succeq 0 \end{array}$$

- ▶ $x \in \mathbb{R}^n$ investment portfolio; x_i fraction invested in asset i
- ▶ $p \in \mathbb{R}^n$ relative asset price changes, random variable with mean \bar{p} and covariance Σ
- ▶ $\mathbb{E}r = \bar{p}^T x$ expected return; $\text{var } r = x^T \Sigma x$ return variance




to find Pareto optimal points, choose $\lambda \succ_{K^*} 0$ and solve scalar problem

$$\begin{array}{ll} \text{minimize} & \lambda^T f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p \end{array}$$

- ▶ if x is optimal for scalar problem, then it is Pareto optimal for vector optimization problem
- ▶ for convex vector optimization problem, can find (almost) all Pareto optimal points by varying $\lambda \succ_{K^*} 0$

example




- ▶ Pareto optimal values $f_0(x_1)$ and $f_0(x_2)$ can both be obtained by scalarization: $f_0(x_1)$ minimizes $\lambda_1^T u$ and $f_0(x_2)$ minimizes $\lambda_2^T u$ over all $u \in \mathcal{O}$
- ▶ $f_0(x_3)$ is Pareto optimal, but cannot be found by scalarization

Scalarization for multicriterion problems

to find Pareto optimal points, minimize positive weighted sum (since $K = \mathbb{R}_+^q$)

$$\lambda^T f_0(x) = \lambda_1 F_1(x) + \cdots + \lambda_q F_q(x)$$

examples



- ▶ regularized least-square problem: take $\lambda = (1, \gamma)$ with $\gamma > 0$

$$\text{minimize} \quad \|Ax - b\|_2^2 + \gamma \|x\|_2^2$$

least-square problem for fixed $\gamma > 0$

- ▶ risk-return trade-off problem: take $\lambda = (1, \gamma)$ with $\gamma > 0$

$$\begin{aligned} &\text{minimize} && -\bar{p}^T x + \gamma x^T \Sigma x \\ &\text{subject to} && \mathbf{1}^T x = 1 \\ &&& x \succeq 0 \end{aligned}$$

quadratic program for each fixed $\gamma > 0$