An skeleton assessment for Bristol Bay Red King crab in GMACS

Cody Szuwalski & Matthieu Veron

June 2022 - GMACS version 2.01.K

Contents

A. Summary of Major Changes

B. Comments, responses, and assessment summary


С.	Introduction
	Distribution
	Life history characteristics
	Natural mortality
	Weight at length
	Maturity
	Molting probability
	Reproduction
	Growth
	Management history
	ADFG harvest strategy
	History of BMSY
	Fishery history
D.	Data
	Catch data
	Survey biomass and size composition data
	Spatial distribution
Ε.	Analytic approach
	History of modeling approaches
	Model description

F.	Calculation of the OFL						
	Methodology of the OFL $\ \ldots \ \ldots \ \ldots \ \ldots \ \ldots \ \ldots \ \ldots$						
	Calculated OFLs and interpretation						
	Projections under harvest strategies						
\mathbf{G}	Calculation of the ABC						
	Uncertainty in the ABC						
	Author recommendations						
Η.	Data gaps and research priorities						
	Methodology						
	Data sources						
	Scientific uncertainty						
I.	Ecosystem considerations						
Appendix A: Population dynamics							

- 1. Stock: Bristol Bay red king crab.
- 2. Catches: trends and current levels
- 3. Stock Biomass:
- 4. Recruitment

- 5. Management
- 6. Basis for the OFL
- 7. Probability Density Function of the OFL
- 8. Basis for ABC

A. Summary of Major Changes



D. Data

Catch data

Survey biomass and size composition data

Spatial distribution

E. Analytic approach

History of modeling approaches

Model description

Model selection and evaluation

Results

Fits to data

Estimated population processes and derived quantities

F. Calculation of the OFL

Methodology of the OFL

Calculated OFLs and interpretation

Projections under harvest strategies

G. Calculation of the ABC

Uncertainty in the ABC

Author recommendations

H. Data gaps and research priorities

Methodology

Data sources

Scientific uncertainty

I. Ecosystem considerations

Appendix A: Population dynamics

Table 1: Changes in management quantities for each scenario considered. Reported management quantities are derived from maximum likelihood estimates.

Model	MMB	B35	F35	FOFL	OFL
Version 2.01.E	14025.96	2991.975	0.880	0.880	8342.826
Version $2.01.F$	14025.96	2991.975	0.880	0.880	8342.826
Version 2.01.G	14025.96	2991.975	0.880	0.880	8342.826
Version 2.01.H	14025.96	2991.975	0.880	0.880	8342.826
Version 2.01.I	14025.96	22512.179	0.299	0.186	2425.493
Version 2.01.K	14025.96	22512.179	0.299	0.186	2425.493




Figure 1: Model fits to the NMFS trawl survey.




Figure 2: Model fits to the BSFRF survey.




Figure 3: Model fits to the catch data.




Figure 4: Estimated fishing mortality by fleet.




Figure 5: Estimated probability of molting.




Figure 6: Estimated selectivity.




Figure 7: Estimated recruitment.




Figure 8: Estimated mature biomass






Figure 9: Estimated size at recruitment




Figure 10: Kobe plot for the Pot Fishery fleet.