Higher-Degree Polynomial Families on $x^4 + y^3 + z^2 + 1 = 0$: Covariant Syzygies, Resolution of Degree-$(3, 8, 12)$ and $(3, 10, 15)$ Systems, Modulo-4 Frobenius Obstructions in $\mathbb{Z}[t]$, and General Theorems for $\deg x \geq 3$

Research Continuation Report

Extending the results of *Integer solutions of* $x^4 + y^3 + z^2 + 1 = 0$ (15 October 2026)

17 October 2026

Abstract

We extend the study of polynomial solutions $(x(t), y(t), z(t))$ to the affine surface $S : x^4+y^3+z^2+1 = 0$ (equivalently $x^4 - Y^3 + z^2 + 1 = 0$ with $Y = -y$) across both $\mathbb{Q}[t]^3$ and $\mathbb{Z}[t]^3$ (as well as $\mathbb{Z}_+[t]^3$) for arbitrary $\deg x \geq 3$. Building on our covariant triangularization of the cubic trace-zero system and universal Davenport–Stothers–Zannier bounds, we establish nine principal results: (1) positivity lower bounds $\deg Y \geq \lceil 4d_x/3 \rceil$ and canonical Galois trace-zero divisorial reductions for all $d_x = \deg x \geq 1$; (2) universal upper bounds $\deg Y = 2\nu \leq 8d_x - 2$ (for $\gcd(Y, z) = 1$, automatic over $\mathbb{Q}$ when $d_x \in \{3, 4\}$) and $\deg Y \leq 10d_x - 4$ unconditionally for all $d_x \geq 3$; (3) the binary cubic covariant syzygy $4p^3 + 27N^2 = (27u^2 + 4s^3)I^2$ and $\mathbb{G}_{\mathrm{m}}^2$-bihomogeneous reduction for $\deg x = 3$; (4) complete resolution of the open $\nu = 4$ ($\deg x = 3, \deg Y = 8, \deg z = 12$) system over $\mathbb{Q}$, together with the discovery of an explicit degree-$(3, 8, 12)$ solution over $\mathbb{Q}(\sqrt{3}, 2^{1/3})$; (5) complete resolution of the $\nu = 5$ ($\deg x = 3, \deg Y = 10, \deg z = 15$) system in both the constant-m branch ($m_1 = 0$) and the linear-m branch ($m_1 \neq 0$) via a denominator-free elimination of g_0, a parity decomposition in $(F, \tilde{b}_2)$, and an explicit resultant curve $H_9(\tilde{B}_1, s) = 0$; (6) infinite towers of polynomial identities over $\mathbb{Q}(\sqrt{3}, 2^{1/3})$ of degrees $(3k, 4k, 6k)$ and $(3k, 8k, 12k)$ arising from the D_4^* rational elliptic surface $\mathcal{E} : \tilde{z}^2 = \tilde{Y}^3 - (w^3 + 1)$; (7) strict equivalence between nonconstant polynomial solutions in $\mathbb{Z}[t]^3$ and in $\mathbb{Z}_+[t]^3$; (8) an unconditional Modulo-4 Frobenius, $\mathbb{F}_4$-Root, and Formal Derivative Theorem in $\mathbb{Z}[t]$ proving that $\bar{Y}(t) = \bar{w}(t)^4 \pmod 2$ with $8 \mid \deg_{\mathbb{F}_2}(Y \bmod 2) \leq 2\deg x - 2$, forcing $Y(t) \equiv 1 + 2x(t)^4 \pmod 4$ and $\deg Y \geq 4\deg_{\mathbb{F}_2}(x \bmod 2)$ whenever $\deg x \leq 4$; and (9) general conditional nonexistence theorems for arbitrary $\deg x \geq 3$ under the Polynomial Hall–Vojta Conjecture and Severi stratum obstructions.

1 Introduction and Context

Let $S \subset \mathbb{A}^3_{\mathbb{Q}}$ denote the smooth affine surface defined by

$$x^4+y^3+z^2+1 = 0, \qquad \text{or equivalently} \qquad x^4-Y^3+z^2+1 = 0 \iff Y^3-z^2 = x^4+1 \quad (Y = -y). \tag{1}$$

Whether $S(\mathbb{Z})$ is infinite is a well-known open problem highlighted by Grechuk [1, 2]. In the companion paper [3], viewing S as the affine complement of the zero section O and the type-IV fibre F_∞ on the rational elliptic surface $\pi : \hat{E} \to \mathbb{P}^1_x$ with Mordell–Weil lattice $\mathrm{MW}(\hat{E}/\overline{\mathbb{Q}}(x)) \cong E_6^*$ [4, 5] and Frobenius-trace verification $\mathrm{MW}(\hat{E}/\mathbb{Q}(x)) = 0$ established two foundational nonexistence theorems for polynomial families $(x(t), y(t), z(t)) \in \mathbb{Q}[t]^3$:

- **Theorem A ($\deg x \leq 2$):** There is no nonconstant polynomial solution $(x(t), y(t), z(t)) \in \mathbb{Q}[t]^3$ with $\deg x \leq 2$ for any y, z.

Result	Tag	Summary and Section Reference
Theorem 2.1 (Lower Bound & Trace)	[E]	For all $d_x = \deg x \geq 1$, $\deg Y \geq \lceil 4d_x/3 \rceil$ over $\mathbb{R}$, and $\sum_{i=1}^{d_x}[P_i] = O$ in $E(\mathbb{Q}(x))$ gives a canonical divisorial reduction of $z(t)$ (Section 2).
Theorem 2.3 (Universal Upper Bound)	[E]	For all $d_x \geq 3$ and any $(Y, z) \in \mathbb{C}[t]^2$, $\deg Y = 2\nu \leq 8d_x - 2$ ($\nu \leq 4d_x - 1$); if $d_x = 3k$, either $\deg Y = 4k$ or $2\nu > 4k$ (Section 2).
Theorem 3.1 (Covariant Syzygy)	[E]	$p = \frac{1}{12}\operatorname{Hess}(I)$ and $N = -\frac{1}{9}\operatorname{Jac}(I, p)$ satisfy $4p^3 + 27N^2 = (27u^2 + 4s^3)I^2$; $\Phi(u) \equiv 0$ is $\mathbb{G}_{\mathrm{m}}^2$-bihomogeneous in degrees $u^{\geq 5}$ (Section 3).
Theorem 4.1 (Explicit $(3, 8, 12)$ Curve)	[E]	Explicit polynomial solution of degrees $(3, 8, 12)$ over $\mathbb{Q}(\sqrt{3}, 2^{1/3})$ for $\nu = 4$, refuting emptiness of $\nu \geq 4$ moduli over $\overline{\mathbb{Q}}$ (Section 4.1).
Theorem 4.2 (Exclusion of $\nu = 4$ over $\mathbb{Q}$)	[E]/[C]	Complete resolution of the open $\deg x = 3, \deg Y = 8, \deg z = 12$ system over $\mathbb{Q}$ in both $m_1 = 0$ and $m_1 \neq 0$ branches (Sections 4.1–4.2).
Theorems 5.1–5.2 (Exclusion of $\nu = 5$)	[E]/[C]	Complete reduction and elimination of $\deg x = 3, \deg Y = 10, \deg z = 15$ in both $m_1 = 0$ and $m_1 \neq 0$ branches (Section 5).
Theorem 6.1 (Pullback Towers for $d_x = 3k$)	[E]	Pullback to $\tilde{z}^2 = \tilde{Y}^3 - (w^3 + 1)$ yields families of degrees $(3k, 4k, 6k)$ and $(3k, 8k, 12k)$ over $\mathbb{Q}(\sqrt{3}, 2^{1/3})$ for all $k \geq 1$ (Section 6).
Theorems 7.1–7.4 ($\mathbb{Z}[t]$ Obstructions)	[E]	Equivalence $\mathbb{Z}[t]^3 \Leftrightarrow \mathbb{Z}_+[t]^3$; Modulo-4 Frobenius & $\mathbb{F}_2[t]$-derivative bound $8 \mid \deg_{\mathbb{F}_2}(Y \bmod 2) \leq 2\deg x - 2$ and $\deg Y \geq 4m_x$ for $d_x \leq 4$ (Section 7).
Theorems 8.1–8.3 (General $d_x \geq 3$)	[E]	General conditional nonexistence for arbitrary $\deg x \geq 3$ via Polynomial Hall–Vojta, Severi codimension, and 2-adic parity (Section 8).

Table 1: Summary of new theorems and verification status ([E] = complete elementary/analytic proof by hand; [C] = exact symbolic computation in SymPy).

- **Theorem B ($\deg x = 3, \deg y \leq 4$):** There is no nonconstant polynomial solution in $\mathbb{Q}[t]^3$ with $\deg x = 3$ and $\deg y \leq 4$ (which forces degrees $(3, 4, 6)$).

However, Section 7 of [3] left two major regimes open:

1. **The higher-degree regime for $\deg x = 3$ ($\deg y \geq 5$):** Proposition 7.1 of [3] showed that any solution with $\deg x = 3$ and $d = \deg Y \geq 5$ must satisfy $d = 2\nu \in \{8, 10, 14, 16, 20, 22\}$ ($\nu \in \{4, 5, 7, 8, 10, 11\}$). For the very first open case $\nu = 4$ ($d = 8, \deg z = 12$), eliminating $n(x)$ via $4m^2e_3 - e_2^2 - 4m^2(x^4 + 1) = 0$ produced a system of 11 equations of degree up to 8 in 10 unknowns for which Gröbner basis computations in Singular modulo 32003 failed to terminate, while boundary dimension counts in Appendix C of [3] gave expected dimension $3 - \nu < 0$, motivating the heuristic conjecture that the moduli spaces for $\nu \geq 4$ are empty even over $\overline{\mathbb{Q}}$.
2. **The regime $\deg x \geq 4$ for arbitrary y and z:** Nothing was proved for $\deg x \geq 4$, where the relevant divisor classes have positive expected dimension growing linearly with $\deg x$.

In this paper, we develop a comprehensive algebraic, geometric, and 2-adic/3-adic framework that extends the theory to $\deg x \geq 3$ for arbitrary $y(t)$ and $z(t)$, completely resolves the open $\nu = 4$ ($\deg Y = 8$) and $\nu = 5$ ($\deg Y = 10$) systems for $\deg x = 3$, proves a powerful unconditional Modulo-4 Frobenius and Derivative Obstruction for polynomial solutions in $\mathbb{Z}[t]^3$ of arbitrary degree, and establishes general conditional nonexistence theorems for all $\deg x \geq 3$.

2 Universal Bounds and Divisorial Reduction for Arbitrary $\deg x \geq 3$

We begin by establishing general structural theorems valid for arbitrary degree $d_x = \deg x \geq 1$ and arbitrary polynomials $Y(t), z(t)$.

2.1 Positivity Lower Bound and Trace-Zero Divisorial Decomposition

Theorem 2.1 (Positivity Lower Bound and Trace-Zero Divisorial Reduction)**.** *Let $(x(t), y(t), z(t)) \in \mathbb{Q}[t]^3$ be a nonconstant solution to $x^4 + y^3 + z^2 + 1 = 0$, and put $Y(t) = -y(t)$, $d_x = \deg x \geq 1$, $d_Y = \deg Y$, and $d_z = \deg z$.*

(i) ***Universal lower bound over $\mathbb{R}$ and $\mathbb{Q}$:*** *For every nonconstant solution in $\mathbb{R}[t]^3$ (and hence in $\mathbb{Q}[t]^3$), the leading terms of $x(t)^4$ and $z(t)^2$ cannot cancel, so $3d_Y = \max(4d_x, 2d_z) \geq 4d_x$. Consequently,*

$$\deg Y = d_Y \geq \left\lceil \frac{4d_x}{3} \right\rceil, \qquad \deg z = d_z \leq \frac{3}{2} d_Y. \tag{2}$$

In particular, if $3 \nmid d_x$, then $3d_Y > 4d_x$, which forces $3d_Y = 2d_z$, so $d_Y = 2\nu$ and $d_z = 3\nu$ for some integer $\nu \geq \lceil (2d_x + 1)/3 \rceil$.

(ii) ***Galois trace and divisorial interpolation:*** *Assume $(x(t), Y(t), z(t))$ is injective on a dense open subset of $\mathbb{A}^1$. Let $t_1, \ldots, t_{d_x}$ be the roots of $x(t) = x$ in $\overline{\mathbb{Q}(x)}$ and $P_i = (Y(t_i), z(t_i)) \in E(\overline{\mathbb{Q}(x)})$ the corresponding points on the generic fibre $E_x : z^2 = Y^3 - (x^4 + 1)$. Then*

$$P_1 + P_2 + \cdots + P_{d_x} = O \quad \text{in } E(\mathbb{Q}(x)). \tag{3}$$

Consequently, there exist unique polynomials $A(x, Y), B(x, Y) \in \mathbb{Q}[x, Y]$ with $\deg_Y A \leq \lfloor d_x/2 \rfloor$, $\deg_Y B \leq \lfloor (d_x - 3)/2 \rfloor$, and $\gcd_{\mathbb{Q}(x)[Y]}(A, B) = 1$ such that the divisor of zeroes of $A(x, Y) - B(x, Y)z$ on E_x is precisely $\sum_{i=1}^{d_x} [P_i]$, and the characteristic polynomial of $Y(t)$ over $\mathbb{Q}(x)$ is given up to a factor in $\mathbb{Q}(x)^*$ by the quotient

$$G_{d_x}(x, Y) = \frac{A(x, Y)^2 - B(x, Y)^2 (Y^3 - x^4 - 1)}{H(x, Y)}, \tag{4}$$

where $H(x, Y) \in \mathbb{Q}[x, Y]$ has $\deg_Y H = 2 \deg_Y B + 3 - d_x \in \{0, 1, 2\}$. In particular, when $B(x, Y) \in \mathbb{Q}[x]$ is independent of Y (which always holds for $d_x = 3$ and $d_x = 4$ whenever $P_1, \ldots, P_{d_x}$ are in general position), $z(t)$ admits the rational representation $z(t) = \frac{A(x(t), Y(t))}{B(x(t))}$.

Proof. (i) Write $x(t) = x_{d_x} t^{d_x} + \ldots$ and $z(t) = z_{d_z} t^{d_z} + \ldots$ with $x_{d_x} \in \mathbb{R}^*$ and $z_{d_z} \in \mathbb{R}$. In $Y(t)^3 = x(t)^4 + z(t)^2 + 1$, the leading coefficient of $x(t)^4$ is $x_{d_x}^4 > 0$ and that of $z(t)^2$ is $z_{d_z}^2 \geq 0$. Since both are non-negative and $x_{d_x}^4 > 0$, the highest-degree term of $x(t)^4 + z(t)^2 + 1$ has degree $\max(4d_x, 2d_z)$ and strictly positive coefficient $x_{d_x}^4$, $z_{d_z}^2$, or $x_{d_x}^4 + z_{d_z}^2$. Thus $3d_Y = \max(4d_x, 2d_z) \geq 4d_x$, giving $d_Y \geq \lceil 4d_x/3 \rceil$. If $3 \nmid d_x$, then $3d_Y \neq 4d_x$, so $3d_Y = 2d_z > 4d_x$, whence $d_Y = 2\nu$ and $d_z = 3\nu$.

(ii) The zero-cycle $\sum_{i=1}^{d_x} [P_i]$ on $E_{\mathbb{Q}(x)}$ is invariant under $\mathrm{Gal}(\overline{\mathbb{Q}(x)}/\mathbb{Q}(x))$, so its Abel–Jacobi sum lies in $E(\mathbb{Q}(x)) = \mathrm{MW}(\hat{E}/\mathbb{Q}(x)) = \{O\}$ by Proposition 2.3 and Corollary 2.4 of [3]. By the Riemann–Roch theorem on the elliptic curve $E_{\mathbb{Q}(x)}$, the linear system $\mathcal{L}(d_x O)$ has dimension d_x with basis $\{1, Y, z, Y^2, Yz, \ldots\}$ of pole orders $\{0, 2, 3, 4, 5, \ldots, d_x\}$ at O. Since $\sum_{i=1}^{d_x} [P_i] \sim d_x O$ as divisors on $E_{\mathbb{Q}(x)}$, there is a unique function (up to $\mathbb{Q}(x)^*$-scaling) $f = A(x, Y) - B(x, Y)z \in \mathcal{L}(d_x O)$ vanishing on $\sum_{i=1}^{d_x} [P_i]$ (or a minimal sub-sum if a proper subset already sums to O). Taking norms from $\mathbb{Q}(x)[Y, z]/(z^2 - Y^3 + x^4 + 1)$ to $\mathbb{Q}(x)[Y]$ yields $\mathrm{N}(f) = A(x, Y)^2 - B(x, Y)^2 (Y^3 - x^4 - 1)$, whose divisor of zeroes on E_x is $\sum_{i=1}^{d_x} ([P_i] + [-P_i]) + R$, where R consists of at most two auxiliary points when $\max(2 \deg_Y A, 2 \deg_Y B + 3) > d_x$, projecting to $H(x, Y)$ of degree ≤ 2 in Y. □

Corollary 2.2 (Exclusion of Pullback and Unramified Families for Arbitrary $d_x \geq 3$)**.** *For any $d_x = \deg x \geq 3$, there is no nonconstant polynomial solution $(x(t), y(t), z(t)) \in \mathbb{Q}[t]^3$ to (1) satisfying any of the following:*

(a) $(x(t), y(t), z(t)) = (\tilde{x}(q(t)), \tilde{y}(q(t)), \tilde{z}(q(t)))$ *factors through a polynomial* $q(t) \in \mathbb{Q}[t]$ *with* $\deg \tilde{x} \in \{1, 2\}$*, or with* $\deg \tilde{x} = 3$ *and* $\deg \tilde{y} \leq 10$*;*

(b) $Y(t) = Y_0(x(t))$ *lies in* $\mathbb{Q}[x(t)]$*;*

(c) $z(t) = z_0(x(t))$ *lies in* $\mathbb{Q}[x(t)]$*.*

Proof. Part (a) follows immediately by substituting $s = q(t)$ and applying Theorem A ($\deg \tilde{x} \leq 2$), Theorem B ($\deg \tilde{x} = 3, \deg \tilde{y} \leq 4$), Theorem 4.2 ($\deg \tilde{x} = 3, \deg \tilde{y} = 8$), and Theorems 5.1–5.2 ($\deg \tilde{x} = 3, \deg \tilde{y} = 10$). Parts (b) and (c) follow from $\mathrm{MW}(\hat{E}/\mathbb{Q}(x)) = 0$ and positivity of $f(x) = x^4 + z_0(x)^2 + 1 \geq 1$ on $\mathbb{R}$. □

2.2 Universal Upper Bounds for Arbitrary $\deg x \geq 3$

Theorem 2.3 (Universal Degree Bound for Arbitrary $\deg x \geq 3$)**.** *Let* $(x(t), Y(t), z(t)) \in \mathbb{C}[t]^3$ *be a nonconstant polynomial solution to* $Y(t)^3 - z(t)^2 = x(t)^4 + 1$ *with* $d_x = \deg x \geq 3$*,* $d_Y = \deg Y$*, and* $d_z = \deg z$*. Assume that* $Y(t)$ *and* $z(t)$ *are coprime in* $\mathbb{C}[t]$ *(which holds automatically over* $\mathbb{Q}$ *whenever every critical value of* $x(t)$ *has degree* ≤ 3 *over* $\mathbb{Q}$*, and in particular for* $d_x \in \{3, 4\}$*). Then* (d_Y, d_z) *falls into one of three finite regimes:*

(1) ***Sub-canonical regime (****$3d_Y < 4d_x$****, over*** $\mathbb{C} \setminus \mathbb{R}$ ***only):*** $d_z = 2d_x$ *and* $1 \leq d_Y \leq \lfloor (4d_x - 1)/3 \rfloor$*. By Theorem 2.1(i), this regime contains* ***no solutions over*** $\mathbb{R}$ ***or*** $\mathbb{Q}$*.*

(2) ***Canonical regime (****$3d_Y = 4d_x$****):*** $d_x = 3k$ *is a multiple of* 3*,* $d_Y = 4k$*, and* $d_z \leq 6k$ *(with* $d_z = 6k$ *over* $\mathbb{R}$ *or* $\mathbb{Q}$*).*

(3) ***Hall–Davenport regime (****$3d_Y = 2d_z > 4d_x$****):*** $d_Y = 2\nu$ *is even and* $d_z = 3\nu$ *for an integer* ν *satisfying the universal bounds*

$$\left\lfloor \frac{2d_x}{3} \right\rfloor + 1 \leq \nu \leq 4d_x - 1, \qquad \text{so} \qquad \frac{4d_x + 2}{3} \leq d_Y \leq 8d_x - 2. \tag{5}$$

Without assuming $\gcd(Y(t), z(t)) = 1$*, let* $g_{\mathrm{cd}} = \deg \gcd(Y(t), z(t))$*. Then* $g_{\mathrm{cd}} \leq d_x - 1$*, and the Mason–Stothers theorem gives* $\nu \leq 5d_x - 2$ *(*$d_Y \leq 10d_x - 4$*) unconditionally for all* $d_x \geq 3$*.*

Proof. When $3d_Y > 4d_x$ and $\gcd(Y, z) = 1$, Davenport's inequality [6, 7, 9] gives $4d_x = \deg(Y^3 - z^2) \geq \frac{1}{2} \deg Y + 1 = \nu + 1$, so $\nu \leq 4d_x - 1$. In general, any root α of $D_0 = \gcd(Y, z)$ is a root of $x(t)^4 + 1$ of multiplicity $m_\alpha \geq 2$, hence a root of $x'(t)$ of multiplicity $m_\alpha - 1 \geq \frac{1}{2} m_\alpha$, giving $g_{\mathrm{cd}} \leq d_x - 1$; Mason–Stothers [8] applied to $(Y^3/D_0^2, -z^2/D_0^2, -(x^4 + 1)/D_0^2)$ yields $6\nu - 2g_{\mathrm{cd}} \leq 5\nu + 4d_x - g_{\mathrm{cd}} - 1$, so $\nu \leq 4d_x + g_{\mathrm{cd}} - 1 \leq 5d_x - 2$. □

3 Binary Cubic Covariants and Bihomogeneous Torus Reduction for $\deg x = 3$

For $\deg x = 3$ and $d = \deg Y = 2\nu \geq 4$, normalise $x(t) = t^3 + st + x_0$ and put $u = x - x_0 = t^3 + st$. By Lemma 6.1 of [3], $z(t) = m(u)Y(t) + n(u)$ with $m(u), n(u) \in \mathbb{Q}[u]$, $m(u) \neq 0$, and $Y(t) = \frac{1}{3} m(u)^2 + W(t)$, where $W(t) = \frac{2}{3} s\gamma(u) + \beta(u)t + \gamma(u)t^2$ has $\mathrm{Tr}_{\mathbb{Q}(t)/\mathbb{Q}(u)}(W) = 0$.

Theorem 3.1 (Binary Cubic Covariant Structure and Unified Bihomogeneous Identity)**.** *Let* $W(t) = \frac{2}{3} s\gamma(u) + \beta(u)t + \gamma(u)t^2 \in \mathbb{Q}[u][t]/(t^3 + st - u)$*, and let* $T^3 + p(u)T - N(u)$ *be the characteristic polynomial of* W *over* $\mathbb{Q}[u]$*.*

(i) ***Covariant representation and discriminant syzygy:*** *Let* $I(\beta, \gamma) = \beta^3 + s\beta\gamma^2 - u\gamma^3$ *be the index cubic of* W*. Then* $p(u)$ *and* $N(u)$ *are the Hessian and Jacobian covariants of* $I(\beta, \gamma)$*:*

$$p(u) = \frac{1}{12} \mathrm{Hess}_{\beta,\gamma}(I) = s\beta^2 - 3u\beta\gamma - \frac{1}{3} s^2\gamma^2, \tag{6}$$

$$N(u) = -\frac{1}{9} \mathrm{Jac}_{\beta,\gamma}(I, p) = u\beta^3 + \frac{2}{3} s^2\beta^2\gamma - su\beta\gamma^2 + \left(u^2 + \frac{2}{27} s^3\right)\gamma^3, \tag{7}$$

satisfying the universal covariant syzygy $4p(u)^3 + 27N(u)^2 = (27u^2 + 4s^3)\, I(\beta(u), \gamma(u))^2$.

(ii) ***Unified polynomial identity:*** *A tuple* $(s, x_0, \beta(u), \gamma(u), m(u))$ *with* $m(u) \neq 0$ *defines a polynomial solution to* (1) *if and only if it satisfies the single polynomial identity in* $\mathbb{Q}[u]$*:*

$$\Phi(u) := \Psi(u) - 4m(u)^2((u + x_0)^4 + 1) \equiv 0, \qquad \Psi(u) := 4m^2N - p^2 + \frac{2}{3}m^4p + \frac{1}{27}m^8, \quad (8)$$

which automatically implies $m(u) \mid p(u)$ *and* $n(u) = -\frac{p(u)}{2m(u)} - \frac{1}{6}m(u)^3 \in \mathbb{Q}[u]$.

(iii) ***Exact degree bounds and*** $\mathbb{G}_{\mathrm{m}}^2$***-bihomogeneity:*** *For* $d = 2\nu \in \{8, 10, 14, 16, 20, 22\}$,

$$\deg_u \beta \leq \left\lfloor \frac{2\nu - 1}{3} \right\rfloor, \quad \deg_u \gamma \leq \left\lfloor \frac{2\nu - 2}{3} \right\rfloor, \quad \mu := \deg_u m \leq \left\lfloor \frac{\nu}{3} \right\rfloor \leq \frac{\nu - 1}{3}, \quad \deg_u n = \nu, \quad (9)$$

and $\Psi(u)$ *is bihomogeneous of weight* μ^8 *under the torus* $\mathbb{G}_{\mathrm{m}}^2 = \{(\lambda, \mu) \in (\overline{\mathbb{Q}}^*)^2\}$ *acting via*

$$(u, s, x_0) \mapsto (\lambda^3 u, \lambda^2 s, \lambda^3 x_0), \qquad (\beta, \gamma, m, n) \mapsto (\lambda^{-1}\mu^2\beta,\ \lambda^{-2}\mu^2\gamma,\ \mu m,\ \mu^3 n). \quad (10)$$

4 Complete Resolution of the Open $\nu = 4$ $(\deg x = 3, \deg Y = 8, \deg z = 12)$ System

For $\nu = 4$ ($d = \deg_t Y = 8, \deg_t z = 12$), writing $\beta(u) = b_2u^2 + b_1u + b_0$, $\gamma(u) = g_2u^2 + g_1u + g_0$ ($g_2 \neq 0$), and $m(u) = m_1u + m_0$, the leading coefficient of $\Phi(u)$ at u^{10} is

$$\mathrm{coeff}_{u^{10}}(\Phi) = -g_2^2(9b_2^2 - 4g_2m_1^2) = 0 \implies 9b_2^2 = 4g_2m_1^2. \quad (11)$$

4.1 The Constant-m Branch ($m_1 = 0$) and the Explicit $(3, 8, 12)$ Curve over $\overline{\mathbb{Q}}$

Theorem 4.1 (Complete Solution of the $m_1 = 0$ Branch for $\nu = 4$)**.** *In the case* $\deg x = 3, \deg Y = 8, \deg z = 12$ *with* $m(u) = m_0 \in \overline{\mathbb{Q}}^*$ *constant (so* $b_2 = 0$*):*

(i) ***No rational solutions:*** *There is no solution* $(x(t), y(t), z(t)) \in \mathbb{Q}[t]^3$ *with* $m_1 = 0$.

(ii) ***Explicit solution over*** $\mathbb{Q}(\sqrt{3}, 2^{1/3})$***:*** *Over* $\overline{\mathbb{Q}}$*, setting* $s = 0, x_0 = 0, b_1 = -2, g_2 = \frac{3}{2}2^{1/3}, m_0 = -2^{1/3}\sqrt{3}$ *yields the explicit polynomial identity of degrees* $(3, 8, 12)$*:*

$$x(t) = t^3, \quad Y(t) = \frac{3}{2}\,2^{1/3}t^8 - 2t^4 + 2^{2/3}, \quad z(t) = \sqrt{3}\left(\frac{3}{2}\,t^{12} - \frac{3}{2}\,2^{2/3}t^8 + 2 \cdot 2^{1/3}t^4 - 1\right), \quad (12)$$

which satisfies $Y(t)^3 - z(t)^2 = x(t)^4 + 1$ *identically in* $\mathbb{Q}(\sqrt{3}, 2^{1/3})[t]$*, disproving the heuristic expectation in Appendix C of [3] that the* $\nu \geq 4$ *moduli spaces are empty over* $\overline{\mathbb{Q}}$.

Proof. From $\mathrm{coeff}_{u^8}(\Phi) = 0$, $m_0^2 = (9b_1 + g_2s^2)^2/(36g_2)$. Then u^7 and u^6 solve linearly for $b_0 = \frac{g_1(9b_1 - g_2s^2)}{18g_2}$ and $g_0 = \frac{324b_1^2g_2s + 243b_1g_1^2 + 36b_1g_2^2s^3 + 27g_1^2g_2s^2 - 8g_2^3s^5}{108g_2(9b_1 - g_2s^2)}$, and u^5 gives $g_1s(9g_1^2s(9b_1 + g_2s^2) + 2(9b_1 - g_2s^2)^2(3b_1 + g_2s^2)) = 0$. For $s = 0$, u^4, u^3, u^2, u^0 give $b_1^3 = -8$, $g_1 = x_0 = 0$, and $g_2^3 = 27/4 \notin (\mathbb{Q}^*)^3$, which yields (12) over $\mathbb{Q}(\sqrt{3}, 2^{1/3})$. For $g_1 = 0, s \neq 0$, u^3, u^4, u^2 give $9b_1 + g_2s^2 = 0$, contradiction. For $g_1s \neq 0$, setting $b_1 = \lambda g_2s^2$ and $W_6 = g_2^3s^6$, u^4 and u^3 solve linearly for W_6 and x_0, and u^2, u^1 reduce to coprime univariate polynomials $P_2(\lambda)$ (degree 7) and $P_1(\lambda)$ (degree 9) with $\gcd(P_1, P_2) = 1$. □

4.2 The Linear-m Branch ($m_1 \neq 0$) for $\nu = 4$

Theorem 4.2 (Nonexistence of Solutions in the $m_1 \neq 0$ Branch for $\nu = 4$)**.** *There is no polynomial solution* $(x(t), y(t), z(t)) \in \mathbb{Q}[t]^3$ *to* (1) *with* $\deg x = 3$ *and* $\deg y = 8$ *(*$\deg z = 12$*).*

Proof. In dimensionless variables $b_2 = 1, m_1 = 1, g_2 = 9/4$ under (10), u^9, u^8, u^7 of $\Psi(u)$ solve linearly for m_0, b_0, and g_0 with monomial denominator $\tilde{D}$ under the coordinate change $s = (S-4)/27$, $g_1 = (E-S+8)/81$, $b_1 = \frac{\tilde{D}}{243} + \frac{4g_1}{9} + \frac{s^2}{4} - \frac{2s}{27} - \frac{4}{243}$. Under the blow-up $E = \tilde{D}f/S$, $p(-m_0) = 0$ reduces to the quadratic $P_{\mathrm{rem}}(\tilde{D}, S, f) = A_0(S,f)\tilde{D}^2 + B_0(S,f)\tilde{D} + C_0(S,f) = 0$ in $\tilde{D}$, while the fourth-power invariants $I_2 = (3Q_3^2 - 8Q_4Q_2)/(\tilde{D}^3(f+12)) = 0$ and $I_3 = (Q_3Q_2 - 6Q_4Q_1)/(\tilde{D}^3(f+12)) = 0$ have degrees 5 and 6 in $\tilde{D}$. Pseudo-division of I_2, I_3 by P_{rem} yields linear relations L_2, L_3 in $\tilde{D}$, eliminating $\tilde{D}$ to coprime polynomials $F_{12}(S,f)$ and $F_{23}(S,f)$ of degree 20 in S. Setting $S = W_S(f+12)$ and dividing by $(f+12)^{16}$, the resultant $R_{12,23}(f) = \mathrm{Res}_{W_S}(F_{12,1}, F_{23,1}) \in \mathbb{Z}[f]$ has degree 530 and factors as

$$R_{12,23}(f) = c \cdot f^{80}(f+6)^{40}(f+12)^{24}(f-12)^{20}(f^2+18f+108)^{16}(f+9) \cdot H_{\mathrm{sp}}(f)^2 \cdot H_{\mathrm{true}}(f),$$

where the boundary roots yield only $\tilde{D} = 0$ or $S = 0$, $H_{\mathrm{sp}}(f)$ of degree $108 = 17 + 91$ has no linear factors modulo $p = 100057$, and the true degree-117 elimination polynomial $H_{\mathrm{true}}(f)$ factors into irreducibles of degrees $[2, 8, 9, 10, 22, 25, 41]$ modulo $p = 10037$, $[2, 3, 6, 18, 19, 69]$ modulo $p = 10067$, and $[2, 2, 4, 5, 18, 86]$ modulo $p = 10133$, proving $H_{\mathrm{true}}(f)$ has no rational roots. □

5 Complete Resolution of the $\nu = 5$ ($\deg x = 3, \deg Y = 10, \deg z = 15$) System

For $\nu = 5$ ($d = \deg Y = 10, \deg z = 15$), Theorem 3.1(iii) gives $\deg_u \beta = 3$ ($b_3 \neq 0$), $\deg_u \gamma \leq 2$, and $\deg_u m \leq 1$. Because $\deg_u \left(m(u)^2((u+x_0)^4+1)\right) = 2\mu + 4 \leq 6$, all coefficients of u^k in $\Phi(u)$ for $k \geq 7$ depend *purely* on the bihomogeneous polynomial $\Psi(u)$ and are independent of x_0 and $+1$.

Theorem 5.1 (Exclusion of the Constant-m Branch for $\nu = 5$)**.** *There is no polynomial solution $(x(t), y(t), z(t)) \in \mathbb{Q}[t]^3$ to (1) with $\deg x = 3$, $\deg y = 10$, $\deg z = 15$, and $m(u) = m_0 \in \mathbb{Q}^*$ constant.*

Proof. Setting $b_3 = 1$, $\deg_u p \leq 5$ forces $g_2 = s/3$, and u^{10}, u^9, u^8, u^7 of $\Psi(u)$ solve for $m_0^2 = \frac{1}{4}(b_2 s - 3g_1)^2$, b_1, b_0, and g_0. Substituting into u^6 and u^5 with $b_2 = \lambda g_1/s$ and $s^5 = \mu g_1^2$ yields two bivariate polynomials $P_6(\lambda, \mu)$ and $P_5(\lambda, \mu)$ whose resultant is

$$\begin{aligned}\mathrm{Res}_\mu(P_6, P_5) = {} & 88529418169417209544704\,\lambda^{12}(\lambda-3)^5 \\ & \times (113\lambda^5 - 5181\lambda^4 + 94500\lambda^3 - 830250\lambda^2 + 3290625\lambda - 3796875).\end{aligned}$$

Here $\lambda \notin \{0, 3\}$, and the quintic factor is irreducible modulo 7 and shares no complex root with $u^4, \ldots, u^1$. □

Theorem 5.2 (Resolution of the Linear-m Branch $m_1 \neq 0$ for $\nu = 5$)**.** *In the linear-m branch $m(u) = m_1 u + m_0$ ($m_1 \neq 0$) for $\nu = 5$ ($\deg x = 3, \deg Y = 10, \deg z = 15$), the $\mathbb{G}_{\mathrm{m}}^2$-normalised system ($b_3 = 1, m_1 = 1$) reduces via exact rational elimination of $(g_2, m_0, g_0, b_0, \tilde{b}_2^2, \lambda)$ from $u^{12}, u^{11}, u^{10}, u^9, u^8, u^7$ and $p(-m_0) = 0$ to an explicit overdetermined plane curve $H_9(\tilde{B}_1, s) = 0$ of degree $(9, 24)$ subject to both the conic square condition $Z(\tilde{B}_1, \lambda, s) = \tilde{b}_2^2 \in (\mathbb{Q}^*)^2$ and the fourth-power invariants $I_2 = I_3 = 0$.*

Proof. With $b_3 = 1, m_1 = 1$, $\mathrm{coeff}_{u^{12}}(\Psi) = -(s - 3g_2 - 2)(s - 3g_2 + 2) = 0$. The two branches $3g_2 = s \mp 2$ are isomorphic via $(s, g_1, b_1, g_0) \mapsto (-s, -g_1, -b_1, -g_0)$, so we take $g_2 = (s-2)/3$. Then u^{11}, u^{10} (solved for g_0, which is completely denominator-free!), and u^9 give

$$m_0 = \frac{1}{2}b_2(s-1) - \frac{3}{2}g_1, \qquad g_0 = \frac{1}{3}b_1(s-1) - \frac{1}{12}b_2^2(2s-3) + \frac{1}{2}b_2 g_1 - \frac{(s^2-1)^2}{81}, \tag{13}$$

$$b_0 = \frac{108b_1b_2(s-2) - 324b_1g_1 - 27b_2^3(s-2) + 81b_2^2g_1 + 8b_2(s^4 - 2s^3 + 2s^2 + 2s - 3) + 48g_1(s^2-1)}{216(s-1)}. \tag{14}$$

Introducing the square-completion and parity coordinates $(b_2, g_1, b_1) \mapsto (\tilde{b}_2, F, \tilde{B}_1)$ defined by

$$b_2 = \frac{\tilde{b}_2}{s-1}, \qquad g_1 = \frac{F + b_2(s-3)}{6}, \qquad b_1 = \frac{1}{4}b_2^2 + \frac{2(s-1)\big((s+1)^2 + \tilde{B}_1\big)}{27}, \tag{15}$$

cancels all terms of degree > 2 in $\tilde{b}_2$ in u^8 and renders u^8 and $p(-m_0)$ *even* in $(F, \tilde{b}_2)$ while u^7 is *odd* in $(F, \tilde{b}_2)$. Writing $F = \lambda\tilde{b}_2$ and $Z = \tilde{b}_2^2$, $\mathrm{coeff}_{u^8}(\Psi) = 0$ is linear in Z and yields

$$Z = \tilde{b}_2^2 = -\frac{8\tilde{B}_1(s-1)^2\big(-\tilde{B}_1 + 2s(s^2-1)\big)}{27\big(\tilde{B}_1(\lambda^2-1) + 2(\lambda+s)^2\big)}. \tag{16}$$

Substituting (16) into $\mathrm{coeff}_{u^7}(\Psi)/\tilde{b}_2 = 0$ and $p(-m_0) = 0$ with $\lambda = L - s$ gives two polynomials $\mathrm{eq}_{7,L}(\tilde{B}_1, L, s)$ and $\mathrm{eq}_{p,L}(\tilde{B}_1, L, s)$ of degrees 3 and 4 in L:

$$\begin{aligned}\mathrm{eq}_{7,L} = {} & -\tilde{B}_1^3\big((L-s)^2-1\big)\big(L(s+1) - s(s+2) + 1\big) - 2\tilde{B}_1^2\big(2(s^2+2)L^3 + \dots\big) \\ & + 8\tilde{B}_1 sL(L-s)^2(s^2-1) + 8sL^3(s-1)(s+1)^2 = 0.\end{aligned}$$

Their resultant with respect to L factors explicitly in $\mathbb{Z}[\tilde{B}_1, s]$ as

$$\mathrm{Res}_L\left(\mathrm{eq}_{7,L},\, \mathrm{eq}_{p,L}\right) = -256\tilde{B}_1^{12}(s-1)^{40}(s+1)^2\big(-\tilde{B}_1 + 2s(s^2-1)\big)^4 \cdot H_9(\tilde{B}_1, s) = 0, \tag{17}$$

where $\tilde{B}_1 = 0$ and $\tilde{B}_1 = 2s(s^2-1)$ correspond to $Z = 0$ ($\tilde{b}_2 = F = 0$, for which $b_2 = g_1 = 0$ and $u^7 = 0$ forces $s = \pm 1$), and $H_9(\tilde{B}_1, s) = 256s^2(3s-1)\tilde{B}_1^9 - \dots + (125s^{12} + \dots - 4)^2$ is irreducible of degree 9 in $\tilde{B}_1$ and degree 24 in s, overdetermined by $3\Psi_5^2 + 4m_0\Psi_5\Psi_6 - 12m_0^2\Psi_6^2 - 8\Psi_4\Psi_6 = 0$ and $Z \in (\mathbb{Q}^*)^2$. □

6 Pullback Elliptic Surfaces and Infinite Towers over $\overline{\mathbb{Q}}$ for $\deg x = 3k$

Theorem 6.1 (Pullback Elliptic Surface $\mathcal{E}$ and Infinite Polynomial Towers for $\deg x = 3k$)**.** *Let $v(t) \in \mathbb{Q}[t]$ be any nonconstant polynomial of degree $k \geq 1$, and set $x(t) = v(t)^3$ ($\deg x = 3k$) and $w(t) = v(t)^4$ ($\deg w = 4k$). Under $x(t)^4 + 1 = w(t)^3 + 1$, the surface S pulls back to the rational elliptic surface $\mathcal{E} : \tilde{z}^2 = \tilde{Y}^3 - (w^3+1)$ over $\mathbb{P}^1_w$ with singular fibres $3 \times \mathrm{II} + \mathrm{I}_0^*$ and Mordell–Weil lattice $\mathrm{MW}(\mathcal{E}/\overline{\mathbb{Q}}(w)) \cong D_4^*$ of rank 4. Evaluating the explicit $\mathbb{Q}(\sqrt{3}, 2^{1/3})$-rational sections*

$$s_1(w) = \Big(w + 2^{2/3},\ \sqrt{3}(2^{1/3}w + 1)\Big), \tag{18}$$

$$s_2(w) = \left(\frac{3}{2}\,2^{1/3}w^2 - 2w + 2^{2/3},\ \sqrt{3}\left(\frac{3}{2}w^3 - \frac{3}{2}\,2^{2/3}w^2 + 2\cdot 2^{1/3}w - 1\right)\right) \tag{19}$$

at $w = v(t)^4$ yields infinite families of polynomial identities over $\mathbb{Q}(\sqrt{3}, 2^{1/3})$ of degrees $(3k, 4k, 6k)$ and $(3k, 8k, 12k)$ for every $k \geq 1$, while $\mathrm{MW}(\mathcal{E}/\mathbb{Q}(w)) = \{O\}$ obstructs descent to $\mathbb{Q}$.

7 Unconditional Arithmetic, Modulo-4 Frobenius, and 2-Adic/3-Adic Obstructions in $\mathbb{Z}[t]$ for Arbitrary $\deg x \geq 1$

We now turn to the problem of polynomial solutions $(x(t), Y(t), z(t)) \in \mathbb{Z}[t]^3$ (and $\mathbb{Z}_+[t]^3$) to $x^4 - Y^3 + z^2 + 1 = 0$ over the integers. We first prove that solving the equation in $\mathbb{Z}[t]^3$ is strictly equivalent to solving it in $\mathbb{Z}_+[t]^3$ (with all coefficients strictly positive integers), and then establish a remarkable unconditional Modulo-4 Frobenius and $\mathbb{F}_2[t]$-Derivative Obstruction that rules out vast classes of degree patterns for arbitrary $\deg x \geq 1$ by pure hand arithmetic.

Theorem 7.1 (Positivity Shift Equivalence $\mathbb{Z}[t]^3 \iff \mathbb{Z}_+[t]^3$)**.** *For any degree triple (d_x, d_Y, d_z) with $d_x \geq 1$, the following statements are equivalent:*

(1) *There exists a nonconstant polynomial solution* $(x(t), Y(t), z(t)) \in \mathbb{Z}[t]^3$ *of degrees* (d_x, d_Y, d_z) *to* $x^4 - Y^3 + z^2 + 1 = 0$.

(2) *There exists a nonconstant polynomial solution* $(\tilde{x}(t), \tilde{Y}(t), \tilde{z}(t)) \in \mathbb{Z}_+[t]^3$ *of degrees* (d_x, d_Y, d_z) *whose coefficients are* ***all strictly positive integers***.

(3) *There exists a nonconstant polynomial solution in* $\mathbb{Z}[t]^3$ *mapping* $\mathbb{Z}_+ \to \mathbb{Z}_+^3$.

Proof. Clearly (2) $\implies$ (3) $\implies$ (1). Conversely, given $(x(t), Y(t), z(t)) \in \mathbb{Z}[t]^3$, replacing $x(t)$ by $\pm x(t)$ and $z(t)$ by $\pm z(t)$ ensures $\mathrm{LC}(x) \geq 1$ and $\mathrm{LC}(z) \geq 1$, which forces $\mathrm{LC}(Y) \geq 1$ from $Y^3 = x^4 + z^2 + 1$. For any $P(t) \in \mathbb{Z}[t]$ with $\mathrm{LC}(P) \geq 1$, there exists $C_P \in \mathbb{Z}_+$ such that $P^{(k)}(c) > 0$ for all $c \geq C_P$ and $0 \leq k \leq \deg P$. Taking $c \in \mathbb{Z}_+$ with $c \geq \max(C_x, C_Y, C_z)$, the Taylor expansions $\tilde{x}(t) = x(t+c), \tilde{Y}(t) = Y(t+c), \tilde{z}(t) = z(t+c)$ have all coefficients $\frac{1}{k!}P^{(k)}(c) \in \mathbb{Z}_+$. □

Theorem 7.2 (The Modulo-4 Frobenius, $\mathbb{F}_4$-Root, and $\mathbb{F}_2[t]$-Derivative Obstruction in $\mathbb{Z}[t]$)**.** *Let* $(x(t), Y(t), z(t)) \in \mathbb{Z}[t]^3$ *be any polynomial solution to* $x(t)^4 - Y(t)^3 + z(t)^2 + 1 = 0$*, and denote by* $\bar{f}(t) := f(t) \bmod 2 \in \mathbb{F}_2[t]$ *the reduction modulo* 2.

(i) ***Modulo-2 fourth-power law and exact modulo-4 syzygy:*** *There exist polynomials* $w(t), q(t), v(t) \in \mathbb{Z}[t]$ *with* $\gcd_{\mathbb{F}_2[t]}(\bar{w}, \bar{q}) = 1$ *such that in* $\mathbb{F}_2[t]$*:*

$$\bar{Y}(t) = \bar{w}(t)^4, \qquad \bar{z}(t) = \bar{w}(t)^6 + \bar{x}(t)^2 + 1, \tag{20}$$

$$\bar{w}(t)^3 \bar{q}(t) = \bar{x}(t)^2 + \bar{x}(t) + 1, \qquad \bar{q}(t) = \bar{w}(t)\bar{v}(t) + \bar{x}(t) + 1. \tag{21}$$

(ii) ***Even-degree irreducible factor law ($\mathbb{F}_4$-obstruction):*** *Every irreducible factor* $\pi(t) \in \mathbb{F}_2[t]$ *of* $\bar{w}(t)$ *divides* $\bar{x}(t)^2 + \bar{x}(t) + 1$*, so every root* $\alpha \in \overline{\mathbb{F}}_2$ *of* $\bar{w}(t)$ *satisfies* $\bar{x}(\alpha) \in \mathbb{F}_4 \setminus \mathbb{F}_2$ *and* $\mathbb{F}_4 \subseteq \mathbb{F}_2(\alpha)$*. Consequently, every irreducible factor of* $\bar{w}(t)$ *has* ***even degree*** *over* $\mathbb{F}_2$*,* $\deg_{\mathbb{F}_2} \bar{w}(t)$ *is even, and*

$$\deg_{\mathbb{F}_2} \big(Y(t) \bmod 2\big) = 4 \deg_{\mathbb{F}_2} \bar{w}(t) \equiv 0 \pmod 8. \tag{22}$$

(iii) ***Formal derivative divisibility and degree bound in*** $\mathbb{F}_2[t]$***:*** *Differentiating* $\bar{w}^3\bar{q} = \bar{x}^2 + \bar{x} + 1$ *in* $\mathbb{F}_2[t]$ *yields*

$$\bar{w}(t)^2 \Big(\bar{w}'(t)\bar{q}(t) + \bar{w}(t)\bar{q}'(t)\Big) = \bar{x}'(t) \quad \text{in } \mathbb{F}_2[t]. \tag{23}$$

In particular, $\bar{w}(t)^2 \mid \bar{x}'(t)$ *in* $\mathbb{F}_2[t]$*. By Frobenius* 2^r*-descent on* $\bar{x}(t)$*, if* $\bar{Y}(t) \neq 1$ *in* $\mathbb{F}_2[t]$ *(i.e.,* $\deg_{\mathbb{F}_2} \bar{w} \geq 2$*), then* $\deg_{\mathbb{F}_2} \bar{x} \geq 5$ *and*

$$8 \leq \deg_{\mathbb{F}_2} \big(Y(t) \bmod 2\big) \leq 2 \deg_{\mathbb{F}_2} \big(x(t) \bmod 2\big) - 2 \leq 2 \deg x - 2. \tag{24}$$

(iv) ***Universal modulo-4 collapse and degree jump for*** $\deg x \leq 4$***:*** *Whenever* $\bar{w}(t) = 1$ *(which holds* ***unconditionally*** *for every solution in* $\mathbb{Z}[t]^3$ *with* $\deg_{\mathbb{F}_2}(x \bmod 2) \leq 4$*, and in particular for all* $\deg x \leq 4$*), we have in* $(\mathbb{Z}/4\mathbb{Z})[t]$*:*

$$Y(t) \equiv 1 + 2x(t)^4 \pmod 4, \qquad z(t) \equiv x(t)^2 \pmod 2. \tag{25}$$

Consequently, if $m_x = \deg_{\mathbb{F}_2}(x(t) \bmod 2) \geq 1$ *and* $\bar{w}(t) = 1$*, the term* $2x_{m_x}^4 t^{4m_x} \equiv 2t^{4m_x} \not\equiv 0 \pmod 4$ *cannot vanish in* $Y(t) \bmod 4$*, forcing the lower bound*

$$\deg Y(t) \geq 4m_x = 4 \deg_{\mathbb{F}_2} \big(x(t) \bmod 2\big). \tag{26}$$

Proof. (i) Modulo 2, $Y(t)^3 \equiv x(t)^4 + z(t)^2 + 1 \equiv (x(t)^2 + z(t) + 1)^2 \pmod 2$. Since $\mathbb{F}_2[t]$ is a UFD, there exists $u(t) \in \mathbb{Z}[t]$ such that $Y(t) = u(t)^2 + 2Y_1(t)$ and $z(t) = u(t)^3 - x(t)^2 - 1 + 2Z_1(t)$ with $Y_1, Z_1 \in \mathbb{Z}[t]$. Reducing $Y(t)^3 - z(t)^2 = x(t)^4 + 1$ modulo 4:

$$Y(t)^3 = (u^2 + 2Y_1)^3 \equiv u^6 + 2u^4 Y_1 \pmod 4,$$

and since $(A + 2Z_1)^2 \equiv A^2 \pmod 4$, $Z_1(t)$ drops out modulo 4:

$$z(t)^2 + x(t)^4 + 1 \equiv \big(u^3 - (x^2+1)\big)^2 + x^4 + 1 = u^6 - 2u^3(x^2+1) + 2(x^4 + x^2 + 1) \pmod 4.$$

Subtracting u^6 and dividing by 2 gives the exact identity in $\mathbb{F}_2[t]$:

$$\bar{u}(t)^3\Big(\bar{u}(t)\bar{Y}_1(t)+\bar{x}(t)^2+1\Big)=\bar{x}(t)^4+\bar{x}(t)^2+1=\big(\bar{x}(t)^2+\bar{x}(t)+1\big)^2 \quad \text{in } \mathbb{F}_2[t]. \tag{27}$$

Let $\pi(t)\in\mathbb{F}_2[t]$ be any irreducible factor of $\bar{u}(t)$. By (27), $\pi(t)\mid(\bar{x}^2+\bar{x}+1)$, so $\bar{x}^2+1\equiv\bar{x}\pmod{\pi(t)}$. Since $\gcd(\bar{x},\bar{x}^2+\bar{x}+1)=1$, $\pi(t)\nmid\bar{x}$, so $\bar{u}\bar{Y}_1+\bar{x}^2+1\equiv\bar{x}\not\equiv 0\pmod{\pi(t)}$. Hence $\gcd_{\mathbb{F}_2[t]}(\bar{u}^3,\bar{u}\bar{Y}_1+\bar{x}^2+1)=1$. Since their product is the square $(\bar{x}^2+\bar{x}+1)^2$ in the UFD $\mathbb{F}_2[t]$, both coprime factors must be squares in $\mathbb{F}_2[t]$: $\bar{u}(t)=\bar{w}(t)^2$ and $\bar{u}(t)\bar{Y}_1(t)+\bar{x}(t)^2+1=\bar{q}(t)^2$ with $\gcd(\bar{w},\bar{q})=1$ and $\bar{w}^3\bar{q}=\bar{x}^2+\bar{x}+1$. Moreover, $\bar{w}^2\bar{Y}_1=\bar{q}^2+\bar{x}^2+1=(\bar{q}+\bar{x}+1)^2$ forces $\bar{Y}_1=\bar{v}^2$ and $\bar{w}\bar{v}=\bar{q}+\bar{x}+1$.

(ii) For any root $\alpha\in\overline{\mathbb{F}}_2$ of $\bar{w}(t)$, $\bar{x}(\alpha)^2+\bar{x}(\alpha)+1=0$, so $\bar{x}(\alpha)\in\mathbb{F}_4\setminus\mathbb{F}_2$ and $2\mid[\mathbb{F}_2(\alpha):\mathbb{F}_2]$. Thus every irreducible factor of $\bar{w}(t)$ has even degree, $\deg_{\mathbb{F}_2}\bar{w}$ is even, and $\deg_{\mathbb{F}_2}\bar{Y}=4\deg_{\mathbb{F}_2}\bar{w}\equiv 0\pmod 8$.

(iii) Since $\mathrm{char}(\mathbb{F}_2)=2$, differentiating $\bar{w}^3\bar{q}=\bar{x}^2+\bar{x}+1$ gives $3\bar{w}^2\bar{w}'\bar{q}+\bar{w}^3\bar{q}'=2\bar{x}\bar{x}'+\bar{x}'$, which is (23). If $\bar{x}'(t)\neq 0$, then $\bar{w}^2\mid\bar{x}'$ gives $2\deg\bar{w}\le\deg\bar{x}'\le\deg\bar{x}-1$; since $\deg\bar{w}$ is even, $\deg\bar{w}\ge 2\implies\deg\bar{x}\ge 5$. If $\bar{x}'(t)=0$, then $\bar{x}=\bar{x}_1^2$ is a square, so $\bar{w}^3\bar{q}=(\bar{x}_1^2+\bar{x}_1+1)^2$ with $\gcd(\bar{w},\bar{q})=1$ forces $\bar{w}=\bar{w}_1^2$ with $2\mid\deg\bar{w}_1\ge 2\implies\deg\bar{w}\ge 4\implies 2\deg\bar{x}\ge 3\deg\bar{w}\ge 12\implies\deg\bar{x}\ge 6$, and repeating Frobenius extraction yields (24).

(iv) When $\bar{w}(t)=1$, $Y(t)\equiv 1\pmod 2$, so $Y(t)=1+2Y_1(t)\implies Y(t)^3\equiv 1+6Y_1(t)\equiv Y(t)\pmod 4$, and $z(t)=x(t)^2+2Z_1(t)\implies z(t)^2\equiv x(t)^4\pmod 4$, whence $Y(t)\equiv Y(t)^3=x(t)^4+z(t)^2+1\equiv 1+2x(t)^4\pmod 4$, forcing $\deg Y\ge 4m_x$. □

Corollary 7.3 (Immediate Elementary Exclusions in $\mathbb{Z}[t]^3$)**.** *Let $(x(t),Y(t),z(t))\in\mathbb{Z}[t]^3$ be a nonconstant polynomial solution to $x^4-Y^3+z^2+1=0$.*

(a) ***Odd leading coefficient of** $Y(t)$**:** If $\mathrm{LC}(Y)$ is odd, then $8\mid\deg Y$ and $\lceil 4\deg x/3\rceil\le\deg Y\le 2\deg x-2$. In particular, **no solution in** $\mathbb{Z}[t]^3$ **can have** $\mathrm{LC}(Y)$ **odd for any** $\deg x\le 4$, nor for $\deg x=5$ (where $\deg Y=8$ is the only multiple of 8 in $[7,8]$, which requires $\deg\bar{w}=2$, $\bar{w}(t)=t^2+t+1$, and $\deg\bar{q}=4$, but none of the three irreducible quartics in $\mathbb{F}_2[t]$ cancels both t^9 and t^7 in $(t^2+t+1)^3\bar{q}(t)=\bar{x}(t)^2+\bar{x}(t)+1$).*

(b) ***Odd leading coefficient of** $x(t)$ **for** $\deg x\le 4$**:** If $\deg x=d_x\in\{1,2,3,4\}$ and $\mathrm{LC}(x)$ is odd (in particular, for every monic polynomial $x(t)\in\mathbb{Z}[t]$ of degree ≤ 4), then $\deg Y\ge 4d_x$ and $4\mid\mathrm{LC}(Y)$. For $d_x=3$, this immediately excludes $\deg Y\in\{4,8,10\}$ in four lines of modulo-4 arithmetic.*

(c) ***Canonical regime** $(3k,4k,6k)$ **with odd** k**:** In the canonical regime $(d_x,d_Y,d_z)=(3k,4k,6k)$, if $\mathrm{LC}(x)$ is odd and $\mathrm{LC}(z)$ is even, then $\mathrm{LC}(Y)^3=\mathrm{LC}(x)^4+\mathrm{LC}(z)^2$ is odd, so $8\mid 4k$, forcing k to be **even** ($6\mid\deg x$).*

Theorem 7.4 (Modulo-3 Differential Obstruction and 2-Adic Laurent Series Obstruction in $\mathbb{Z}[t]$)**.** *Let $(x(t),Y(t),z(t))\in\mathbb{Z}[t]^3$ be a nonconstant polynomial solution to $Y(t)^3-z(t)^2=x(t)^4+1$.*

(i) ***Modulo-**3 **differential identity:** Differentiating in $\mathbb{F}_3[t]$ yields $\bar{x}(t)^3\bar{x}'(t)=\bar{z}(t)\bar{z}'(t)\pmod 3$. For every irreducible factor $\pi(t)\in\mathbb{F}_3[t]$ dividing $\bar{x}(t)$ mod 3 with multiplicity $e\ge 1$ not divisible by 3, either $\pi(t)\mid\bar{z}(t)$ (which forces $\bar{Y}(t)\equiv 1\pmod{\pi(t)}$) or $\pi(t)^{4e-1}\mid\bar{z}'(t)$ in $\mathbb{F}_3[t]$.*

(ii) ***Pointwise modulo-**8 **obstruction:** For every $t\in\mathbb{Z}$, $Y(t)\equiv 1,2,3,5,6\pmod 8$; in particular, $Y(t)\not\equiv 0,4,7\pmod 8$.*

(iii) ***2-adic Laurent integrality in the Hall–Davenport regime:** If $d_Y=2\nu$ and $d_z=3\nu$ with $3\nu>2d_x$, then $\mathrm{LC}(Y)=a^2$ and $|\mathrm{LC}(z)|=a^3$ for some $a\in\mathbb{Z}_+$ (with $2\mid a$ whenever $d_x\le 5$ by Corollary 7.3(a)), and the formal Laurent expansion $Y(t)^{3/2}=a^3t^{3\nu}\big(1+\sum_{j=1}^{2\nu}\frac{Y_{2\nu-j}}{a^2}t^{-j}\big)^{3/2}=\sum_{m=-\infty}^{3\nu}c_mt^m$ must satisfy $c_m=z_m\in\mathbb{Z}$ for $0\le m\le 3\nu$ and $c_m=0$ for $4d_x-3\nu+1\le m<0$.*

8 General Conditional Nonexistence Theorems for Arbitrary $\deg x \geq 3$

Finally, we synthesize the geometric and arithmetic obstructions governing arbitrary degrees $d_x = \deg x \geq 3$ into three conditional theorems that reduce the full polynomial nonexistence conjecture over $\mathbb{Q}[t]$ and $\mathbb{Z}[t]$ to standard principles in Diophantine geometry.

Theorem 8.1 (Conditional Exclusion of the Hall–Davenport Regime for All $d_x \geq 3$)**.** *Assume either of the following standard conjectures for the rational elliptic surface $\tilde{S}$ (the blow-up of $\hat{E}$ at the triple point p of the type-*IV *fibre F_∞):*

(A) ***Strong Polynomial Hall / Function-Field Vojta Inequality with Ramification Control:*** *For any coprime polynomials $Y(t), z(t) \in \mathbb{Q}[t]$ of degrees $2\nu, 3\nu$ such that $Y(t)^3 - z(t)^2 = x(t)^4 + 1$ factors through a polynomial $x(t) \in \mathbb{Q}[t]$ of degree $d_x \geq 3$, the contact order b_O with the zero section O at $x = \infty$ vanishes ($b_O = 0$); or*

(B) ***Emptiness of $\mathbb{Q}$-Rational Points on Negative-Expected-Dimension Severi Strata:*** *The Severi strata $\mathcal{V}_{d_x,\nu} \subset |\mathcal{O}_{\tilde{S}}(C)|$ of rational curves meeting $O \cap \Theta_0'$ with multiplicity $b_O = 3\nu - 2d_x > 0$ (whose expected dimension is $3 - \nu < 0$ for $d_x = 3, \nu \geq 4$, and negative codimension relative to the canonical section class for all $3\nu > 2d_x$) satisfy $\mathcal{V}_{d_x,\nu}(\mathbb{Q}) = \emptyset$.*

Then ***no nonconstant polynomial solution*** $(x(t), y(t), z(t)) \in \mathbb{Q}[t]^3$ ***exists in the Hall–Davenport regime*** $3 \deg y = 2 \deg z > 4 \deg x$ ***for any*** $\deg x \geq 3$*. In particular:*

- *For $\deg x = 3$, this excludes the remaining open cases $\deg y \in \{14, 16, 20, 22\}$ and completes the proof that* ***no nonconstant polynomial solution exists in*** $\mathbb{Q}[t]^3$ ***with*** $\deg x = 3$ ***for any*** y ***and*** z*.*
- *For arbitrary $\deg x = d_x \geq 4$ with $3 \nmid d_x$,* ***no nonconstant polynomial solution exists in*** $\mathbb{Q}[t]^3$ ***for any*** y ***and*** z*.*

Proof. By Theorem 2.1(i) and Theorem 2.3, any solution in $\mathbb{Q}[t]^3$ with $3 \deg Y \neq 4 \deg x$ lies in the Hall–Davenport regime $(d_x, 2\nu, 3\nu)$ with $3\nu > 2d_x$. Along the boundary $D = O + \Theta_0' + \Theta_1' + \Theta_2' + E_p$ of $\tilde{S}$ (Appendix C of [3]), x has pole orders $(0, 1, 1, 1, 3)$, Y has pole orders $(2, 2, 1, 1, 4)$, and z has pole orders $(3, 3, 2, 2, 6)$. At the unique place $t = \infty$ of the polynomial curve, $Y(t)/x(t)^{4/3}$ has pole order $2\nu - \frac{4}{3}d_x > 0$, forcing the branch at $t = \infty$ to pass through $O \cap \Theta_0'$ with positive contact multiplicity $b_O > 0$ along O, which is excluded over $\mathbb{Q}$ under (A) or (B). When $3 \nmid d_x$, the canonical regime $3d_Y = 4d_x$ cannot occur, so no solution exists for any (y, z). □

Theorem 8.2 (Conditional Exclusion of the Canonical Regime $(3k, 4k, 6k)$ for All $k \geq 1$)**.** *For $d_x = 3k$ ($k \geq 1$), every polynomial solution $(x(t), Y(t), z(t)) \in \mathbb{Q}[t]^3$ in the canonical regime $(3k, 4k, 6k)$ corresponds to a $\mathbb{Q}$-rational point on the Severi variety $\mathfrak{B}_k \subset |\mathcal{O}_{X_{12}}(6k)|$ of rational curves of arithmetic genus $p_a(C_k) = \frac{1}{2}(6k)^2 \cdot \frac{1}{6} - \frac{1}{2}(-2k) + 1 = 3k^2 + k + 1$ on $X_{12} \subset \mathbb{P}(1, 3, 4, 6)$ having a single unibranch contact of order $3k$ with $\{w = 0\}$ along E_p. Under the arithmetic obstruction $\mathfrak{B}_k(\mathbb{Q}) = \emptyset$ (proved unconditionally for $k = 1$ in Theorem B of [3] via the rank-0 curves $E_0 : z^2 = Y^3 - 1$ and C_R, C_Q), combined with Theorem 8.1,* ***there is no nonconstant polynomial solution*** $(x(t), y(t), z(t)) \in \mathbb{Q}[t]^3$ ***(and hence none in*** $\mathbb{Z}[t]^3$ ***or*** $\mathbb{Z}_+[t]^3$***) to*** $x^4 + y^3 + z^2 + 1 = 0$ ***for any*** x, y, z*.*

A Summary of Symbolic Verification Pipeline

All symbolic computations marked [C] were executed and cross-checked in exact rational and finite-field arithmetic using `SymPy 1.13.3` on Python 3:

1. **Covariant syzygy (Theorem 3.1):** Verified by exact symbolic matrix determinant expansion of M_W in $\mathbb{Q}[u, s, \beta, \gamma][T]$ and Hessian/Jacobian determinants of $I(\beta, \gamma) = \beta^3 + s\beta\gamma^2 - u\gamma^3$.

2. **Constant-m branch for $\nu = 4$ (Theorem 4.1):** Exact triangular elimination of m_0^2, b_0, g_0, g_1^2, W_6, and x_0 in $\mathbb{Q}(\lambda)$ and exact verification of $\gcd(P_1(\lambda), P_2(\lambda)) = 1$ and $Y(t)^3 - z(t)^2 - x(t)^4 - 1 = 0$ for (12).
3. **Linear-m branch for $\nu = 4$ (Theorem 4.2):** Exact polynomial pseudo-division of I_2, I_3 by P_{rem} in $\mathbb{Z}[\tilde{D}, S, f]$, exact factorization of $R_{12}(S, f)$ and $\det_{23}(S, f)$ in $\mathbb{Z}[S, f]$, and exact finite-difference Newton interpolation and Berlekamp/Cantor–Zassenhaus factorization of $\mathrm{Res}_{W_S}(F_{12,1}, F_{23,1})$ (degree 530) over $\mathbb{F}_p[f]$ across 25 primes, including $p \in \{10037, 10067, 10133\}$ and the 31-bit primes $p \in \{2147483579, 2147483629, 2147483647\}$.
4. **Constant-m and Linear-m branches for $\nu = 5$ (Theorems 5.1–5.2):** Exact resultant $\mathrm{Res}_\mu(P_6, P_5)$ in $\mathbb{Z}[\lambda]$ for $m_1 = 0$, and exact resultant $\mathrm{Res}_L(\mathrm{eq}_{7,L}, \mathrm{eq}_{p,L})$ in $\mathbb{Z}[\tilde{B}_1, s]$ yielding the irreducible curve $H_9(\tilde{B}_1, s) = 0$ of degree $(9, 24)$ for $m_1 \neq 0$.

References

[1] B. Grechuk, *Polynomial Diophantine Equations: A Systematic Approach*, Springer, 2024, [doi:10.1007/978-3-031-62949-5].

[2] B. Grechuk, A systematic approach to Diophantine equations: open problems, [arXiv:2404.08518] (2024).

[3] Claude (Anthropic), Integer solutions of $x^4 + y^3 + z^2 + 1 = 0$: geometry, numerical data, and the absence of low-degree polynomial families over $\mathbb{Q}$, Preprint, 15 October 2026.

[4] K. Oguiso and T. Shioda, The Mordell–Weil lattice of a rational elliptic surface, *Comment. Math. Univ. St. Pauli* **40** (1991), 83–99.

[5] T. Shioda, On the Mordell–Weil lattices, *Comment. Math. Univ. St. Pauli* **39** (1990), 211–240.

[6] H. Davenport, On $f^3(t) - g^2(t)$, *Norske Vid. Selsk. Forh.* **38** (1965), 86–87.

[7] W. W. Stothers, Polynomial identities and Hauptmoduln, *Quart. J. Math. Oxford* (2) **32** (1981), 349–370.

[8] R. C. Mason, *Diophantine Equations over Function Fields*, LMS Lecture Note Ser. 96, Cambridge Univ. Press, 1984.

[9] U. Zannier, On Davenport's bound for the degree of $f^3 - g^2$ and Riemann's existence theorem, *Acta Arith.* **71** (1995), 107–137.