



Whole-task peak correlation versus Eco Position (primary analysis)











One point per participant; labels show participant IDs. Each point uses the same eligible attempts for Eco and peak correlation.
Spearman 95% CIs: exploratory participant percentile bootstrap, unadjusted. p: permutation; q: BH; Pearson r/fit: descriptive.
CI unreliable: <97.5% defined bootstrap draws. BH family spans phases and features; phase scores are whole-attempt.
Eco uses the supplied column values directly; attempts with missing Eco values are excluded.
Axes shared within each L/R phase row; ranges can vary between rows. Whole-task panels share axes.

Task A: valid-phase peak correlation versus Eco Position (primary analysis)


Task A: Reach Ring / Left PSM
Pearson r = +0.14; Spearman ρ = +0.27
Spearman 95% CI: [-0.40, +0.64]
p = 0.417; BH q = 0.942; n = 11 participants




Task A: Reach Ring / Right PSM
Pearson r = +0.36; Spearman ρ = +0.31
Spearman 95% CI: [-0.42, +0.78]
p = 0.36; BH q = 0.942; n = 11 participants




Task A: Grab Ring / Left PSM
Pearson r = +0.19; Spearman ρ = +0.35
Spearman 95% CI: [-0.38, +0.83]
p = 0.301; BH q = 0.929; n = 11 participants




Task A: Grab Ring / Right PSM
Pearson r = +0.41; Spearman ρ = +0.42
Spearman 95% CI: [-0.24, +0.83]
p = 0.208; BH q = 0.816; n = 11 participants




Task A: Transfer Ring / Left PSM
Pearson r = +0.38; Spearman ρ = +0.42
Spearman 95% CI: [-0.25, +0.87]
p = 0.204; BH q = 0.816; n = 11 participants




Task A: Transfer Ring / Right PSM
Pearson r = +0.54; Spearman ρ = +0.55
Spearman 95% CI: [-0.01, +0.87]
p = 0.0842; BH q = 0.758; n = 11 participants



Task A: Place Ring / Left PSM
Pearson r = +0.11; Spearman ρ = -0.19
Spearman 95% CI: [-0.77, +0.69]
p = 0.576; BH q = 0.942; n = 11 participants





Task A: Place Ring / Right PSM
Pearson r = +0.16; Spearman ρ = +0.26
Spearman 95% CI: [-0.47, +0.73]
p = 0.433; BH q = 0.942; n = 11 participants





One point per participant; labels show participant IDs. Each point uses the same eligible attempts for Eco and peak correlation.
Spearman 95% CIs: exploratory participant percentile bootstrap, unadjusted. p: permutation; q: BH; Pearson r/fit: descriptive.
CI unreliable: <97.5% defined bootstrap draws. BH family spans phases and features; phase scores are whole-attempt.
Eco uses the supplied column values directly; attempts with missing Eco values are excluded.
Axes shared within each L/R phase row; ranges can vary between rows. Whole-task panels share axes.


Task C: valid-phase peak correlation versus Eco Position (primary analysis)














One point per participant; labels show participant IDs. Each point uses the same eligible attempts for Eco and peak correlation.
Spearman 95% CIs: exploratory participant percentile bootstrap, unadjusted. p: permutation; q: BH; Pearson r/fit: descriptive.
CI unreliable: <97.5% defined bootstrap draws. BH family spans phases and features; phase scores are whole-attempt.
Eco uses the supplied column values directly; attempts with missing Eco values are excluded.
Axes shared within each L/R phase row; ranges can vary between rows. Whole-task panels share axes.