
EcoRaw vs selected-PSM stage features (position)
* BH q < 0.05 within outcome/scenario/method; all three predictors and stages
Between participants
Within participant
Within participant
adjusted for task order + attempt
A: Reach Ring (selected PSM)
A: Grab Ring (selected PSM)
A: Transfer Ring (selected PSM)
A: Place Ring (selected PSM)
A: Drop Ring (selected PSM)
C: Reach String (selected PSM)
C: Threading (selected PSM)
C: Wrap around PSM (selected PSM)
C: Pull/cinch (selected PSM)
+0.40 -0.28 -0.41
-0.12 -0.33 -0.16
-0.20 -0.22 +0.00
-0.26 -0.16 -0.35
-0.02 -0.02 -0.18
+0.32 -0.10 -0.28
+0.50 +0.50 -0.65
-0.35 -0.34 -0.13
+0.05 +0.00 -0.15
-0.11 -0.14 -0.33
+0.18 +0.02 +0.17
+0.03 -0.00 +0.20
excluded excluded excluded
excluded excluded excluded
excluded excluded excluded
-0.13 +0.17 +0.52
-0.00 -0.42 -0.26
+0.01 -0.45 -0.36
low N low N low N
low N low N low N
low N low N low N
-0.18 +0.33 -0.53
+0.17 -0.33 -0.26
+0.15 -0.21 -0.21
excluded excluded excluded
excluded excluded excluded
excluded excluded excluded
Signed lag
Absolute lag
Peak r
Rank association
1.00
0.75
0.50
0.25
0.00
−0.25
−0.50
−0.75
−1.00
One PSM chosen per recording/stage by largest signed xcorr peak, before score matching.
Positive signed lag = gaze leads. Scores describe whole attempts. Missing outcomes are excluded.
Selected PSM indicates stronger correlation, not necessarily the operated instrument.